

Ozone is bad for health – but only for some?

Claudia Spix

In recent years there has been accumulating evidence that high levels of ozone exacerbate pre-existing respiratory disease and cause increases in emergency attendances, admissions to hospital, and mortality.¹⁻¹² In this issue of *Thorax* Stedman *et al* present a first attempt to derive from such studies an estimate of the number of additional hospital admissions attributable to ozone effects, given the distribution of respiratory admissions and ozone levels in Great Britain in the summers of 1993 and 1995.¹³ They present an interpolated estimate of each day's spatial ozone concentration distribution in Great Britain, use an average number of respiratory admissions per small region per day, and use a published respiratory hospital admissions/ozone dose response curve from a study based on London data to predict the number of extra cases per day and region caused by ozone values above a chosen cut off point. They then present the reader with the sum of those extra cases. This approach to the public health problems caused by ozone is innovative and, in principle, is important. However, the estimates produced are dependent on the accuracy of the predicted exposure estimate, which is difficult to assess from the methodological information given by the authors, and the use of a dose response curve from a slightly different time and place (London 1987-91) may also be a problem, though in fact effect estimates tend to be similar between places at least in Europe.¹² Alternatively, it might be more appropriate to use a coefficient from a model with a threshold assumption to predict effects above this threshold instead of a model with no threshold assumption. Whether this would improve the procedure needs to be further investigated. Finally, in the future one might want to see confidence intervals for those estimates of additional cases that take into account the error of both the spatial interpolation (exposure prediction) and the temporal effect prediction.

Apart from these purely methodological considerations, two medical questions come to mind. (1) Why can it not be said, for sure, how many additional cases there are? After all, the paper gives estimates between 317 and 11 195 additional admissions for the summer of 1995. (2) Why is the figure the authors give as the main result (+0.1% in 1993 and +0.35% in 1995) so low when there is so much evidence of the dangers of air pollution in general and ozone in particular?

Health care planners would certainly like to be able to put a price tag on certain levels of ozone. Statisticians and epidemiologists would very much like to do this too, so let me try to explain why this is so difficult and why it is consequently so important that Stedman *et al* do present an attempt to put this on a sound scientific basis – something rarely done before.^{14,15} The first question is related to the problem of thresholds. The authors present estimates for various choices of cut off points on a monotonic dose response curve. Further research – and not only epidemiological research – is needed to identify which of those values does come close to a medically justified effect limit. Attempts to find such a threshold have rarely been made with this type of data, and some may not have been published due to lack of success in identifying a threshold. Any such threshold is likely to be at the lower end of the ozone value distribution and this is the region where the

signal to noise ratio in such data sets tends to become unfavourable. A threshold found may really be an “effect detection limit”, which is dependent mostly on the size of the data set. The existing evidence is inconclusive to say the least.^{6,16} Thus, the question of determining a threshold even for small homogenous risk groups is currently open and a topic for further toxicological and epidemiological research. Consequently, the researchers presented estimates for a choice of possible thresholds.

The second question relates to the specific type of study results used here. These are not “ozone effects on respiratory health” but are a subset of all possible “short term ozone effects on respiratory health”. Probably few – maybe none – of the persons admitted to hospital on or following a high ozone day moved from a perfectly healthy respiratory tract to needing stationary treatment for a respiratory condition in one day. Practically all cases would be exacerbations of an existing chronic problem. For some patients this might mean entering a worse stage of their disease. It is likely that most chronic cases would react to an irritant by increasing their medication or reducing their activity, some of them thus causing work or school absences, while only a small fraction of the most sensitive ones would actually end up in hospital.¹⁷ This might explain why the effects seen on hospital admissions are, indeed, fairly small – usually 2-5% per 25 ppb.¹² The effect estimates for all summer in this study are further diminished by the fact that only a small fraction of the summer days exceeded any one of the thresholds examined (except for “zero”).

However, it is likely that short term effects are only part of the profile of the effects of ozone. There are a few studies that link lifetime exposure to ozone to the development of chronic disease, but the exposure assessment is necessarily very difficult.¹⁸⁻²⁰ Some studies and experiments point out that individuals and populations seem to be able to adapt to higher levels of ozone in terms of short term effects on lung function parameters, while in animal experiments long term exposure is seen to cause lasting damage to the lung tissue.²¹⁻²⁴ Without being able to quantify this it must be assumed that a certain fraction of the cases exacerbated by an ozone episode would not – or not yet – have been suffering from that condition at that time had the person not been repeatedly exposed to increased levels of ozone previously. In terms of costs, several years of additional treatment (and possibly work absences or early retirement) are perhaps more relevant than some days with a few additional patients in hospital.

More research on the quantification of disease and costs caused by air pollution must be greatly encouraged. The paper by Stedman *et al* may not be the ultimate solution as to how to conduct such studies methodologically, but it is certainly a good starting point for the necessary discussion on these matters.

GSF National Research Center for Environment and Health, Institute for Epidemiology, Neuherberg, 85758 Oberschleissheim, Germany

CLAUDIA SPIX

1 Lippmann M. Health effects of tropospheric ozone: review of recent research findings and their implications to ambient air quality standards. *J Expo Anal Environ Epidemiol* 1993;3:103-29.

- 2 Brunekreef B, Dockery DW, Krzyzanowski M. Epidemiologic studies on short term effects of low levels of major ambient air pollution components. *Environ Health Perspect* 1995;103(Suppl):3–13.
- 3 Schwartz J. Short term fluctuations in air pollution and hospital admissions of the elderly for respiratory disease. *Thorax* 1995;50:531–8.
- 4 Koenig JQ. Effect of ozone on respiratory responses in subjects with asthma. *Environ Health Perspect* 1995;103(Suppl):103–5.
- 5 Schwartz J. Air pollution and hospital admissions for respiratory disease. *Epidemiology* 1996;7:20–8.
- 6 Ponce de Leon A, Anderson A, Bland JM, Strachan DP, Bower J. Effects of air pollution on daily hospital admissions for respiratory disease in London between 1987/88 and 1991/92. *J Epidemiol Community Health* 1996;50(Suppl):63–70.
- 7 Quenel P, Zmirou D, Le Tertre A, Balducci F, Medina S, Barumandzadeh T, et al. Impact de la pollution atmosphérique urbaine de type acido-particulaire sur la mortalité quotidienne à Lyon et dans l'agglomération Parisienne. *Santé Publ* 1996;6:1–14.
- 8 Anderson HR, Spix C, Medina S, Schouten JP, Castellsague J, Rossi G, et al. Air pollution and daily admissions for chronic obstructive pulmonary disease in 6 European cities: results from the APHEA project. *Eur Respir J* 1997;10:1064–71.
- 9 Yang W, Jennison BL, Omaye ST. Air pollution and asthma emergency room visits in Reno, Nevada. *Inhalation Toxicol* 1997;9:15–29.
- 10 Borja-Aburto VH, Loomis DP, Bangdiwala SI, Shy CM, Rascon Pacheco RA. Ozone, suspended particulates and daily mortality in Mexico City. *Am J Epidemiol* 1997;145:258–68.
- 11 Burnett RT, Brook JR, Yung WT, Dales RE, Krewski D. Association between ozone and hospitalization for respiratory diseases in 16 Canadian cities. *Environ Res* 1997;72:24–31.
- 12 Spix C, Anderson HR, Schwartz J, Vigotti MA, Le Tertre A, Vonk J, et al. Short term effects of air pollution on hospital admissions of respiratory diseases in Europe – a quantitative summary of APHEA study results. *Arch Environ Health* 1997 (in press).
- 13 Stedman JR, Anderson HR, Atkinson RW, Maynard RL. Emergency hospital admissions for respiratory disorders attributable to summer time ozone episodes in Great Britain. *Thorax* 1997;52:958–63.
- 14 Heinz I, et al. Krankheitskosten durch Luftverschmutzung, Studie im Auftrag des Bundesministeriums für Umwelt, Naturschutz und Reaktorsicherheit. *Wirtschafts Beiträge* 1990:28.
- 15 ISPM Institut für Sozial- und Präventivmedizin der Universität Basel. Monetarisierung der verkehrsbedingten externen Gesundheitskosten, Teilbericht Epidemiologie. Studie im Auftrag des Dienstes für Gesamtverkehrstragen des Eidgenössischen Verkehrs- und Energiewirtschaftsdepartements, Basel, 1996.
- 16 Schwartz J. Air pollution and hospital admissions for the elderly in Detroit, Michigan. *Am J Respir Crit Care Med* 1994;150:648–55.
- 17 Franke K, Boeriu A, Degens P, et al. A 3-year cohort study on short term effects of air pollution in Germany. 1. Influences of medication and season. *The Science of the Total Environment* 1992;127:69–78.
- 18 Ackermann-Lieblich U, Leuenberger P, Schwartz J, Schindler C, Monn C, Bolognini C, et al. Lung function and long term exposure to air pollutants in Switzerland. *Am J Respir Crit Care Med* 1997;155:122–9.
- 19 Abbey DE, Lebowitz MD, Mills PK, Petersen FF, Beeson WL, Burchette RJ. Long term ambient concentrations of particulates and oxidants and development of chronic disease in a cohort of nonsmoking California residents. *Inhalation Toxicol* 1995;7:19–34.
- 20 Kunzli N, Lurmann F, Segal M, Ngo L, Balmes J, Tager IB. Reliability of lifetime residential history and activity measures as elements of cumulative ambient ozone exposure assessment. *J Expos Anal Environ Epidemiol* 1996; 6:289–310.
- 21 Linn WS, Avol EL, Shamoo DA, Peng RC, Valencia LM, Little DE, et al. Repeated laboratory ozone exposure of volunteer Los Angeles residents: an apparent seasonal variation in response. *Toxicol Ind Health* 1988;4: 505–20.
- 22 Stockstill BL, Chang LY, Menache MG, Mellick PW, Mercer RR, Crapo JD. Bronchiolarized metaplasia and interstitial fibrosis in rat lungs chronically exposed to high ambient levels of ozone. *Toxicol Appl Pharmacol* 1995; 134:251–63.
- 23 Drechsler-Parks DM, Bedi JF, Horvath SM. Pulmonary function responses of older men and women to ozone exposure. *Environ Gerontol* 1987;22: 91–101.
- 24 Hackney JD, Linn WS, Shamoo DA, Avol EL. Responses of selected reactive and nonreactive volunteers to ozone exposure in high and low pollution seasons. In: Schneider T, Lee SD, Wolters GJR, Grant LD, et al, eds. *Atmospheric ozone research and its policy implications*. Amsterdam: Elsevier, 1989:311–8.