26. **Bergeron A**, Soler P, Kambouchner M, *et al*. Cytokine profiles in idiopathic pulmonary fibrosis suggest an important role for TGF-beta and IL-10. *Eur Respir J* 2003;**22**:69–76.
27. **Carre P**, Leophonte P. [Cytokines and pulmonary fibrosis]. *Rev Mal Respir* 1993;**10**:193–207.
28. **Hodge G**, Hodge S, Chambers D, *et al*. Acute lung transplant rejection is associated with localized increase in T-cell IFNgamma and TNFalpha proinflammatory cytokines in the airways. *Transplantation* 2007;**84**:1452–8.
29. **Bates RC**, Mercurio AM. Tumor necrosis factor-alpha stimulates the epithelial-to-mesenchymal transition of human colonic organoids. *Mol Biol Cell* 2003;**14**:1790–800.
30. **Armendariz-Borunda J**, Katayama K, Seyer JM. Transcriptional mechanisms of type I collagen gene expression are differentially regulated by interleukin-1 beta, tumor necrosis factor alpha, and transforming growth factor beta in Ito cells. *J Biol Chem* 1992;**267**:14316–21.
31. **Regan MC**, Kirk SJ, Hurson M, *et al*. Tumor necrosis factor-alpha inhibits in vivo collagen synthesis. *Surgery* 1993;**113**:173–7.
32. **Solis-Herruzo JA**, Brenner DA, Chojkier M. Tumor necrosis factor alpha inhibits collagen gene transcription and collagen synthesis in cultured human fibroblasts. *J Biol Chem* 1988;**263**:5841–5.

Pulmonary puzzle

ANSWER

From the question on page 756

The history of the farmer's eating habits and diet had to be taken into consideration and questioned. It was found that he had a history of eating raw meat, with frequent consumption of uncooked dog and cow liver. The level of eosinophil cationic protein was 12.50 µg/l and the bronchoalveolar lavage fluid contained 20.8% monocytes, 4.6% neutrophils, 69.4% lymphocytes and 5.2% eosinophils. An open lung biopsy revealed extensive eosinophilic infiltration and a larval nematode near the pleural surface (fig 1). An ELISA with the larval excretory–secretory antigen of *Toxocara canis* was positive. He was treated with albendazole at 13 mg/kg/day (800 mg/day) for 6 weeks. After 6 months, he had improved chest radiographs and no respiratory symptoms.

Human infection by the nematode *T canis* occurs by ingesting embryonated eggs from the environment or encapsulated larvae from uncooked animal liver.¹ In 1952, Beaver *et al* identified larvae of *T canis* in a liver biopsy specimen and proposed the term visceral larva migrans (VLM).² The larvae penetrate the intestinal wall and begin a migration through the tissues. VLM involves several organ systems, such as the liver, lung, skin and, rarely, the central nervous system.³ Since *T canis* larvae are rarely detected by pathological examination, the most reliable diagnostic method is ELISA with excretory–secretory antigens.⁴ We believe that this is the first reported case of toxocariasis by direct detection of a *Toxocara* larva in an adult lung biopsy specimen. Toxocariasis should be considered as a differential diagnosis for adult patients with hypereosinophilic syndrome and a history of eating raw foods.

Figure 1 The subpleural granuloma associated with chronic organising pneumonia and a multifocal eosinophilic microabscess reveals *Toxocara* larva (inset) with a smooth and thin cuticle (scale bar: 50 µm).

Thorax 2009;**64**:777. doi:10.1136/thx.2008.101352a

REFERENCES

1. **Aragane K**, Akao N, Matsuyama T, *et al*. Fever, cough, and nodules on ankles. *Lancet* 1999;**354**:1872.
2. **Beaver PC**, Snyder CH, Carrera GM, *et al*. Chronic eosinophilia due to visceral larva migrans; report of three cases. *Pediatrics* 1952;**9**:7–19.
3. **Despommier D**. Toxocariasis: clinical aspects, epidemiology, medical ecology, and molecular aspects. *Clin Microbiol Rev* 2003;**16**:265–72.
4. **Jacquier P**, Gottstein B, Stingelin Y, *et al*. Immunodiagnosis of toxocarosis in humans: evaluation of a new enzyme-linked immunosorbent assay kit. *J Clin Microbiol* 1991;**29**:1831–5.