

and when NPPV should be applied in patients with stable hypercapnic COPD. Recent research indicates that the technique of how NPPV is applied is the crucial issue regarding the acceptance and effectiveness of NPPV. Here, the new concept of high-intensity NPPV has provided promising results with respect to gas exchange, lung function, treatment compliance, dyspnoea and HRQL. In addition, NPPV applied during physical activity and rehabilitation has emerged as a new potential indication for NPPV aiming to improve gas exchange, functional status and exercise-induced dyspnoea. Although the importance of proven NPPV-associated improvements in gas exchange, symptoms, functional status and HRQL is undisputed, we need an answer as soon as possible to the question: 'Are these new approaches also capable of improving long-term survival?'

Competing interest BS is sponsored by Resmed and Heinen & Löwenstein to give talks on mechanical ventilation.

Provenance and peer review Commissioned; not externally peer reviewed.

Thorax 2010;65:765–767.
doi:10.1136/thx.2010.145276

REFERENCES

- Elliott MW, Simonds AK, Carroll MP, et al. Domiciliary nocturnal nasal intermittent positive pressure ventilation in hypercapnic respiratory failure
- Wijkstra PJ, Lacasse Y, Guyatt GH, et al. A meta-analysis of nocturnal noninvasive positive pressure ventilation in patients with stable COPD. *Chest* 2003;124:337–43.
- Simonds AK, Elliott MW. Outcome of domiciliary nasal intermittent positive pressure ventilation in restrictive and obstructive disorders. *Thorax* 1995;50:604–9.
- Casanova C, Celli BR, Tost L, et al. Long-term controlled trial of nocturnal nasal positive pressure ventilation in patients with severe COPD. *Chest* 2000;118:1582–90.
- Clini E, Sturani C, Rossi A, et al. The Italian multicentre study on noninvasive ventilation in chronic obstructive pulmonary disease patients. *Eur Respir J* 2002;20:529–38.
- McEvoy RD, Pierce RJ, Hillman D, et al. Nocturnal non-invasive nasal ventilation in stable hypercapnic COPD: a randomised controlled trial. *Thorax* 2009;64:561–6.
- Ware JE Jr, Sherbourne CD. The MOS 36-item Short-Form Health Survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;30:473–83.
- Jones PW, Quirk FH, Baveystock CM. The St George's Respiratory Questionnaire. *Respir Med* 1991;85(Suppl B):25–31.
- Testa MA, Simonson DC. Assessment of quality-of-life outcomes. *N Engl J Med* 1996;334:835–40.
- Windisch W, Freidel K, Schucher B, et al. The Severe Respiratory Insufficiency (SRI) Questionnaire: a specific measure of health-related quality of life in patients receiving home mechanical ventilation. *J Clin Epidemiol* 2003;56:752–9.
- Windisch W, Budweiser S, Heinemann F, et al. The Severe Respiratory Insufficiency Questionnaire was valid for COPD patients with severe chronic respiratory failure. *J Clin Epidemiol* 2008;61:848–53.
- Windisch W, Vogel M, Sorichter S, et al. Normocapnia during nNPPV in chronic hypercapnic COPD reduces subsequent spontaneous PaCO₂. *Respir Med* 2002;96:572–9.
- Windisch W, Storre JH, Sorichter S, et al. Comparison of volume- and pressure-limited NPPV at night: a prospective randomized cross-over trial. *Respir Med* 2005;99:52–9.
- Dreher M, Storre JH, Schmoor C, et al. High-intensity versus low-intensity non-invasive ventilation in patients with stable hypercapnic COPD: a randomised crossover trial. *Thorax* 2010;65:303–8.
- Dreher M, Storre JH, Windisch W. Noninvasive ventilation during walking in patients with severe COPD: a randomised crossover trial. *Eur Respir J* 2007;29:930–6.
- Duiverman ML, Wempe JB, Bladde G, et al. Nocturnal non-invasive ventilation in addition to rehabilitation in hypercapnic patients with COPD. *Thorax* 2008;63:1052–7.

Lung Alerts: promoting education and encouraging new authors

Angshu Bhowmik,¹ Jenni Quint²

Lung Alerts were introduced by Wisia Wedzicha, Editor-in-Chief in 2003 to increase the educational content of the journal. The aim was to commission brief summaries of papers on respiratory topics published in non-respiratory journals, thereby attracting a broader interest particularly among doctors in training. It

was also intended to allow readers to keep abreast of publications in journals to which Respiratory Specialists are less likely to subscribe.

At the time of writing, 372 Lung Alerts have so far been published from 2003 until June 2010 at an average of 4.25 per month. A further 15 are in press.

Of the 387 Lung Alerts we have edited, looking at contributors from a regional perspective, the largest proportion (44%) were from London and the Kent, Surrey and Sussex Deaneries. 8% of contributors have been from outside the UK. The North Western (6%), South West Peninsula (5%), Eastern (5%) and LNR (Leices-

¹Homerton Hospital, London, UK; ²Department of Academic Respiratory Medicine, Royal Free and University College Medical School, Royal Free Hospital, London, UK

Correspondence to Angshu Bhowmik, Homerton Hospital, Homerton Row, London E9 6SR, UK; a.bhowmik@qmul.ac.uk

tershire, Northamptonshire and Rutland) (4%) Deaneries have been the other major sources of authors of Lung Alerts.

Fifty per cent have been written by Specialist Registrars or equivalent, 20% by doctors in various grades which previously fell under the category of 'Senior House Officer', 10% by 'Research Fellows', 7.5% by Consultant or equivalent doctors, 1.6% by Foundation 1 doctors and three of the Alerts have been written by students. The remainder were written by doctors of various other grades including Speciality doctors and Associate Specialists. We are hopeful that the experience of being able to publish in *Thorax* will have led to many of those who were not already committed to careers in Respiratory Medicine and Science being drawn to this speciality!

Over the years, 70 journals have provided the sources for the papers summarised in Lung Alerts. The largest number has been from the *New England Journal of Medicine* (99) with other major contributors being the *Lancet* (48) and

JAMA (24). The *BMJ*, with 19 papers, has also provided us with much material of interest.

One hundred and sixty-four of the Lung Alerts were about treatments and management strategies of various lung diseases. Ninety-four concerned pathological mechanisms of disease, with an increasing number in the last 2 years about genetic factors relating to lung disease, reflecting the increasing interest and knowledge about genetics in the published literature. We are confident that this trend will continue and we will gain a greater understanding of the genetics of pulmonary disease in the coming years.

The diseases of greatest interest in the general medical journals seem to be asthma and allergy, with 19% of the Lung Alerts. This was followed closely by lung cancer (16%), infection (excluding tuberculosis (TB)) (15%) and chronic obstructive pulmonary disease (COPD) (12%). TB garnered 7% of the alerts, with cystic fibrosis following at 4%.

Some of the Lung Alerts which we published are memorable to us for particular points of interest. For example, we learned that allergic sensitisation to

peanuts may occur through inflamed skin rather than only the oral route.¹ Other Alerts have reported early studies which have led to greater understanding of new procedures or treatments—for example, radio-frequency ablation^{2 3} and gefitinib⁴ in the treatment of lung cancer. We alerted our readers to a useful new method of assessing patients with COPD: the BODE index described in the *New England Journal of Medicine* in 2004.⁵ We have reported studies that reinforce good practice—for example, PET-CT to reduce futile thoracotomies⁶ and the use of large particle talc for safe pleurodesis.⁷ We have tried to incorporate some important scientific topics to balance the clinical content and described important pathological mechanisms of disease⁸ and emerging genetic mechanisms.⁹

It has been a privilege to edit Lung Alerts as this job has allowed us to keep up to date with the literature and put us in contact with many readers of *Thorax* everywhere. We hope others find Lung Alerts as interesting as we do. We would like to take the opportunity to thank Angshu's previous Lung Alert co-editors—Terence Seemungal and John Hurst.

Competing interests

None. **Provenance and peer review** Commissioned; not externally peer reviewed.

Thorax 2010;65:767–768.
doi:10.1136/thx.2010.146100

REFERENCES

- Quint J. Allergic sensitisation to peanut may occur through inflamed skin. *Thorax* 2003;58:469.
- Bhatia P. Radiofrequency ablation of lung cancer. *Thorax* 2003;58:771.
- Roy-Choudhury S. Inflammation caused by radiofrequency ablation for lung cancer is worse after radiotherapy and in large tumours. *Thorax* 2008;63:998.
- Banerjee AK. Oral gefitinib produces symptomatic and radiological benefit in advanced NSCLC. *Thorax* 2004;59:104.
- Powrie DJ. The BODE index: a new grading system in COPD. *Thorax* 2004;59:427.
- Navani N. Preoperative integrated PET-CT scanning reduces the number of futile thoracotomies for lung cancer. *Thorax* 2009;64:1089.
- Najib M. Pleurodesis with large-particle talc poudrage is safe in malignant pleural effusion. *Thorax* 2007;62:897.
- Fletcher CE. IL6 and dendritic cells in allergic asthma. *Thorax* 2008;63:731.
- Curley R. Mutations in the LKB1 tumour suppressor gene linked to non-small cell lung cancer in Caucasians. *Thorax* 2009;64:267.