

Abstract P69 Table 1.

Confidence levels in:	Number of trainees	Mean Improvement (SD)	% Improvement	P Value
Knowing the indications for thoracoscopy	Pre=11, Post=11	1.27(0.65)	31.82	0.004
Knowing the contra-indications for thoracoscopy	Pre=11, Post=11	1.36(0.67)	34.09	0.004
Consenting the patient and quoting accurate complication rates	Pre=11, Post=11	1.36(0.5)	34.09	0.002
Being aware of the complications of thoracoscopy	Pre=10, Post=11	1.3(0.67)	32.50	0.006
Knowing the limits of safe conscious sedation and the complications	Pre=11, Post=11	1.54(0.93)	38.64	0.004
Technical ability in handling the thoracoscopy equipment	Pre=11, Post=11	2.54(0.82)	63.64	0.004
Knowing the indications for IPC insertion	Pre=10, Post=11	2.1(1.1)	52.50	0.007
Technical ability in performing an IPC insertion	Pre=11, Post=11	2.45(1.69)	61.36	0.007

Figure 1. Trainee improvement in confidence after attending the Medical Thoracoscopy and IPC insertion

the large airways. Knowledge testing was carried out using a preparing the patient metal algorithm taught during the course. The data generated was recorded on an assessment proforma entitled the Basic Bronchoscopy Simulation Assessment Tool (BBSAT).

Results Statistically significant confidence levels were found in 9 domains in all trainees after the course. Confidence levels fell significantly after assessment in the majority of domains (Table 1). Dexterity was retained at assessment, but a small fall in anatomical knowledge was demonstrated in Anaesthetic and Respiratory trainees. MCQ knowledge also showed a general decrease at assessment. The average BBSAT score was 67.2/72 (93%) indicating that all trainees had retained the basic level of skills required to perform bronchoscopy in a simulated environment.

Conclusion Overall, skills and knowledge are retained to a level required to perform bronchoscopy in a simulated environment, however confidence of trainees is significantly affected at assessment two months later. We propose that trainees are provided with mandatory bronchoscopy simulation training accompanied by assessment to ensure that skills are retained. We believe that basic bronchoscopy should be mastered before diagnostic bronchoscopy is broached; hence the use of a mastery assessment similar to the BBSAT may helpful in guiding simulation-based assessment.

P70 MANAGEMENT OF PLEURAL EFFUSIONS: ARE HEALTHCARE PROFESSIONALS ADEQUATELY TRAINED?

CL Ross, R Dimock, A Chotai, R Meeajan, T Wan, L Finney, SL Elkin; *Imperial College Healthcare NHS Trust, London, UK*

10.1136/thoraxjnl-2013-204457.220

Introduction and Objectives Management of pleural effusions is evolving with the introduction of a Best Practice Tariff for unilateral effusions and increasing use of thoracic ultrasound. However, transitions have been slow to filter through to clinical practice. This improvement project investigated the current management of pleural effusions within our Trust.

Methods We conducted: 1) An audit of all procedures for pleural effusions between 01.09.12 and 30.11.12; 2) A survey of junior doctors to establish their training provision, current practice and knowledge; and 3) A survey of acute medicine, respiratory, cardiothoracic and oncology nurses who regularly care for patients with effusions.

Results The audit reviewed 52 chest drains and 20 aspirations across 3 sites. Documentation of consent occurred in 39% of cases. Ultrasound was used in 79% of drains, but 26% of cases still used a remote 'X marks the spot' approach. 27% of procedures were done out-of-hours and 34% of patients waited over 12 hours for a chest x-ray following drain insertion. 61 junior doctors responded to the survey: 44% did not use sterile gowns when inserting drains; 11% did not use sterile gloves. No medical doctors had acquired level 1 ultrasound competence. Of the

31 nurses surveyed: 19% did not know the significance of a 'bubbling' drain; 39% did not know what 'swinging' indicated; 48% did not equate closing a 3-way tap with clamping a drain; and 26% were not aware chest drain bottles should be kept below the insertion point. 54% had never received any training regarding chest drain management. 45% of doctors and 58% of nurses were unaware of local guidelines.

Conclusions It is unlikely our findings are grossly different to practice in other inner city Trusts. With high staff turnaround, regular training for those caring for people with pleural effusions is clearly needed. Training should focus on aspects of patient safety such as image-guidance, aseptic technique and drain management. This study has led to the introduction of new local guidelines and pathways, creation of a pleural procedure kit to include aseptic equipment, implementation of an effusion bundle, and training sessions for doctors and nurses.

P71 TRAINING OF JUNIOR DOCTORS INTO THORACIC ULTRASOUND AND PLEURAL PROCEDURES—IMPACT OF A DEDICATED RESPIRATORY CONSULTANT—SUPERVISED ADVANCED NURSE PRACTITIONER-SUPPORTED SESSION

F Thaivalappil, D O'Neill, A Ionescu; *Royal Gwent Hospital, Newport, UK*

10.1136/thoraxjnl-2013-204457.221

The BTS made recommendations on pleural services. In order to stream-line the elective pleural procedures (other than thoracoscopy) a mid-week session was initiated on the pleural unit at a teaching hospital providing for over 500,000.

Aims To review the impact of a mid-week dedicated pleural session (DPS) Consultant-led and advanced nurse practitioner (ANP)-supported on the ward-based thoracic ultrasound (US) and pleural procedure training of respiratory trainees and impact on other hospital services.

Methods This is an analysis of data collected prospectively at the time of the US and procedures over 1 year. US referred to radiology were obtained from a search on radiology database. Changes over a period of six months after the DPS was started were assessed against the six months before. SPSS programme and Chi² test were used.

Results The numbers of US and procedures over 6 months was similar before and after the DPS (N = 297 vs. 260). Procedures performed on a Monday were less after DPS (Chi² = 9.21, p < 0.01), but not for Thursdays and Fridays.

Procedures done by trainees under Consultant supervision increased with the DPS (Chi² = 5.45, p = 0.019), so did those performed by the ANP (from 25 to 66 procedures, Chi² = 23.4, p < 0.001).

The proportion of chest drains inserted out of hours was 1.8% all were done by trained respiratory juniors. 20% US and procedures were referred by other departments.