

CORRESPONDENCE

Efficacy of lower-limb muscle training modalities in severely dyspnoeic individuals with COPD and quadriceps muscle weakness: response from the authors

We thank Chaplin and Latimer for their interest in the DICES study.¹ The correspondents question the validity of the power calculation, and in turn, state that we cannot conclude that high-frequency neuromuscular electrical stimulation (HF-NMES) and strength training have equivalent effects in severely dyspnoeic patients with COPD and quadriceps muscle weakness at baseline.¹ At the time of the start of the preparations of the DICES study in September 2009, we were not able to find statistical formulas where results from studies comparing two groups can be used for power calculations for three groups. Therefore, for the initial power calculation, we have used a power calculation of two groups to estimate the sample size for a three-arm study design. Initially, we did not expect a difference in change in isokinetic quadriceps peak torque between HF-NMES

and strength training. A posteriori analyses revealed that an extra ± 15 patients per group would be needed to detect a significant difference for the primary outcome, with HF-NMES being superior. To conclude, future trials are needed to investigate optimal strategies for exercise training in pulmonary rehabilitation for specific subgroups of patients with COPD, to enable development of exercise guidelines.² Based on the results of the DICES study,¹ we still believe that HF-NMES may be a good alternative for strength training in severely dyspnoeic COPD patients with quadriceps muscle weakness.

Maurice J H Sillen,¹ Frits M E Franssen,¹ Jeannet M L Delbressine,¹ Anouk W Vaes,¹ Emiel F M Wouters,^{1,2} Martijn A Spruit¹

¹Department of Research & Education, CIRO+, Centre of expertise for chronic organ failure, Horn, The Netherlands

²Department of Respiratory Medicine, Maastricht University Medical Centre (MUMC+), Maastricht, The Netherlands

Correspondence to Maurice Sillen, Department of Research & Education, CIRO+, Centre of expertise for chronic organ failure, Hornerheide 1, Horn 6085 NM, The Netherlands; mauricesillen@ciro-horn.nl

Contributors Trial concept and design: MJHS, EFMW and MAS; acquisition of data: MJHS, JMLD, AWV; analysis and interpretation of data, and drafting the article: MJHS, FMEF and MAS; revising it critically for

important intellectual content and final approval of the version to be published: all authors. MJHS had full access to all trial data and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Competing interests None.

Ethics approval Maastricht Ethical Committee, Maastricht, the Netherlands.

Provenance and peer review Not commissioned; internally peer reviewed.

To cite Sillen MJH, Franssen FME, Delbressine J M L, *et al.* *Thorax* Published Online First: [please include Day Month Year] doi:10.1136/thoraxjnl-2014-205781

Received 21 May 2014

Accepted 24 May 2014



► <http://dx.doi.org/10.1136/thoraxjnl-2014-205345>

Thorax 2014;0:1. doi:10.1136/thoraxjnl-2014-205781

REFERENCES

- 1 Sillen MJ, Franssen FM, Delbressine JM, *et al.* Efficacy of lower-limb muscle training modalities in severely dyspnoeic individuals with COPD and quadriceps muscle weakness: results from the DICES trial. *Thorax* 2014;69:525–31.
- 2 Spruit MA, Singh SJ, Garvey C, *et al.* An Official American Thoracic Society/European Respiratory Society Statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med* 2013;188:e13–64.