

Correspondence

Parachute deformity of the tricuspid valve

Sir,—We have read with interest the report¹ in which parachute deformity of the tricuspid valve is considered to be the first reported case. In 1972 we published a report of two cases of cor triatriatum.² One of them proved at necropsy to have a tricuspid valve whose anterior and posterior leaflets were fused. The chordae tendineae of the resulting valve (septal and antero-posterior) were inserted into a single smooth papillary muscle that originated in the apex of the ventricle. The tricuspid valve thus acquired the typical aspect of a parachute valve.

We think that anomalies of the tricuspid valve are more frequent than published reports indicate. This is probably because the clinical findings and pathological anatomical features are masked by the severe anatomical lesions with which they are usually associated.³

MJ MAITRE AZCARATE
Cardiología Pediátrica
Centro Especial "Ramon y Cajal"
Madrid, Spain

References

- 1 Milo L, Stark J, Macartney FJ, Anderson RH. Parachute deformity of the tricuspid valve. *Thorax* 1979; **34**:543–6.
- 2 Maitre MJ, Otero E, Quero M *et al*. Cor triatriatum: estudio anatómico-clínico. *Rev Esp Cardiol* 1972; **25**:119.
- 3 Ariza S, Cintado C, Castillo JA *et al*. Valvula tricuspid en parachute asocié a une tetralogie de Fallot. *Arch Mal Coeur* 1979; **72**:317–20.

Reply,—Dr MJ Maitre Azcarate is absolutely correct.

RH ANDERSON
Cardiothoracic Institute
London