

## Validity of a modified shuttle test in adult cystic fibrosis

Judy Bradley, Jennifer Howard, Eric Wallace, Stuart Elborn

### Abstract

**Background**—The purpose of this study was to provide some evidence of the validity of a modified shuttle test (MST) by comparing performance on the MST with peak oxygen consumption ( $\dot{V}O_{2peak}$ ) measured during a treadmill test in a group of adult patients with cystic fibrosis.

**Method**—Twenty patients with stable cystic fibrosis performed a ramped maximal treadmill test (STEEP protocol) and the MST using a randomised balanced design.

**Results**—The relationship between the distance achieved on the MST and  $\dot{V}O_{2peak}$  was strong ( $r = 0.95$ ,  $p < 0.01$ ) with 90% of the variance in  $\dot{V}O_{2peak}$  explained by the variance in MST distance. The relationship was represented by the regression equation (with 95% confidence intervals)  $\dot{V}O_{2peak} = 6.83 (2.85 \text{ to } 10.80) + 0.028 (0.019 \text{ to } 0.024) \times \text{MST distance}$ .

**Conclusion**—This study provides evidence of the construct validity of the MST as an objective measure of exercise capacity in adults with cystic fibrosis.

(Thorax 1999;54:437-439)

Keywords: modified shuttle test; cystic fibrosis

Exercise testing is useful in determining prognosis, exercise prescription, and in the evaluation of new and existing treatments in patients with cystic fibrosis. The most precise method of assessing exercise capacity is by formal laboratory tests with online analysis of expired air. However, formal tests are not widely available to clinicians working in cystic fibrosis centres, and there is debate as to the most appropriate protocol for testing. Furthermore, many patients with cystic fibrosis find these tests excessively stressful and are reluctant to perform such tests on a routine basis. Some centres have attempted to use informal tests to assess and monitor exercise capacity in patients with cystic fibrosis, but some studies have highlighted the controversy surrounding the reliability, validity, and sensitivity of many of these tests.<sup>1 2</sup>

The shuttle walking test is an incremental externally paced informal exercise test which overcomes many of the problems associated

with existing informal exercise tests. The original authors of the shuttle walking test have shown this test to be a reliable (after just one practice test), valid, and sensitive measure of exercise capacity in patients with chronic obstructive pulmonary disease.<sup>3-5</sup> We have carried out preliminary work with the shuttle walking test in adult patients with cystic fibrosis, and have shown that the walking speeds in the original test (up to a maximum of 2.37 m/s) do not elicit a maximal response in adult patients with cystic fibrosis and minimal disability as well as in patients with more severe disability. On the basis of these preliminary findings the original test was modified by the addition of three levels and, further, by permitting the patients to run. The additional stages to the original 12 stage test were: level 13, 5.63 mph, 15 shuttles; level 14, 6.00 mph, 16 shuttles; and level 15, 6.38 mph, 17 shuttles. It was hypothesised that this modified shuttle test (MST) could be used to measure peak exercise capacity objectively in adult patients with cystic fibrosis. The aim of this study was therefore to compare patients' performance on the MST with peak oxygen consumption ( $\dot{V}O_{2peak}$ ) measured directly during a treadmill test.

### Methods

Twenty patients (14 men) of mean (SD) age 25 (7) years, weight 58 (8) kg, height 1.68 (0.08) metres volunteered for the study. All patients had been familiarised with the MST and the treadmill test prior to entry into the study. Patients undertook the treadmill test on one visit to the hospital and the MST on a separate visit. The order of the tests was randomised in a counterbalanced design. The mean (SD) duration between visits was 7 (4) days. The tests were performed at approximately the same time each day. Baseline spirometric measurements (Vitalograph Alpha), resting oxygen saturations and resting heart rate (Ohmeda  $SpO_2$  monitor with ear probe), and rating of perceived breathlessness (Borg scale<sup>6</sup>) were recorded before the exercise test on each study day. The study was approved by the hospital ethical committee and informed consent was obtained from all patients.

### TREADMILL TEST

The treadmill test was a symptom limited maximal exercise test performed according to

Sport and Exercise Sciences, University of Ulster at Jordanstown, Co. Antrim BT37 0QB, UK

J Bradley  
E Wallace

Adult Cystic Fibrosis Unit, Belfast City Hospital, Belfast, Co. Antrim BT9 7AB, UK

J Howard  
S Elborn

Correspondence to:  
Dr S Elborn.

Received 21 April 1998  
Returned to author  
8 July 1998  
Revised manuscript received  
2 October 1998  
Accepted for publication  
1 December 1998

the standardised treadmill exponential exercise protocol (STEEP).<sup>7</sup> During the treadmill test measures of  $\dot{V}O_2$  (ml/min),  $\dot{V}CO_2$  (ml/min), and minute ventilation ( $\dot{V}E$ , l/min) were recorded at 15 s intervals (PK Morgan). The equation forced expiratory volume in one second ( $FEV_1$ )  $\times$  35 was used to predict maximum voluntary ventilation (MVV).<sup>8</sup> The heart rate was measured at one minute intervals using a 12 lead electrocardiogram (Model Marguette Case 15) and  $SaO_2$  was continuously monitored using an Ohmeda  $SaO_2$  monitor. At the end of the test the peak heart rate,  $SaO_2$ , and peak rate of perceived breathlessness were recorded. Reasons for stopping or failing to maintain the correct pace were also recorded.

#### SHUTTLE TEST

Using the 15 level MST, patients were required to walk/run at increasing speeds back and forth on a 10 metre course.<sup>3</sup> They were accompanied by an operator during the first minute of the test to help them pace themselves with the audiosignal. At the end of each level the patients were also told to go a little faster and were reminded that they were permitted to run at any time during the test. Patients continued with the test until they were unable to do so or failed to maintain the set pace.<sup>3</sup> Heart rate was measured at 15 s intervals using a short range telemetry device (Polar Sports Tester) and  $SaO_2$  was continuously monitored using an Ohmeda  $SaO_2$  monitor. At the end of the test the peak heart rate,  $SaO_2$ , and peak rating of perceived breathlessness were recorded. Reasons for stopping or failing to maintain the correct pace were also recorded.

#### Results

Lung function ( $FEV_1$ ) ranged from 17% to 96% predicted normal, indicating that the patients exhibited a wide variety of disease impairment. Table 1 shows that there were no significant differences between study days in baseline test parameters ( $FEV_1$ , resting heart rate, resting rating of perceived breathlessness, resting  $SaO_2$ ). Furthermore, there were no significant differences between tests in comparable physiological responses to exercise.

Table 1 Mean (SD) baseline characteristics and physiological responses to exercise testing of patients

Parameter	Shuttle test	Treadmill test	p value
Baseline characteristics			
Age (years)	25 (7)	—	—
Weight (kg)	58 (8)	—	—
Height (m)	1.67 (0.008)	—	—
$FEV_1$ (% pred)	49 (23)	48 (25)	0.31
$SaO_2$	94 (3)	94 (3)	0.39
Resting heart rate (beats/min)	95 (12)	97 (17)	0.56
Resting heart rate (% pred max)	49 (7)	50 (8)	0.46
Resting rating perceived breathlessness	0 (1)	0 (1)	0.89
Physiological responses			
$\dot{V}O_{2peak}$ (ml/kg/min)	32.85 (10.36)	—	—
$\dot{V}O_{2peak}$ (% pred)	73.45 (26.27)	—	—
$\dot{V}E_{max}$ (l/min)	57.76 (20.01)	—	—
$\dot{V}E$ (% pred)	101 (40)	—	—
MST distance (m)	—	929 (335)	—
Max heart rate (beats/min)	171 (23)	169 (24)	0.90
Max heart rate (% pred)	87 (10)	86 (11)	0.95
Peak rate perceived breathlessness	6 (1)	6 (1)	0.90
End $SaO_2$	89 (7)	88 (7)	0.10

$FEV_1$  = forced expiratory volume in one second;  $SaO_2$  = oxygen saturation;  $\dot{V}O_{2peak}$  = peak oxygen consumption;  $\dot{V}E$  = minute ventilation; MST = modified shuttle test.




Figure 1 Regression line (with 95% confidence intervals) for the relationship between performance on the modified shuttle test (MST) and peak oxygen consumption ( $\dot{V}O_{2peak}$ ) measured during treadmill testing.

A significant and moderately strong relationship was found between the distance achieved on the MST and lung function (MST vs  $FEV_1$ % predicted:  $r = 0.70$ ,  $p = 0.001$ ) and between  $\dot{V}O_{2peak}$  and lung function ( $\dot{V}O_{2peak}$  vs  $FEV_1$ % predicted:  $r = 0.78$ ,  $p < 0.001$ ). The relationship between the distance achieved on the MST and directly measured  $\dot{V}O_{2peak}$  was strong ( $r = 0.95$ ,  $p < 0.00$ ) with 90% of the variance in  $\dot{V}O_{2peak}$  explained by the variance in the MST distance. Using methods adopted from the authors of the original test,<sup>4</sup> regression analysis was used to further describe the nature of the relationship between  $\dot{V}O_{2peak}$  and MST performance. The relationship was represented by the regression equation and 95% confidence intervals:

$$\dot{V}O_{2peak} = 6.83 (2.85 \text{ to } 10.80) + 0.028 (0.019 \text{ to } 0.024) \times \text{MST distance (fig 1)}.$$

#### Discussion

The purpose of this study was to determine the validity of the MST as a measure of exercise capacity in adults with cystic fibrosis. The results show that there was a strong relationship between  $\dot{V}O_{2peak}$  and MST performance in patients with cystic fibrosis and varying degrees of lung function impairment. Ninety percent of the variation of directly measured  $\dot{V}O_{2peak}$  is explained by the variation in MST performance. This compares very favourably with the original test in which the shuttle walking test performance explained 77.4% of the variance in directly measured  $\dot{V}O_2$ . Many cystic fibrosis clinicians have no access to formal exercise testing equipment and the regression analysis used in this study provides additional information to such individuals on the nature of the relationship between  $\dot{V}O_{2peak}$  and MST performance. There was no significant difference between peak heart rate and peak rating of perceived breathlessness recorded during both exercise tests, which indicates the effectiveness of the MST to evoke a symptom limited exercise response in both mildly and more severely compromised adults with cystic fibrosis.

Most of the patients in the present study encroached on their pulmonary reserve during exercise testing (mean peak  $\dot{V}E > 70\%$  MVV). None of the patients reached their maximum predicted heart rate, as determined by the age related equation ( $220 - \text{age}$ ), and "shortness of

breath" and "fatigue" were the most common reasons reported for stopping the exercise test. These findings support the assertion that ventilatory factors rather than cardiovascular factors limit exercise tolerance in cystic fibrosis.<sup>9</sup>

The relationship between lung function and  $\dot{V}O_{2peak}$  was moderate. This finding supports previous work which indicated that impaired pulmonary function limits exercise capacity.<sup>10</sup> Lung function is not a good predictor of exercise capacity because of wide intersubject variability of exercise capacity in patients with comparable lung function. In the present study oxygen desaturation (more than 5% fall in  $SaO_2$ )<sup>11</sup> occurred in all patients with  $FEV_1$  less than 35% predicted, and in two of the patients with  $FEV_1$  of 43% and 50%. Exercises tests to establish if exercise induced desaturation occurs are therefore a necessary prerequisite to exercise prescription in cystic fibrosis. Oxygen saturation should also be intermittently monitored during exercise programmes and, if necessary, supplemental oxygen should be used to avoid oxygen desaturation.

This study has shown that there is a strong relationship between MST performance and  $\dot{V}O_{2peak}$  in adults with cystic fibrosis and thus provides evidence of the validity of this test as a measure of peak exercise capacity in adult

cystic fibrosis. Further work is required to establish the intertest reliability, test-retest reliability, and the sensitivity to change of the MST in adult patients with cystic fibrosis.

The authors would like to thank Mrs Jackie Megarry for her technical assistance with this research study.

- 1 Shephard RJ. Tests of maximum oxygen intake. A critical review. *Sports Med* 1984;1:99-124.
- 2 Swinburn CR, Wakefield JM, Jones PW. Performance, ventilation and oxygen consumption in three different types of exercise test in patients with chronic obstructive lung disease. *Thorax* 1985;40:581-6.
- 3 Singh SJ, Morgan MDL, Scott S, et al. Development of a shuttle walking test of disability in patients with chronic airways obstruction. *Thorax* 1992;47:1019-24.
- 4 Singh SJ, Morgan MDL, Hardman AE, et al. Comparison of oxygen uptake during a conventional treadmill test and the shuttle walk test in chronic airflow limitation. *Eur Respir J* 1994;7:2016-20.
- 5 Singh SJ, Morgan MDL. The use of the shuttle walking test to assess disability before and after rehabilitation in chronic respiratory disease. *Pediatr Pulmonology* 1996;340:303.
- 6 Borg GAV. Psychophysical basis of perceived exertion. *Med Sci Sports Exerc* 1982;14:377-81.
- 7 Riley M, Northridge DB, Henderson E, et al. The use of the exponential protocol for bicycle and treadmill testing in patients with chronic heart failure. *Eur Heart J* 1992;13:1363-7.
- 8 Gabdevia B, Hugh-Jones P. Terminology for measurements of ventilatory capacity. *Thorax* 1957;1:290-3.
- 9 Cerny FJ, Pullano TP, Cropp GJA. Cardiorespiratory adaptations to exercise in cystic fibrosis. *Am Rev Respir Dis* 1982;126:217-20.
- 10 Webb AK, Dodd ME. Exercise and cystic fibrosis. *J R Soc Med* 1995;88(Suppl 25):30-6.
- 11 Marcotte JE, Grisdale R, Levison H, et al. Multiple factors limit exercise tolerance in cystic fibrosis. *Pediatr Pulmonol* 1986;2:274-81.

## Collagen degrading activity associated with *Mycobacterium* species

F Massó, A Paéz, E Varela, L Díaz de León, E Zenteno, L F Montaña

Departamento de Biología Celular, Instituto Nacional de Cardiología "Ignacio Chávez", Juan Badiano 1, Tlalpan 14080, Mexico  
F Massó  
A Paéz  
E Varela  
L F Montaña

Departamento de Biología Desarrollo, Instituto de Investigaciones Biomédicas, UNAM, Mexico  
L Díaz de León

Departamento de Bioquímica, Instituto Nacional Enfermedades Respiratorias, Tlalpan, México  
E Zenteno

Correspondence to:  
Dr L F Montaña.

Received 13 August 1998  
Returned to author  
28 September 1998  
Revised manuscript received  
20 November 1998  
Accepted for publication  
5 January 1999

### Abstract

**Background**—The mechanism of *Mycobacterium tuberculosis* penetration into tissues is poorly understood but it is reasonable to assume that there is a contribution from proteases capable of disrupting the extracellular matrix of the pulmonary epithelium and the blood vessels. A study was undertaken to identify and characterise collagen degrading activity of *M tuberculosis*.

**Methods**—Culture filtrate protein extract (CFPE) was obtained from reference mycobacterial strains and mycobacteria isolated from patients with tuberculosis. The collagen degrading activity of CFPE was determined according to the method of Johnson-Wint using <sup>3</sup>H-type I collagen. The enzyme was identified by the Birkedal-Hansen and Taylor method and its molecular mass determined by SDS-PAGE and Sephacryl S-300 gel filtration chromatography using an electroelution purified enzyme.

**Results**—CFPE from *Mycobacterium tuberculosis* strain H37Rv showed collagenolytic activity that was four times

higher than that of the avirulent strain H37Ra. The 75 kDa enzyme responsible was divalent cation dependent. Other mycobacterial species and those isolated from patients with tuberculosis also had collagen degrading activity.

**Conclusions**—*Mycobacterium* species possess a metalloprotease with collagen degrading activity. The highest enzymatic activity was found in the virulent reference strain H37Rv.

(*Thorax* 1999;54:439-441)

Keywords: *Mycobacterium tuberculosis*; collagenase; metalloprotease

An increasing number of microorganisms, many of which are putative human pathogens, produce enzymes which degrade collagen.<sup>1</sup> The mechanism of penetration of *Mycobacterium tuberculosis* into the tissues and bloodstream is poorly understood but, as with other lung diseases,<sup>2,3</sup> gastrointestinal infections,<sup>4</sup> and necrotic conditions,<sup>5</sup> it is reasonable to assume that there is a contribution from proteases capable of disrupting the






Figure 1 Collagenolytic activity of *M tuberculosis* H37Rv and H37Ra culture filtrate protein extract in the presence of calcium or EDTA as calcium chelant. Results are expressed as mean (SE). Background counts per minute (cpm) have already been subtracted. Maximum activity, calculated assuming that all the radioactivity (12 500 cpm/well) was liberated, corresponded to *Cl histolyticum* (75.5).

extracellular matrix of the pulmonary epithelium and the blood vessels. The aim of this study was to identify and characterise collagenolytic activity in *Mycobacterium* reference strains and in those isolated from patients with tuberculosis.

## Methods

### MYCOBACTERIAL CULTURES

First early morning expectoration samples from two patients with pulmonary tuberculosis and two with miliary tuberculosis, pleural fluid from two patients with pulmonary tuberculosis and one with miliary tuberculosis, and the first early morning urine sample from one patient with renal tuberculosis were collected, seeded, grown, and identified as *M tuberculosis* by the National Tuberculosis Reference Laboratory. The following ATCC reference strains were donated by the referred laboratory and CENID Microbiología, México: *M tuberculosis* H37Rv and H37Ra (virulent and attenuated mycobacteria isolated from a human lung, used for susceptibility testing<sup>6,7</sup>), *M kansasii*, *M fortuitum*, *M microti*, *M terrae*, *M avium*, and *M goodnae*. These strains were seeded in protein-free Proskauer-Beck-Youmans (PBY) medium and maintained at 37°C for 7–9 weeks before bacteria were harvested.<sup>8</sup> Some cultures were maintained for as long as 12 weeks before harvesting.

### CULTURE FILTRATE PROTEIN EXTRACT (CFPE)

The culture medium was separated from the bacterial mass by filtration through 0.45 µm and 0.22 µm membrane filters (Millipore Corp). Proteins were precipitated with solid ammonium sulphate crystals (80% final saturation) and centrifuged for 30 minutes at 16 000g. The precipitate, suspended in 5 ml phosphate buffered saline (PBS) containing 0.2 mg/ml phenylmethylsulphonyl fluoride (Sigma Chemicals), was dialysed extensively against PBS and 0.5 ml aliquots (6 mg/ml) were stored in liquid nitrogen until used. Some filtrates were precipitated at 40% and 60% (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> final saturation.

### COLLAGEN

Collagen was obtained from the skin of two month old female Wistar rats according to the method described by Epstein.<sup>9</sup> The specific activity of radiolabelled collagen was 1 × 10<sup>6</sup> cpm/mg of protein.

### COLLAGENOLYTIC ASSAY

The enzymatic activity was determined according to the method of Johnson-Wint<sup>10</sup> using 120 µg CFPE and 12 500 cpm radioactive collagen/well. Some experiments were performed with 20 mM/well EDTA tetrasodium salt (Sigma Chemicals). Controls included buffer, 0.001% trypsin (type IX from porcine pancreas), and *Clostridium histolyticum* collagenase (high purity, type III fraction A).

### COLLAGENASE IDENTIFICATION

The enzyme was identified using a modification of the method described by Taylor and Birkedal-Hansen<sup>11</sup> that included elimination of SDS from the gel with 3% Triton X-100 and overnight incubation at 37°C. The region with collagenolytic activity in the CFPE containing gel (β-mercaptoethanol reduced and boiled sample/lane in 10% slab gels<sup>12</sup>) was located, purified by electroelution, dialysed, lyophilised, and kept at –20°C until used. The non-denatured state of the substrate was corroborated by the inclusion of trypsin, an enzyme incapable of breaking down collagen but highly active upon gelatin.

### MOLECULAR MASS DETERMINATION

The molecular mass of the enzyme (75 000 by SDS-PAGE) was determined by Sephacryl S-300 gel filtration chromatography. 500 µg of enzyme partially purified by electroelution were suspended in 0.05 M Tris HCl buffer (pH 7.5) containing 4 mM CaCl<sub>2</sub>, poured onto a 1.2 × 100 cm glass column previously equilibrated with the same buffer, and eluted with the Tris HCl buffer at a flow rate of 12 ml/hour. Standards included horse heart myoglobin (17 kDa), bovine serum albumin (66 kDa), purified human IgG (150 kDa), and bovine liver catalase (240 kDa). The activity of each fraction was tested as described above.

## Results

The collagenolytic activity of CFPE from H37Rv was four times higher than that from H37Ra (43.4% and 9.1%, respectively). Addition of EDTA to the H37Rv CFPE blocked the enzymatic activity by 88%, thus establishing that the enzyme in the virulent strain was dependent on divalent cations (fig 1). Similar results were obtained with replicate cultures. The enzymatic activity was secondary to a 75 kDa protein which was clearly established by the degradation of collagen containing gels and molecular mass determination by Sephacryl S-300 gel filtration chromatography.

The collagenolytic activity was not changed by the use of ammonium sulphate saturation to precipitate the CFPE, the amount of H37Rv CFPE precipitated with a final ammonium sulphate saturation of 40%, 60% and 80% being

Table 1 Mean (SE) collagenolytic activity of culture filtrates obtained from various mycobacterial species

	With Ca		With EDTA	
	Counts per minute (cpm)	Enzyme activity	Counts per minute (cpm)	Enzyme activity
Trypsin	1088 (203)	8.7	528 (18)	4.2
<i>Cl histolyticum</i>	10982 (904)	87.8	2888 (878)	23.1
<i>M microti</i>	3234 (113)	25.8	1094 (140)	8.7
<i>M kansasii</i>	1038 (348)	8.3	888 (282)	7.1
<i>M avium</i>	2621 (229)	20.9	1277 (305)	10.2
<i>M terrae</i>	768 (63)	6.1	433 (69)	3.4
<i>M gordonae</i>	542 (74)	4.3	485 (45)	3.8
<i>M fortuitum</i>	272 (33)	2.1	174 (46)	1.3

Background cpm have already been subtracted. The percentage of activity was calculated assuming that all the radioactivity (12 500 cpm/well) was liberated by the enzyme.

Table 2 Mean (SE) collagenolytic activity of culture filtrates obtained from mycobacteria isolated from patients with tuberculosis

Diagnosis*	Source†	With Ca		With EDTA	
		Counts per minute (cpm)	Enzyme activity	Counts per minute (cpm)	Enzyme activity
Trypsin		792 (130)	6.3	404 (103)	3.2
<i>Cl histolyticum</i>		9763 (559)	78.1	2367 (362)	18.9
Pulmonary TB	Sputum	1160 (284)	9.2	510 (462)	4.0
Pulmonary TB	Sputum	4198 (204)	33.5	684 (642)	5.4
Pulmonary TB	Pleural fluid	1783 (462)	14.2	348 (90)	2.7
Pulmonary TB	Pleural fluid	1376 (248)	11.0	1142 (108)	9.1
Miliary TB	Sputum	2276 (98)	18.2	874 (138)	6.9
Miliary TB	Sputum	4986 (230)	39.8	566 (32)	4.5
Miliary TB	Pleural fluid	2864 (127)	22.9	814 (175)	6.5
Renal TB	Urine	2316 (224)	18.5	1238 (148)	9.9

\*All patients were PPD positive.

†Source refers to the secretion from which the mycobacteria were isolated.

Background cpm have already been subtracted. The percentage of activity was calculated assuming that all the radioactivity (12 500 cpm/well) was liberated by the enzyme.

4380 (304) cpm, 4224 (215) cpm, and 4146 (182) cpm, respectively. To determine the influence of culture time on enzymatic activity, five H37Rv cultures were seeded and harvested after 32, 38, 46, 70, and 103 days of culture and had CFPE activity of 20%, 29%, 34%, 33%, and 15%, respectively, indicating that the collagenolytic activity decays with time.

The collagenolytic activity of CFPE obtained from mycobacteria other than *M tuberculosis* was also tested. *M microti* showed the highest level of activity followed by *M avium*, *M kansasii*, *M terrae*, *M gordonae*, and *M fortuitum* which had the least activity (table 1). Enzymatic variability was also observed in culture filtrates obtained from mycobacteria isolated from patients with tuberculosis (table 2).

## Discussion

The presence of collagenolytic activity in the CFPE of *M tuberculosis* H37Rv strain was secondary to a 75 kDa enzyme whose activity was dependent on divalent cations and was blocked by EDTA, suggesting a mechanism of non-competitive inhibition. These characteristics are similar to those reported for *Clostridium histolyticum* collagenase<sup>13</sup> and the majority of collagenases.<sup>14</sup> Even though we recognise the possible influence of collagenase degradation before harvesting, the differences between the CFPE samples of H37Rv virulent and H37Ra avirulent mycobacterial strains were striking and were independent of batch to batch variations, source of seed, and time of culture.

None of the mycobacteria we tested had activity greater than *M tuberculosis* H37Rv. As in the case of catalase activity, once considered

a virulence factor,<sup>15</sup> collagen degrading activity is important but not vital in the pathogenesis of the disease since there was no uniform activity in *M avium* and *M kansasii*, both of which induce tuberculosis. *M fortuitum* and the non-disease inducers *M terrae* and *M gordonae* had very poor enzyme activity. Although tissue destruction is secondary to an enhanced cellular immune response, mycobacterial metalloproteinases might be important for bacterial penetration<sup>16</sup> as has been shown with other infectious agents.<sup>4 17</sup> Despite the above considerations, it was interesting to observe that *M tuberculosis* isolated from tuberculosis patients with different clinical syndromes (renal, miliary, pulmonary) had greater activity than the avirulent reference strain H37Ra.

Our results confirm the presence of collagen degrading enzymatic activity in the culture filtrate protein extract of all the mycobacteria tested. Extracellular proteases may also play a role in the pathogenesis of mycobacterial infection. However, the recent resolution of the genome for *M tuberculosis* H37Rv (website: www.sanger.ac.uk) predicts at least 38 genes coding for proteins of virulence and more than 250 macromolecular and/or micro-molecular degrading proteins. One of these, the product of gene Rv0198c, a zinc metalloprotease with a molecular weight of 73.8 kDa, is similar to the enzyme we describe in this work.

- Harrington DJ. Bacterial collagenases and collagen-degrading enzymes and their potential role in human disease. *Infect Immun* 1996;64:1885-91.
- Bruce MC, Poncz I, Klinger JD, et al. Biochemical and pathologic evidence for proteolytic destruction of lung connective tissue in cystic fibrosis. *Am Rev Respir Dis* 1985; 132:529-35.
- Liener IE, Twumasi DY. Proteases from purulent sputum. Purification and properties of the elastase and chemotrypsin-like enzymes. *J Biol Chem* 1977;252:1917-26.
- Serrano JJ, De la Garza M, Moreno MA, et al. *Entamoeba histolytica*: electrodense granule secretion, collagenase activity and virulence are altered in the cytoskeleton mutant BG-3. *Mol Microbiol* 1994;11:787-92.
- Wright LG, Atkinson RK. The involvement of collagenase in the necrosis induced by the bites of some spiders. *Comp Biochem Physiol* 1992;102:125-8.
- Van Soelingen D, Hoogenboezem T, de Haas PE, et al. A novel pathogenic taxon of the *Mycobacterium tuberculosis* complex, Canetti: characterization of an exceptional isolate from Africa. *Int J Syst Bacteriol* 1997;47:1236-45.
- McMurray DN, Carlomagno MA, Cumberland PA. Respiratory infection with attenuated *Mycobacterium tuberculosis* H37Ra in malnourished guinea pigs. *Infect Immun* 1983;39: 793-9.
- Parra C, Montano LF, Huesca M, et al. Inhibition of mitogenesis induced by phytohemagglutinin and Lens culinaris lectin in adherent-cell supernatants treated with protein extract of *Mycobacterium tuberculosis*. *Infect Immun* 1986;52:309-13.
- Epstein EH. [ $\alpha_1$ (III)]<sub>3</sub> human skin collagen. *J Biol Chem* 1974;249:3225-31.
- Johnson-Wint B. A quantitative collagen film collagenase assay for large number of samples. *Anal Biochem* 1980;104: 175-81.
- Taylor RE, Birkedal-Hansen H. Detergent activation of latent collagenase and resolution of its component molecules. *Biochem Biophys Res Commun* 1982;107:1173-8.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680-5.
- Van Wart HE, De Bond M. Characterization of the individual collagenases from *Clostridium histolyticum*. *Biochemistry* 1984;23:3085-91.
- Harper E. Collagenases. *Ann Rev Biochem* 1980;49:1063-78.
- Youmans GP. Virulence of mycobacteria. In: Youmans GP, ed. *Tuberculosis*. Philadelphia: WB Saunders, 1979: 194-201.
- Chang JC, Wysocki A, Tchou Wong KM, et al. Effect of *Mycobacterium tuberculosis* and its components on macrophages and the release of matrix metalloproteinases. *Thorax* 1996;51:306-11.
- Merkel JR, Smith GC. Collagenolytic activity of *Vibrio vulnificus*: potential contribution on its invasiveness. *Infect Immun* 1982;35:1155-6.