

Conclusions Most healthcare professionals understand the indication for the use of oxygen. However, knowledge of the appropriate use of delivery devices and target saturations was noted to be poor. Targeted educational sessions have been developed to address the above issues. We are currently in the process of trialling a new Kardex to ensure that all oxygen used on the ward within our Trust is prescribed. A Hospital Oxygen Steering Group is being set up with the function of overseeing the management and prescription of oxygen.

REFERENCES

- Oxygen safety in hospitals: Rapid Response Report; NPSA/2009/RRR006, September 2009.
- BTS guideline for emergency oxygen use in adult patients. *Thorax* 2008 Oct; **63**(Suppl. 6):vi1–68.

Clinical studies in cystic fibrosis

P96 HAPLOTYPE VARIABILITY IN PSEUDOMONAS AERUGINOSA (PSA) STRAINS IN THE ADULT CF POPULATION

doi:10.1136/thx.2010.150979.47

¹A Ashish, ²J Fothergill, ²E Mowat, ¹J Greenwood, ²C Winstanley, ¹M Walshaw. ¹Liverpool Heart and Chest Hospital, Liverpool, UK; ²University of Liverpool, Liverpool, UK

Introduction Morphological type variation in Psa is thought to aid its survival in hostile environments, including the CF lung. Previous studies using PGFE have shown wide heterogeneity in LES strain, despite this, little is known about their phenotypic diversity (haplotypes). To look at this further, we compared the haplotypes of the most successful and prevalent UK transmissible Psa (the Liverpool epidemic strain, LES) with those of sporadic strains and assessed their response to antibiotic pressure.

Methods Sputum from two groups of adult CF patients chronically infected (for at least 4 years) with sporadic single Psa strains or LES was analysed at the beginning and end of an intravenous antibiotic-treated exacerbation, where every patient had subjective and spirometric improvement, and also for LES patients over a period of time (at least 4 months) when they were in a stable state. From each sample, 40 single Psa colonies were selected (with every morphological type proportionately represented), and colony morphology, susceptibility to six antibiotics (ciprofloxacin, ceftazidime, colomycin, meropenem, tobramycin, piperacillin/tazobactam), hypermutability (rate of spontaneous mutation to rifampicin resistance) and auxotrophy (ability to grow on glucose M9 media) determined. Each Psa colony could potentially exhibit any of nine combinations: antibiotic susceptibility patterns, auxotrophy, hypermutability, and six colony morphologies. The resulting haplotypes were compared for LES and unique strains using 'e-burst' software.

Results Overall, 151 unique haplotypes were defined (96 LES, 75 sporadic, 20 shared), with LES demonstrating the greater number ($X^2=4.67$, $p<0.03$). Following antibiotic pressure, there was an alteration in haplotypes which was similar in both groups (LES 54%, sporadic 64%, $p=0.28$): but during the stable state LES demonstrated a significantly greater change in haplotypes (89%, $p=0.004$).

Conclusion A change in haplotypes occurs in all CF Psa strains during treatment for clinical exacerbations, presumably as a defence mechanism against the challenge of hostile antibiotic therapy. However, the enhanced haplotype variability of transmissible strains with time when there is no antibiotic pressure may be a

strategy that confers a survival advantage compared to sporadic strains: studies are under way to explore this further.

P97 DO INTRAVENOUS ANTIBIOTICS INFLUENCE ARTERIAL STIFFNESS IN ADULTS WITH CYSTIC FIBROSIS?

doi:10.1136/thx.2010.150979.48

¹J H Hull, ²R Garrod, ³T B Ho, ³R K Knight, ⁴J R Cockcroft, ⁴D J Shale, ⁵C E Bolton. ¹Kingston University and St George's Hospital, London, UK; ²Kings College Hospital, London, UK; ³Frimley Park NHS Foundation Trust, Frimley, UK; ⁴Wales Heart Research Institute Cardiff University, Cardiff, UK; ⁵NIHR Nottingham Respiratory BMU, Nottingham, UK

Introduction and objectives Adults with cystic fibrosis (CF) have evidence of increased arterial stiffness which is associated with a chronic systemic inflammatory status (1). We hypothesised that an intervention which decreased inflammation in patients with CF would improve arterial stiffness and we therefore evaluated a 2-week course of intravenous antibiotics on large artery haemodynamics in adults with CF.

Methods We recruited 18 adult patients with CF immediately preceding (visit 1) and following 14 days of intravenous antibiotics (visit 2); prompted by clinical indication. Arterial stiffness was determined by supine applanation tonometry to obtain augmentation index (AIx) which was heart rate adjusted, and aortic pulse wave velocity (aPWV) (SphygmoCor). In addition, heart rate (HR), blood pressure (BP), spirometry and CRP were measured.

Results Complete data was available for 15 patients ($n=3$ not suitable); mean (SD) age 28 (6) years and BMI 21.2 (2.4) kg/m². All but one patient reported symptomatic improvement following treatment. Percent predicted forced expiratory volume in one second (FEV₁) improved and CRP was reduced at visit 2 (Abstract P97 Table 1). There was a trend towards reduction in HR ($p=0.06$) whilst peripheral and central BP and aPWV were similar between visits. However, of particular note, HR adjusted AIx was reduced (Abstract P97 Table 1). The change in AIx was related to FEV₁ % predicted ($r=0.77$) and BMI ($r=0.71$) at visit 1 (both $p<0.01$), but not CRP ($p=0.13$).

Abstract P97 Table 1

Variable	Visit 1	Visit 2
FEV ₁ % predicted	54 (20)	60 (20)*
CRP (mg/l)†	6.9 (6.4)	2.6 (7.6)*
HR (bpm)	75 (15)	70 (12)
Peripheral MAP (mm Hg)	89 (9)	90 (8)
AIx (%)	10.9 (10.9)	8.1 (10.9)*
aPWV (m/s)	6.3 (1.0)	6.1 (1.1)

Values are Mean (SD).

*Represents $p<0.05$.

†Geometric mean (SD).

Conclusions Intervention with intravenous antibiotics in adults with CF is associated with a reduction in AIx, a composite measure of systemic arterial stiffness and wave reflection. Modulation of systemic inflammation results in beneficial haemodynamic changes that may be key in maintaining cardiovascular health in adults with CF.

REFERENCE

- Hull JH, et al. *Eur Respir J* 2009;**34**(6):1322–8.