

centres studied. The data suggest capreomycin might be associated with less ototoxicity when compared to amikacin. There are no UK guidelines to inform best practice and better evidence, including clinical and cost-effectiveness studies, is needed to inform the implementation of current technology including genetic testing.

S3 FACTORS FOR SUCCESSFUL TREATMENT COMPLETION AMONG MDR TB CASES IN THE UK

doi:10.1136/thx.2010.150912.3

¹L F A Anderson, ¹S T Tamme, ²J W Watson, ¹I A Abubakar. ¹Health Protection Agency, London, UK; ²Department of Respiratory Medicine, St James University Hospital, Leeds, UK

Background UK guidelines recommend at least 18 months treatment for patients with multi-drug resistant tuberculosis (MDR TB). Prior to 2008, treatment completion was only available for 12 months nationally in the UK, therefore the proportion completing treatment was unknown. There is also a lack of recent guidance for the treatment and management of MDR TB cases in resource rich low TB incidence settings.

Aims To report the treatment outcome at 24 months for MDR TB patients between 2004 and 2007 and examine treatment regimens and management associated with successful outcomes, taking clinical, social and demographic factors into account.

Methods A retrospective cohort study was used to follow-up MDR TB patients at 24 months. Patients were identified using the national surveillance system which contains demographic and clinical characteristics of cases and is matched routinely to reference laboratory data. Questionnaires were sent out to case managers to collect information on outcome of care and associated risk factors for treatment completion.

Results 69.6% (142/204) of patients completed treatment at 24 months and 2.9% (6) of these completed treatment within 12 months. 4.4% (9) were still on treatment, 6.9% (14) had their treatment stopped, 6.9% (14) died, 7.8% (16) were lost to follow-up and 4.4% (9) were transferred overseas. Only 40.3% (77/191) patients started on treatment received directly observed therapy at any time. Treatment with a recommended fluoroquinolone (OR=2.3; 95% CI 1.2 to 4.2, $p<0.000$) or a bacteriostatic (OR 2.86; 95% CI 1.6 to 5.3, $p=0.001$), a change in treatment regimen (OR=2.2; 95% CI 1.2 to 4.0, $p=0.01$) and treatment with four or more effective drugs (OR=2.0; 95% CI 1.1 to 3.8, $p=0.02$) were significantly associated with a successful treatment outcome.

Conclusion The proportion of MDR TB cases completing treatment is similar to cases with drug susceptible disease due to the use of individualised treatment regimens. However, treatment completion still remains below World Health Organization targets.

S4 EVALUATION OF TB PEER EDUCATORS—ESSENTIAL PARTNERS IN METROPOLITAN TB CONTROL

doi:10.1136/thx.2010.150912.4

¹J Hall, ²S Bethell, ²S Hellenen, ¹A Story, ³M Lipman. ¹Find & Treat, Department of Health TB Projects, London, UK; ²Groundswell, London, UK; ³Royal Free Hampstead NHS Trust, London, UK

Background One in six of all notified tuberculosis (TB) cases in London are among homeless people, problem drug and alcohol users and prisoners (hard-to-reach). These groups are at high risk of delayed diagnosis, infectious and drug resistant TB and poor treatment outcomes. The Berlin Declaration (2007) stated that affected communities are essential partners in TB control. While initiatives involving hard-to-reach communities in HIV control have proven effective, evidence to

support their contribution to TB control activities is lacking. We aimed to improve service access and uptake of TB screening among hard-to-reach groups by harnessing the authentic voice and experience of former TB patients from these affected communities.

Method Seven former TB patients with a history of homelessness and drug/alcohol dependence were recruited and trained as peer educators to work alongside TB clinics and a mobile x-ray screening service. Qualitative and quantitative methods were used to evaluate their impact on service access and screening uptake.

Results Peers recruited 3200 hard-to-reach clients at 101 screening sessions resulting in 45 hospital referrals between May 2009 and February 2010. Following TB peer training of homeless shelter hostel workers, screening uptake increased from 44% to 75%. Subsequent structured interviews with service users highlighted the importance of peer educators in raising TB awareness and promoting service access.

Conclusion Our evaluation demonstrated that trained peer educators can improve service access and TB screening uptake in the short and medium term in hard-to-reach groups. The success of this approach argues for greater peer educator involvement in strategies to control metropolitan TB.

Funding This project was supported by the Department of Health.

S5 PREVALENCE OF LATENT TUBERCULOSIS INFECTION IN IMMIGRANTS TO THE UK: FINDINGS FROM A MULTI-CENTRE STUDY

doi:10.1136/thx.2010.150912.5

¹M Pareek, ²J P Watson, ³O M Kon, ⁴G Woltmann, ¹A Lalvani. ¹Imperial College London, London, UK; ²Leeds Teaching Hospital NHS Trust, Leeds, UK; ³Imperial College Healthcare NHS Trust, London, UK; ⁴University Hospitals Leicester NHS Trust, Leicester, UK

Background Notifications for tuberculosis (TB) in the United Kingdom (UK) increased by 30% between 1992 and 2007 with over-seas-born migrants accounting for 72% of cases in 2007. Much of this increase is due to the synergy of migration from high-TB burden countries and the reactivation of pre-existing latent TB infection (LTBI). Since 2006, NICE guidance advocates screening for LTBI in all children from countries with a TB incidence $>40/100\,000$ and adults from Sub-Saharan Africa and countries with a TB incidence $>500/100\,000$. The rationale underpinning these guidelines remains unclear particularly as there are little data on the prevalence of LTBI in immigrants to the UK from regions with different TB incidence rates.

Aims To quantify the prevalence of LTBI in immigrants, assess factors associated with IGRA positivity and to determine LTBI yield from current screening thresholds recommended by NICE.

Methods Analysis of data prospectively obtained through IGRA testing (QuantiFERON-Gold/QuantiFERON-Gold-in-tube) of immigrants in three different centres in the UK during 2008–2010. Descriptive analyses were undertaken. Yields at different screening thresholds were calculated. Subsequent univariate and multivariate analyses were undertaken to assess independent factors associated with IGRA positivity; $p<0.05$ considered significant.

Results In total 915 immigrants were screened during the study period. Median age was 26.8 years (interquartile range 22–33), 50.6% were female and 72.5% had previously been BCG vaccinated. 48.9% and 28.5% of migrants screened originated from the Indian Subcontinent and Sub-Saharan Africa, respectively. Overall 911/915 individuals had determinate IGRA results—199 (21.7%) tested positive and 712 (77.8%) tested negative; 4 (0.4%) individuals had indeterminate results. Multivariate analysis revealed that increasing age ($p<0.0001$) and increasing TB incidence of country of origin ($p=0.0014$) were independently associated with IGRA positivity. Applying current NICE guidance resulted in a yield of 76/311 (24.4%) but would miss 61.8% of the immigrant population with LTBI (Abstract S5 Table 1).