

Cell signalling and cell responses in pulmonary vascular disease

S150 SMAD-DEPENDENT AND SMAD-INDEPENDENT INDUCTION OF ID1 BY PROSTACYCLIN ANALOGUES INHIBITS PROLIFERATION OF PULMONARY ARTERY SMOOTH MUSCLE CELLS IN VITRO AND IN VIVO

doi:10.1136/thx.2010.150953.1

Jun Yang, Xiaohui Li, Rafia S Al-Lamki, Mark Southwood, Jing Zhao, Andrew M Lever, Friedrich Grimminger, Ralph T Schermuly, Nicholas W Morrell. *University of Cambridge, Cambridge, UK*

Introduction and Objectives Mutations in the bone morphogenetic protein type II receptor (BMPR-II) are responsible for the majority of cases of heritable pulmonary arterial hypertension (PAH). Mutations lead to reduced Smad1/5-driven expression of inhibitor of DNA binding protein 1 (Id1) and loss of the growth suppressive effects of BMPs. The impact of existing PAH therapies on BMP signalling is lacking. Because prostacyclin analogues are effective treatments for clinical PAH, we hypothesised that these agents enhance Smad1/Id1 signalling.

Methods Iloprost alone induced Id1 expression in human pulmonary artery smooth muscle cells (PASMCs), an effect that was independent of Smad1/5 activation but dependent on a cAMP-responsive element in the Id1 promoter. In addition, iloprost and treprostinil enhanced BMP-induced phosphorylation of Smad1/5 and Id1 expression in a cAMP-dependent manner. The mechanism involved suppression of inhibitory Smad, Smad6. Furthermore, iloprost rescued the deficit in Smad1/5 phosphorylation and Id gene expression in PASMCs harbouring mutations in BMPR-II and restored growth suppression to BMP4 in mutant PASMCs.

Results We confirmed a critical role for Id1 in PASMC proliferation. Reduced expression of Id1 was observed in concentric intimal lesions of heritable PAH cases. In the monocrotaline rat model of PAH, associated with reduced BMPR-II expression, we confirmed that treprostinil inhibited smooth muscle cell proliferation and prevented progression of PAH while enhancing Smad1/5 phosphorylation and Id1 gene expression.

Conclusions Prostacyclin analogues enhance Id1 expression in vitro and in vivo and restore deficient BMP signalling in BMPR-II mutant PASMCs.

S151 TRAIL DEFICIENCY IS PROTECTIVE IN EXPERIMENTAL PULMONARY ARTERIAL HYPERTENSION

doi:10.1136/thx.2010.150953.2

¹A G Hameed, ¹J Chamberlain, ¹N D Arnold, ¹S E Francis, ¹C M H Newman, ²D C Crossman, ¹A Lawrie. ¹University of Sheffield, Sheffield, UK; ²NIHR Cardiovascular Biomedical Research Unit, Sheffield, UK

Introduction and objectives Despite advances in the overall management of Pulmonary Arterial Hypertension (PAH) significant morbidity and poor prognosis remain a major clinical problem. Identifying key pathways in the pathogenesis of this disease will allow development of more targeted therapies aimed at treating PAH. There is emerging evidence to support that Tumour Necrosis Factor (TNF)-Related Apoptosis-Inducing Ligand (TRAIL) plays an important role in vascular biology. We have recently reported expression of TRAIL from lesions of patients with advanced Idiopathic PAH. To further determine the role of TRAIL in the pathogenesis of PAH we used a diet-induced murine model of PAH.

Methods ApoE^{-/-}, TRAIL^{-/-} and ApoE^{-/-}/TRAIL^{-/-} double null mice were fed chow or Paigen (high fat, cholate-containing diet) for 8 weeks. They underwent echocardiographic assessment prior to

right heart catheterisation using the internal jugular venous route. The heart and right lung were perfusion fixed with 10% formalin for subsequent determination of right ventricular mass and Immunohistochemistry of pulmonary vascular lesions. The left lung was immediately frozen in liquid nitrogen for subsequent determination of protein and RNA by western Immunoblotting and quantitative PCR. Identical to above ApoE^{-/-}/TRAIL^{-/-} double null mice were also treated with recombinant murine TRAIL, or saline (4 week infusion via an osmotic mini pump) at the time of commencing the diet for 8 weeks.

Results Compared to control chow fed mice, ApoE^{-/-} mice fed the Paigen diet developed significant elevation of Right Ventricular Systolic Pressure (RVSP) (23 mm Hg vs 50 mm Hg n=7 p<0.001), pulmonary vascular resistance and arteriolar remodelling. ApoE^{-/-}/TRAIL^{-/-} double null mice fed the Paigen diet, were protected from these haemodynamic (RVSP 27 mm Hg n=6 p<0.05) and pulmonary vascular remodelling changes. Moreover, the PAH phenotype was re-established in the ApoE^{-/-}/TRAIL^{-/-} double null mice by the administration of exogenous recombinant TRAIL.

Conclusions TRAIL is a critical mediator in disease pathogenesis of PAH in the diet-induced murine model of PAH. Targeting TRAIL could provide a novel therapeutic approach to the treatment of PAH. Work is ongoing to determine if this approach can stabilise or reverse established disease, in both this, and rat, experimental models of PAH.

S152 DEXAMETHASONE REVERSES ESTABLISHED MONOCROTALINE-INDUCED PULMONARY HYPERTENSION IN RATS AND INCREASES PULMONARY BMPR2 EXPRESSION

doi:10.1136/thx.2010.150953.3

¹L C Price, ¹S J Wort, ²D Montani, ²C Tcherakian, ²P Dorfmueller, ²R Souza, ¹D Shao, ²G Simonneau, ³L S Howard, ¹I Adcock, ²M Humbert, ²F Perros. ¹National Heart & Lung Institute, Royal Brompton Hospital, London, UK; ²Centre National de Référence de l'Hypertension Pulmonaire Sévère, Hôpital Antoine Bécélère, Clamart, F-92140, France; ³Université Paris-Sud, Paris, France; ⁴Hammersmith Hospital, Imperial College Healthcare NHS Trust, Du Cane Road, London, UK

Background Pulmonary arterial hypertension (PAH) is associated with pulmonary vascular inflammation and dysregulated bone morphogenetic protein receptor type 2 (BMPR2) signalling in both human and experimental PAH. We evaluated the effects of dexamethasone on established monocrotaline-induced PAH in rats for potential reversal of PAH, at time points when pulmonary vascular remodelling has already developed (from day 14 after a single injection of monocrotaline at day 0), and for the effects on pulmonary IL6 and BMPR2 expression.

Methods Saline-treated controls, monocrotaline-exposed, monocrotaline-exposed and dexamethasone-treated rats (5 mg/kg/day, 1.25 mg/kg and 2.5 mg/kg/48 h given from day 14–28 and day 21–35) were evaluated at day 28 and day 35 following monocrotaline for pulmonary vascular haemodynamic parameters, right ventricular hypertrophy, morphometry, immunohistochemistry, whole lung IL-6 and BMPR2 expression by quantitative real-time PCR (qRT-PCR).

Results Dexamethasone significantly improved pulmonary haemodynamics and morphometric indices of pulmonary vascular remodelling, reversing PAH when given at day 14–28, day 21–35 following monocrotaline, as well as improving survival in monocrotaline-exposed rats compared to controls (log rank p<0.0001). Dexamethasone reduced both monocrotaline-induced whole lung IL-6 overexpression (p<0.05), as well as reducing IL-6-expressing adventitial inflammatory cell infiltration as assessed by immunohistochemistry. This was associated with pulmonary BMPR2