

## Images in Thorax

### Multiple pulmonary nodules: and if it was not a cancer

A 72-year-old woman was admitted in the Cardiology Department for respiratory insufficiency. ECG revealed atrial fibrillation, and echocardiography revealed heart failure with an estimated left ventricular ejection fraction of 25%. Chest roentgenogram revealed multiple nodules, and chest CT scan revealed multiple dense nodules surrounded by ground-glass attenuation (halo sign) with unilateral pleural effusion (figure 1). A bronchoalveolar lavage was performed to rule out any infection.

Anticoagulation was begun before a successful external cardioversion, followed by oral amiodarone treatment. Diuretics were increased without any other medication. One week later, a control chest CT scan revealed the disappearance of all nodules, as well as of the pleural effusion (figure 2).

#### LEARNING POINTS

Multiple pulmonary nodules evoke metastatic cancer, and the rate of malignancy in nodules >20 mm is 81%.<sup>1</sup> The halo sign, defined as ground-glass attenuation surrounding a solid nodule, is usually associated with haemorrhagic nodules.<sup>2</sup> First described in immunocompromised patients with invasive aspergillosis, it has been reported in metastases from hypervascular tumours and vasculitis. The halo sign may also correspond to tumorous or inflammatory infiltration, as in bronchiolo-alveolar carcinoma, lymphoma or viral infection. Although the association of ground-glass opacities, interlobular septal and peri-



**Figure 1** CT scan on arrival, revealing multiple nodules.



**Figure 2** CT scan after cardiac therapy.

bronchovascular interstitial thickening and predominantly right-sided pleural effusion is the most common presentation, cardiac insufficiency may occasionally take the form of multiple ill-defined centrilobular opacities that may have a nodular appearance.<sup>3 4</sup>

**Raphael Borie,<sup>1</sup> Marie-Pierre Debray,<sup>2</sup> Guillaume Jondeau,<sup>3</sup> Bruno Crestani<sup>1</sup>**

<sup>1</sup>Service de Pneumologie A, Centre de compétence des maladies pulmonaires rares, Hôpital Bichat, Assistance Publique-Hopitaux de Paris, Paris, France; <sup>2</sup>Service de Radiologie, Hôpital Bichat, Assistance Publique-Hopitaux de Paris, Paris, France; <sup>3</sup>Département de Cardiologie, Hôpital Bichat, Assistance Publique-Hopitaux de Paris, Paris, France

**Competing interests** None.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

Accepted 6 May 2010

Published Online First 29 July 2010

*Thorax* 2011;**66**:1018. doi:10.1136/thx.2010.134726

#### REFERENCES

1. **McWilliams A**, Mayo J. Computed tomography-detected noncalcified pulmonary nodules: a review of evidence for significance and management. *Proc Am Thorac Soc* 2008;**5**:900–4.
2. **Pinto PS**. The CT halo sign. *Radiology* 2004;**230**:109–10.
3. **Gluecker T**, Capasso P, Schnyder P, *et al*. Clinical and radiologic features of pulmonary edema. *Radiographics* 1999;**19**:1507–31; discussion 32–3.
4. **Ribeiro CM**, Marchiori E, Rodrigues R, *et al*. Hydrostatic pulmonary edema: high-resolution computed tomography aspects. *J Bras Pneumol* 2006;**32**:515–22.