

Finger-in-glove sign in bronchial atresia

A 31-year-old, HIV-positive man presented with a history of chronic dry cough. Chest x-ray showed a rounded, branching opacity in the left upper lung (finger-in-glove sign, figure 1A). CT showed mucoid impaction, segmental hyperlucency and decreased vascularity of the left upper lobe (figure 1B). 3D reconstruction of the bronchial tree revealed an atretic apico-posterior segmental bronchus of the left upper lobe confirming the diagnosis of congenital bronchial atresia (figure 1C).

Congenital bronchial atresia is a rare, mostly incidental radiographic finding, and consists of atresia of a lobar, segmental or subsegmental bronchus at or near its origin.^{1 2} The apico-posterior segmental bronchus of the left upper lobe is most commonly affected. Distal to the bronchial atresia secretions accumulate, leading to mucoid impaction surrounded by segmental hyperlucency caused by a combination of trapped air and oligoemia. The differential diagnosis of the finger-in-glove sign includes mucus impaction due to cystic fibrosis, allergic bronchopulmonary aspergillosis, broncholithiasis, foreign body aspiration and malignancies.

Yvonne Nussbaumer-Ochsner, Malcolm Kohler

Pulmonary Division, University Hospital of Zurich, Switzerland

Correspondence to Malcolm Kohler, Pulmonary Division, University Hospital of Zurich, Raemistrasse 100, CH-8091 Zurich, Switzerland; malcolm.kohler@usz.ch

Learning points

- The finger-in-glove sign is a radiographic feature and refers to mucoid impaction in central airways and typically radiates from the hilum to the periphery. Mucoid impaction is defined as airway filling by mucoid secretions.
- Physicians should be aware of the most frequent differential diagnoses of this radiographic sign which includes congenital (cystic fibrosis and congenital bronchial atresia) and acquired conditions (allergic bronchopulmonary aspergillosis, neoplastic conditions, broncholithiasis and foreign body aspiration).

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Accepted 16 February 2010

Published Online First 2 August 2010

Thorax 2011;66:182. doi:10.1136/thx.2009.130179

REFERENCES

1. **Ghaye B**, Szapiro D, Fanchamps J, *et al*. Congenital bronchial abnormalities revisited. *Radiographics* 2001;21:105–19.
2. **Martinez S**, Heyneman L, McAdams H, *et al*. Mucoid impactions: finger-in-glove sign and other CT and radiographic features. *Radiographics* 2008;28:1369–82.




Figure 1 (A) Posteroanterior radiograph showing a branching opacity in the left upper lung (finger-in-glove sign). (B) Axial CT image with tubular opacity and adjacent hyperlucency in the left upper lobe. (C) 3D-reconstruction of the bronchial tree. There is no division of corresponding bronchi, indicating atresia (see arrows).