

ORIGINAL ARTICLE

Poor air quality in classrooms related to asthma and rhinitis in primary schoolchildren of the French 6 Cities Study

Isabella Annesi-Maesano,^{1,2} Marion Hulin,^{1,2} François Lavaud,³ Chantal Raheison,^{4,5} Christine Kopferschmitt,⁶ Frederic de Blay,⁶ Denis André Charpin,⁷ Caillaud Denis⁸

► An additional material is published online only. To view this file please visit the journal online (<http://thorax.bmj.com/content/67/8.toc>).

¹EPAR, INSERM, Paris, France

²EPAR, Université Pierre et Marie Curie, Paris, France

³Hôpital Maison Blanche, Reims, France

⁴Hôpital du Haut-Lévêque, Bordeaux, France

⁵ISPED Université, Bordeaux, France

⁶Hôpitaux Civils, Strasbourg, France

⁷Hôpital Nord, Marseille, France

⁸Hôpital Montpied, Clermont-Ferrand, France

Correspondence to

Dr Isabella Annesi-Maesano, Epidemiology of Allergic and Respiratory Diseases (EPAR), UMR S 707, INSERM-UPMC-Paris 6, Medical School St Antoine, 27 rue Chaligny, Paris 75571 Cedex 12, France; isabella.annesimaesano@inserm.fr

Received 25 April 2011

Accepted 16 January 2012

Published Online First

21 March 2012

ABSTRACT

Background Relationships between indoor air quality (IAQ) found in schools and the allergic and respiratory health of schoolchildren have been insufficiently explored. A survey was conducted in a large sample of classrooms of primary schools in France to provide objective assessments of IAQ to which young schoolchildren are exposed in classrooms, and to relate exposure to major air pollutants found in classrooms to asthma and allergies of schoolchildren.

Methods Concentrations of fine particles with aerodynamic diameter $\leq 2.5 \mu\text{m}$ (PM_{2.5}), nitrogen dioxide (NO₂) and three aldehydes were objectively assessed in 401 randomly chosen classrooms in 108 primary schools attended by 6590 children (mean age 10.4 years, SD ± 0.7) in the French 6 Cities Study. The survey incorporated a medical visit including skin prick testing (SPT) for common allergens, a test for screening exercise-induced asthma (EIA) and a standardised health questionnaire completed by parents.

Results Children were differently exposed to poor air quality in classrooms, with almost 30% being highly exposed according to available standards. After adjusting for confounders, past year rhinoconjunctivitis was significantly associated with high levels of formaldehyde in classrooms (OR 1.19; 95% CI 1.04 to 1.36).

Additionally, an increased prevalence of past year asthma was found in the classrooms with high levels of PM_{2.5} (OR 1.21; 95% CI 1.05 to 1.39), acrolein (OR 1.22; 95% CI 1.09 to 1.38) and NO₂ (OR 1.16; 95% CI 0.95 to 1.41) compared with others. The relationship was observed mostly for allergic asthma as defined using SPT. A significant positive correlation was found between EIA and the levels of PM_{2.5} and acrolein in the same week.

Conclusions In this random sample, air quality in classrooms was poor, varied significantly among schools and cities, and was related to an increased prevalence of clinical manifestations of asthma and rhinitis in schoolchildren. Children with a background of allergies seemed at increased risk.

INTRODUCTION

Children are frail during their growth and because of their physical constitution and breathing pattern are more susceptible to the health effects of air pollution than adults.¹ In industrialised countries, children spend more time (up to 80%) indoors, mostly in schools, than in other places except home. In the USA, data from the Environmental

Key messages

What is the key question?

► How bad is air quality at school for children' respiratory health?

What is the bottom line?

► In our random sample, air quality in classrooms was poor, varied significantly among schools and cities.

► Almost 30% of school children were highly exposed according to WHO standards. Exposure to elevated concentrations of particulate matter and volatile organic compounds found in classrooms was significantly related to current asthma and rhino-conjunctivitis in schoolchildren.

► Children with a background of allergies were at increased risk.

Why read on?

► Our results are of relevance for public health because the population of exposed school children is huge due to the fact that primary school is compulsory and because childhood asthma and allergies are very common.

Protection Agency (<http://nces.ed.gov/surveys/frss/publications/2000032>) have shown that many schools have problems linked to poor indoor air quality (IAQ) with indoor air pollutant levels two to five times higher than outdoor levels. Indoor air pollutants can cause or contribute to short-term and long-term health problems, including nasal congestion, eye and skin irritations, coughing, sneezing, respiratory tract infection, allergic reactions, asthma, headaches, fatigue, dizziness and nausea.^{2,3} Moreover, indoor air pollutants can cause discomfort and reduce school attendance and productivity.⁴ However, data on air quality found in classrooms and its health effects are scant and inconsistent. Furthermore, few studies have used objective assessments of air quality and health indicators.⁵

To bridge the knowledge gap on the health effects of air pollution in children at school, we conducted a survey in a large representative sample of classrooms in six French cities. The survey protocol was intended to provide data on IAQ in primary schools in France objectively measured in



This paper is freely available online under the BMJ Journals unlocked scheme, see <http://thorax.bmj.com/site/about/unlocked.xhtml>

the frame of a field campaign, and on the relationships between major air pollutants found in the classrooms and asthma and rhinitis in schoolchildren using these classrooms. The air pollutants considered included fine particles with aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$), nitrogen dioxide (NO_2), formaldehyde, acetaldehyde and acrolein. In the analysis, various cofactors were taken into account as potential confounders.

METHODS

Population and survey

The 6 Cities Study (<http://www.epar.fr>) aimed to investigate the impact on childhood asthma and allergies of air pollutants objectively assessed in different settings, including schools in six cities (Bordeaux, Clermont-Ferrand, Créteil, Marseille, Strasbourg and Reims) chosen for heterogeneity of air pollution. Study participants were children aged 9–10 years who were invited to take part in a health survey according to a standardised protocol^{6–8} (for detailed information on the study protocol, see the online data supplement). Times of school visits for air quality assessment and medical examination of the children were randomly chosen.

Air quality

School visits were conducted during class time in the academic year. Concentrations of NO_2 , formaldehyde, acetaldehyde and acrolein were measured with passive diffusion samplers (Radiello, Padua, Italy). Concentrations of $\text{PM}_{2.5}$ were measured from Monday to Friday (night and day) using filter-based samples, charged with a pump, from representative points of the classroom. These measurements were made during the week when most of the children using the classroom underwent medical examination using standardised methodology.⁷ However, for logistic reasons, in some children the examination was realised beforehand or postponed. Analyses were performed in Padua for NO_2 and aldehyde samples, and in Paris for $\text{PM}_{2.5}$ filters. Methods, reliability and reproducibility of air pollution assessment have been presented elsewhere.^{7 9 10}

Health status

Parents gave written informed consent for their children to have a medical examination, including skin prick testing (SPT) to 10 common allergens and exercise-induced asthma (EIA) according to the standardised protocol of the run test.⁷ Parents also completed an enriched version of the International Study of Asthma and Allergies in Childhood (ISAAC) phase II questionnaire (for details, see the online data supplement). SPT positivity was defined as a weal at least 3 mm greater than that of the negative control for any of the allergens, 15 min after pricking. EIA was calculated using the formula $\text{PEF}_{\text{in}} - \text{PEF}_{\text{in}} / \text{PEF}_{\text{in}} \geq 10\%$ (PEF =peak expiratory flow).^{9–11} All steps of the study protocol received approval from the French National Ethics Board.

Studied variables

For each pollutant, a 5-day mean concentration ($\mu\text{g}/\text{m}^3$) in the classroom was computed and a three-class variable of exposure (high, medium or low) was defined with respect to the tertiles of the concentration distribution in the classroom, independent of the city. To investigate the impact of air quality in classrooms, health outcomes included EIA during the period of the survey and rhinoconjunctivitis and asthma in the past year ('sneezing and runny nose accompanied by itchy eyes out of cold in the past year' and 'ever had asthma'; having responded positively to at least one of these questions, 'wheezing or whistling in the

chest' or 'wheezing or whistling in the chest at night-time' or 'having taken asthma treatment' in the past year). Asthma was classified as allergic, when accompanied by SPT positivity to at least one allergen and non-allergic otherwise. Potential confounders assessed through the standardised ISAAC questions included age, gender, passive smoking, paternal or maternal history of asthma and allergic diseases, dampness, gas appliance, ethnicity and socio-economic status (SES) coded using the French Institut National de la Santé et des Etudes Economiques classification (<http://www.insee.fr>).

Statistical and epidemiological analyses

Between-school and within-school variability ('schools' variance and 'classrooms' variance, respectively) of the measured indoor pollutants were estimated using linear mixed models for longitudinal data (SAS MIXED procedure). Comparisons between groups for population characteristics were made using the Student t test for continuous variables and the Pearson χ^2 test or exact Fisher test for categorical variables. Pearson correlations were estimated between 5-day mean concentrations of the various air pollutants and the prevalence of EIA assessed during the same week of investigation. To investigate the relationship between each health outcome and each air pollutant, a marginal model was used,¹¹ taking into account the non-independence of data for children living in the same residential area (as defined by the school) and sharing the same environment (climate, pollens, social factors, food products, diet etc). This model calculated the adjusted OR for each health outcome at different levels of pollutant exposure. The parameters of the marginal model were estimated using the generalised estimating equation approach (SAS GENMOD) with independent working correlation structure using the city as stratum. These analyses were stratified using SPT positivity to investigate the role of atopic status. Association analyses were restricted to the subsample of schoolchildren who underwent a medical examination in the same week as air quality assessments were performed to diminish exposure misclassification (sensitivity analysis). SAS System for AIX V.8.2 was used for statistical analyses. A p value < 0.05 indicated statistical significance.

RESULTS

Overall, 7780 children out of 9615 who were initially selected participated in the study. Their mean age was 10.4 years. They originated from 401 classrooms of the 108 targeted schools in the six cities with the questionnaire filled out by the parents (response rate 81%). A total of 6683 school children (69%) completed the entire survey protocol. The analyses were limited to 6590 schoolchildren for whom all the variables of interest were available. Overall, 4643 schoolchildren underwent a medical examination during the same week that air quality assessments were performed (restricted sample, sensitivity analysis). Children of the restricted sample did not differ significantly from the others in terms of age, sex, weight, height and baseline peak flow, but they had more educated fathers (table 1). Approximately one in three children had at least one positive SPT (table 2). Only 31 children were positive to trophoallergens. Monosensitisation was observed in 11.2% of children and polysensitisation in 16.8%. Rhinoconjunctivitis was the most common condition, followed by EIA and asthma. As expected, allergic asthma was more frequent than non-allergic asthma.

Asthma and the environment

Table 1 Characteristics of the schoolchildren having participated in the French 6 Cities Study

Factors	All (n=6590)	Restricted sample (n=4643)*
Age (years) (mean±SD)	10.4±0.7	10.4±0.7
Sex (men) (%)	49.3	49.2
Weight (kg) (mean±SD)	36.2±8.2	36.3±8.2
Height (m) (mean±SD)	1.42±0.08	1.42±0.08
Body mass index (m/kg ²) (mean±SD)	17.9±2.9	17.9±2.9
Peak flow (ml/min) (mean±SD)	330.3±49.1	331.7±49.6
Parental history of allergy (%)	36.2	36.6
Exposure to tobacco smoke (%)	43.8	43.0

Body mass index: weight/height².

*Underwent medical examination during the same week as air quality assessments.

Air quality and health indicators

Air pollutants were variously distributed in the sample (figure 1). The lowest within-school (classrooms) variance was observed for acrolein (variance=2.55 µg/m³; SE=0.19 µg/m³; p<0.001) and the highest for formaldehyde (variance=183.52 µg/m³; SE=12.47 µg/m³; p<0.001) whereas the lowest between-school (schools) variance was observed for acrolein (variance=3.08 µg/m³; SE=0.53 µg/m³; p<0.001) and the highest for NO₂ (variance=107.26 µg/m³; SE=15.54 µg/m³; p<0.001) (for details, see the online data supplement). Overall, about one-third of the children were exposed to high concentrations of air pollutants according to standards indicated by WHO for PM_{2.5} (10 µg/m³) and NO₂ (40 µg/m³) (figure 2). In some classrooms, very high concentrations were found for formaldehyde, PM_{2.5} and NO₂. After adjusting for confounders, rhinoconjunctivitis was more frequent in classrooms with high formaldehyde concentrations (figure 3) and asthma was more common in classrooms with high PM_{2.5} and acrolein levels (figure 4). The results are consistent after accounting for SES and ethnicity in the marginal models. Stratifying the population according to SPT positivity showed that PM_{2.5}, acrolein and NO₂ were significantly related to allergic asthma (figure 5). In contrast, acrolein was negatively associated with non-allergic asthma (figure 6). Taking into account the time of year when the survey was conducted (spring and summer: 227 classrooms, 3609 schoolchildren; autumn and winter: 175 classrooms, 2981 schoolchildren), the following significant relationships were found: PM_{2.5} and asthma (OR 1.28, 95% CI 1.00 to 1.65), acrolein and asthma (OR 1.37, 95% CI 1.14 to 1.66) and PM_{2.5} and allergic asthma (OR 1.41, 95% CI

1.16 to 1.73) during the warm season; NO₂ and asthma (OR 1.18, 95% CI 1.01 to 1.39) and FA and rhinoconjunctivitis (OR 1.41, 95% CI 1.08 to 1.85) during the cold season. The sensitivity analysis confirmed previous results, although with a diminished statistical significance, probably because of the reduced sample size (data not shown). However, non-allergic asthma was found to be inversely related to acrolein (OR 0.36, 95% CI 0.29 to 0.44). The results were consistent after stratifying the restricted sample according to atopic status. Lastly, levels of PM_{2.5} and acrolein in classrooms correlated positively with the prevalence of EIA, which was assessed simultaneously (table 3).

DISCUSSION

In a random sample of primary schools, air quality in classrooms was poor and varied significantly among buildings and cities. Elevated levels of major pollutants were found to be related to a higher prevalence among schoolchildren of rhinoconjunctivitis and asthma in the year of the survey, and of EIA on the day of assessment. Children with a background of allergies were at increased risk.


In some classrooms, the mean concentrations of fine particles and NO₂ were higher than the threshold limits proposed for the control of health hazards by WHO (10 µg/m³ and 40 µg/m³ respectively). Elevated concentrations were also found for formaldehyde compared with the limits proposed by the French Agency for Environmental Safety (50 µg/m³ in the short term (2 h)), 10 µg/m³ in the long term (1 year)). Thirty per cent of schoolchildren were exposed to such elevated levels of air pollution in classrooms, which confirms previous observations. In 27 primary schools located in the city centre and suburbs of Antwerp, Belgium indoor PM_{2.5} and BTEX (benzene, toluene, ethylbenzene and xylenes) concentrations were elevated and exceeded the ambient concentrations, thus raising concerns about possible adverse health effects in susceptible children.⁹ In the cross-sectional EU-funded Health Effects of School Environment (HESE) Study conducted in five European countries, the proportions of schoolchildren highly exposed to PM₁₀ and CO₂ according to the standards for good IAQ (>50 µg/m³ and >1000 ppm, respectively) were 78% and 66%, respectively.¹⁰ This is in line with the report by the European Federation of Asthma and Allergy Associations, indicating that the right to breathe clean air in European schools is not encountered throughout Europe (<http://paginas.fe.up.pt>).

Table 2 Prevalence rates (%) of allergic sensitisation and allergic and respiratory morbidity in the French 6 Cities Study in the entire sample population (n=6590) and in the restricted sample of children who had a clinical examination during the same week as air quality assessments (n=4643)

Health outcome	All (n=6590)			Restricted sample (n=4643)				
	Boys	Girls	p Value	All	Boys	Girls	p Value	All
During air pollution assessment								
EIA	9.2	8.3	0.2269	8.7	10.0	8.5	0.0872	9.2
Past year (current)								
Rhinoconjunctivitis	13.1	10.7	0.0074	11.8		11.2	0.0840	12.1
Asthma	9.0	4.9	<0.0001	6.9	9.5	5.0	<0.0001	7.2
Allergic asthma	5.4	2.6	<0.0001	4.2	6.2	2.9	<0.0001	4.7
Non-allergic asthma	2.8	1.8	0.0111	2.3	2.8	1.5	0.0051	2.1
SPT positivity								
All allergens	32.7	23.8	<0.0001	28.0	33.3	25.3	<0.0001	29.2
Indoor allergens	24.5	17.6	<0.0001	21.0	22.4	18.7	<0.0001	22.0
Outdoor allergens	14.8	10.0	<0.0001	12.4	14.7	10.9	0.0002	12.8
Moulds allergens	3.7	1.8	<0.0001	2.7	3.8	1.7	<0.0001	2.7

EIA, exercise-induced asthma; SPT, skin prick test.

Figure 1 Concentration of indoor air pollutants in classrooms in the 6 Cities Study (N=408 classrooms). The median value is displayed as a line in each box, the mean as a cross, and box limits represent the 25th and 75th percentiles. The whiskers extend to the 5th and 95th percentile. NO₂, nitrogen dioxide; PM_{2.5}, fine particles with aerodynamic diameter ≤2.5 μm.




In this study, the alarming consequence of poor air quality in classrooms was a deterioration of respiratory health. A high prevalence of asthma in the past year was found in children using classrooms with 'high' pollution levels of PM_{2.5}, NO₂ and acrolein. Similarly, a high prevalence of rhinoconjunctivitis in the past year was found in children using classrooms with high levels of formaldehyde. In addition, schoolchildren experienced significantly more EIA during the run test when attending classrooms with elevated levels of PM_{2.5} and acrolein.

The association between air pollution in classrooms and respiratory health of schoolchildren was predictable¹² on the basis of previous findings obtained in experimental and epidemiological settings. This suggests that there is potential for air pollution to be a contributory factor to asthma and allergies, including allergic sensitisation and morbidity in some causal sequences.¹³ However, while evidence is accumulating on the adverse impact of outdoor air pollution exposure on respiratory health,¹⁴ few data exist on indoor air pollution and resulting


respiratory health effects,⁵ and even less in the case of school air quality. This lack of data is in spite of the fact that there is growing concern about the school environment, where children spend up to 8 h/day.^{15–19} Air pollutants have been found in classrooms^{20–22} that are known to be associated with deteriorating respiratory health in other indoor settings.⁵ In a subsample of the 6 Cities Study, exposure to acetaldehyde, toluene and formaldehyde at home was significantly associated with a higher risk of asthma.²³ Associations between respirable dust concentrations or amount of settled dust in the classroom with asthma symptoms, new onset of self-reported pet allergy and new onset of asthma diagnoses were reported in Swedish schoolchildren.^{18–24} In the HESE Study, PM₁₀ and CO₂ levels in a day of normal activity (full classroom) were related to dry cough at night and rhinitis in 654 children aged 10 years and to nasal patency in the subsample having performed this clinical test.¹⁰ All disorders were more prevalent in children attending poorly ventilated classrooms than in others. In our study, none

Figure 2 Exposure of schoolchildren in the 6 Cities Study the entire sample population (N=6590). Tertiles are defined according to the distribution of air pollutants in the classrooms (in μg/m³). Nitrogen dioxide (NO₂): low ≤23.7, medium >23.7 to ≤31.6, high >31.6; fine particles with aerodynamic diameter ≤2.5 μm (PM_{2.5}): low ≤12.2, medium >12.2 to ≤17.5, high >17.5; formaldehyde: low ≤19.1, medium >19.1 to ≤28.4, high >28.4; acetaldehyde: low ≤6.5, medium >6.5 to ≤9.9, high >9.9; acrolein: low=limit of detection (LD), medium >LD to ≤1.55, high >1.55.



Asthma and the environment

Figure 3 Relationships between air pollutants in classrooms and current rhinoconjunctivitis in the entire sample population of schoolchildren (N=6590) of the 6 Cities Study. Tertiles are defined according to the distribution of air pollutants in the classrooms (in $\mu\text{g}/\text{m}^3$). Nitrogen dioxide (NO_2): low ≤ 23.7 , medium >23.7 to ≤ 31.6 , high >31.6 ; fine particles with aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$): low ≤ 12.2 , medium >12.2 to ≤ 17.5 , high >17.5 ; formaldehyde: low ≤ 19.1 , medium >19.1 to ≤ 28.4 , high >28.4 ; acetaldehyde: low ≤ 6.5 , medium >6.5 to ≤ 9.9 , high >9.9 ; acrolein: low=limit of detection (LD), medium $>\text{LD}$ to ≤ 1.55 , high >1.55 . Dots are OR adjusted for age, gender, passive smoking and paternal or maternal history of asthma and allergic diseases; bars are 95% CIs.



of the classrooms were mechanically ventilated, which is common in French schools where ventilation is still provided by opening windows, so comparison is not possible with other studies. In 30 classrooms in 10 naturally ventilated schools in Shanghai, indoor NO_2 was associated with current asthma and asthma medication (OR 1.45 for $10 \mu\text{g}/\text{m}^3$; $p < 0.01$).²¹ For the first time, in our study we examined the effects of acrolein in classrooms. Interestingly, acrolein, which is a powerful irritant,²⁵ was significantly associated with EIA and past year asthma. However, it cannot be excluded that acrolein is a proxy of other indoor air pollutants that were not assessed in our study.

Our study on air pollution exposure was for a limited period of time and was conducted in late childhood, therefore it does not allow clarification of the role of air pollution in the development of asthma and allergies. However, it highlights the effect that school air quality may have on asthma and rhinitis

related morbidity. In our study, schoolchildren from classrooms with elevated levels of air pollutants experienced more clinical manifestations of asthma and rhinitis than others. This suggests that in the school setting, even in the absence of environmental smoking and other combustion sources, children may be equally exposed to air pollution. The fact that children attend school with associated environmental exposures early in life is also relevant because childhood is a particularly vulnerable period, when air pollution and other exposures are encountered.^{1 26} Other significant factors at this stage of life, but not investigated in our study, are air-pollutant-related disturbances observed in schoolchildren, such as learning impairment and poor cultural and social development.²⁷ As a consequence, good air quality at school has to be maintained so as not to expose children to an increased risk of these conditions.

Our study had several limitations. Although air pollution measurements were avoided during the summer, end-of-term

Figure 4 Relationships between air pollutants in classrooms and past year asthma in the entire sample population of schoolchildren (N=6590) of the 6 Cities Study. Tertiles are defined according to the distribution of air pollutants in the classrooms (in $\mu\text{g}/\text{m}^3$). Nitrogen dioxide (NO_2): low ≤ 23.7 , medium >23.7 to ≤ 31.6 , high >31.6 ; fine particles with aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$): low ≤ 12.2 , medium >12.2 to ≤ 17.5 , high >17.5 ; formaldehyde: low ≤ 19.1 , medium >19.1 to ≤ 28.4 , high >28.4 ; acetaldehyde: low ≤ 6.5 , medium >6.5 to ≤ 9.9 , high >9.9 ; acrolein: low=limit of detection (LD), medium $>\text{LD}$ to ≤ 1.55 , high >1.55 . Dots are OR adjusted for age, gender, passive smoking and paternal or maternal history of asthma and allergic diseases; bars are 95% CIs.





Figure 5 Relationships between air pollutants in classrooms and past year asthma among atopic children (N=1719) of the 6 Cities Study. Tertiles are defined according to the distribution of air pollutants in the classrooms (in $\mu\text{g}/\text{m}^3$). Nitrogen dioxide (NO_2): low ≤ 23.7 , medium >23.7 to ≤ 31.6 , high >31.6 ; fine particles with aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$): low ≤ 12.2 , medium >12.2 to ≤ 17.5 , high >17.5 ; formaldehyde: low ≤ 19.1 , medium >19.1 to ≤ 28.4 , high >28.4 ; acetaldehyde: low ≤ 6.5 , medium >6.5 to ≤ 9.9 , high >9.9 ; acrolein: low=limit of detection (LD), medium $>\text{LD}$ to ≤ 1.55 , high >1.55 . Dots are OR adjusted for age, gender, passive smoking and paternal or maternal history of asthma and allergic diseases; bars are 95% CIs.




vacations and at weekends, and the teachers were asked not to modify activities to reduce exposure misclassification, the 5-day measurements may not have been representative of children's usual exposure at school. In addition, the school environment could not have been the only factor responsible for the observed health effects because children might have been exposed to air pollution at home and outdoors. However, taking into account domestic sources of air pollution (environmental smoking, dampness, gas appliances) did not modify the relationship between air pollution at school and respiratory health in our study. The noxious effect of outdoor air pollutants including aldehydes has already been established in our population using various exposure assessments.⁶⁻⁸ Nonetheless, the indoors (classrooms)/outdoors (cities) ratio for aldehydes was higher than 1 (10), indicating that the levels of these pollutants were higher indoors than outdoors. This suggests that indoor exposure to formaldehydes and acrolein plays a role in the development of asthma and rhinitis. Health outcomes other than EIA

and allergic sensitisation were reported by parents using the ISAAC questionnaire, which is a robust measure. Nevertheless, using these reports in the absence of objective validation could lead to misclassification of asthma and rhinitis. The definition of asthma used in this study, which combined ever asthma and current symptoms, could also have influenced the results by excluding children with asthma who were not aware of their condition because they had not received a diagnosis. However, figures for current asthma-like symptoms in the absence of ever asthma showed similar trends. Finally, due to the cross-sectional study design and the extent of air pollution measurements, our data do not allow causal relationships between air pollution and asthma and rhinitis to be established.

The strengths of our study are the sampling design, the large number of schools and classrooms, the objective assessments of various air pollutants and the sensitivity analysis, which was restricted to children for whom clinical examination and air pollution assessment were conducted simultaneously,

Figure 6 Relationships between air pollutants in classrooms and past year asthma among non-atopic children (N=4377) of the 6 Cities Study. Tertiles are defined according to the distribution of air pollutants in the classrooms (in $\mu\text{g}/\text{m}^3$). Nitrogen dioxide (NO_2): low ≤ 23.7 , medium >23.7 to ≤ 31.6 , high >31.6 ; fine particles with aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$): low ≤ 12.2 , medium >12.2 to ≤ 17.5 , high >17.5 ; formaldehyde: low ≤ 19.1 , medium >19.1 to ≤ 28.4 , high >28.4 ; acetaldehyde: low ≤ 6.5 , medium >6.5 to ≤ 9.9 , high >9.9 ; acrolein: low=limit of detection (LD), medium $>\text{LD}$ to ≤ 1.55 , high >1.55 . Dots are OR adjusted for age, gender, passive smoking and paternal or maternal history of asthma and allergic diseases; bars are 95% CIs.



Asthma and the environment

Table 3 Correlations between concentrations of air pollution and exercise-induced asthma (EIA) in the sample of children who had a clinical examination during the same week as air quality assessments (n=4643)

Air pollutant	n	Correlation coefficient*	p Value
NO ₂	4254	0.022	0.1555
PM _{2.5}	4051	0.061	<0.001
Formaldehyde	4374	-0.018	0.2257
Acetaldehyde	4374	-0.025	0.0945
Acrolein	3551	0.038	<0.0253

*When less than 10 children were examined by the doctor in a classroom during a week, the median value of the concentrations of pollutants measured during the same week and the mean prevalence of EIA in the entire school were taken.

NO₂, nitrogen dioxide; PM_{2.5}, fine particles with aerodynamic diameter ≤2.5 µm.

confirming the observed associations. Restricting the analysis to these children not only reduced exposure misclassification bias but was also appropriate to investigate short-term effects of air pollution on EIA. Although small, the significant correlations between EIA and PM_{2.5} and acrolein represent an important message for public health as between 40% and 90% of people with asthma usually have EIA and could be at even higher risk when exposed to air pollution. In our population, asthma and EIA were highly related (OR 10.49, 95% CI 7.31 to 15.05).

In conclusion, poor air quality was found in randomly chosen classrooms of primary schools in metropolitan France and was related to an increased prevalence of clinical manifestations of asthma and rhinitis among schoolchildren. Whether and how exposure at school plays a role in the development of asthma and allergies needs further investigation, taking into account individual daily activity patterns.

Acknowledgements We are particularly indebted to pupils and parents without whom the study would not have been possible. We thank Yvon Le Moulec and Anne-Marie Laurent from Laboratoire d'Hygiène de la Ville de Paris, who centralised air pollution assessments and took charge of PM assessments, and Gilles Perron, Joseph Kleinpeter and Alain Target from ASPA in Strasbourg, who assessed the reproducibility of air pollution assessment. We thank David Moreau from the EPAR Department, UMR-S 707, Paris who created the database and contributed to the statistical analysis, and Soutrik Banerjee from the EPAR Department, UMR-S 707, Paris who also contributed to the statistical analysis. We thank the study field team and staff at the participating air quality districts in the six cities, who helped to establish the quality of air in schools. We also thank Vincenzo Cocheo and Paolo Sacco (Fondazione Maugeri, Italy) who helped in air pollutants assessment through passive samplers, the Education Nationale, the school doctors, school principals and teachers. We thank Professor Mercier from Comité Contre la Tuberculose et les Maladies Respiratoires, on the aegis of which the survey was conducted. Allergen extracts for skin prick testing were kindly provided by Stallergènes Laboratoires (Antony, France).

Contributors IAM is the PI of the 6 Cities Study. All the authors but MH implemented the rationale of the study and conducted the survey in their city. IAM and MH conducted the statistical analysis presented in the paper. IAM wrote the paper. All the authors contributed to the study and worked on the paper.

Funding The French 6 Cities Study was supported by National Institute for Health and Medical Research (INSERM) (Programme Déterminants de la Santé), Ministry of Health (DGS), Environmental Programme PRIMEQUAL-PREDIT, Agence de la Maîtrise de l'Energie (ADEME), Mutuelle Générale de l'Education Nationale (MGEN), Agence Française de Sécurité Sanitaire de l'Environnement et du Travail (AFSSET) and ANTADIR.

Competing interests None.

Patient consent obtained.

Ethics approval French CCPPRB.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The 6 Cities Study should allow investigation of the relationship between stress, air pollution, asthma and allergies.

REFERENCES

1. Annesi-Maesano I, Agabiti N, Pistelli R, *et al*. Subpopulations at increased risk of adverse health outcomes from air pollution. *Eur Respir J Suppl* 2003;**40**: 57s–63s.
2. Geller RJ, Rubin IL, Nodvin JT, *et al*. Safe and healthy school environments. *Pediatr Clin North Am* 2007;**54**:351–73, ix.
3. Bernstein JA, Alexis N, Bacchus H, *et al*. The health effects of non-industrial indoor air pollution. *J Allergy Clin Immunol* 2008;**121**:585–91.
4. Mendell MJ, Heath GA. Do indoor pollutants and thermal conditions in schools influence student performance? A critical review of the literature. *Indoor Air* 2005;**15**:27–52.
5. Viegi G, Simoni M, Scognamiglio A, *et al*. Indoor air pollution and airway disease. *Int J Tuberc Lung Dis* 2004;**8**:1401–15.
6. Penard-Morand C, Charpin D, Raheison C, *et al*. Long-term exposure to background air pollution related to respiratory and allergic health in schoolchildren. *Clin Exp Allergy* 2005;**35**:1279–87.
7. Annesi-Maesano I, Moreau D, Caillaud D, *et al*. Residential proximity fine particles related to allergic sensitisation and asthma in primary school children. *Respir Med* 2007;**101**:1721–9.
8. Penard-Morand C, Raheison C, Charpin D, *et al*. Long-term exposure to close-proximity air pollution and asthma and allergies in urban children. *Eur Respir J* 2010;**36**:33–40.
9. Stranger M, Potgieter-Vermaak SS, Van Grieken R. Characterization of indoor air quality in primary schools in Antwerp, Belgium. *Indoor Air* 2008;**18**:454–63.
10. Simoni M, Annesi-Maesano I, Sigsgaard T, *et al*. School air quality related to dry cough, rhinitis and nasal patency in children. *Eur Respir J* 2010;**35**:742–9.
11. Liang H, Wang S, Carroll RJ. Partially linear models with missing response variables and error-prone covariates. *Biometrika* 2007;**94**:185–98.
12. Susser M. What is a cause and how do we know one? A grammar for pragmatic epidemiology. *Am J Epidemiol* 1991;**133**:635–48.
13. Penard-Morand C, Annesi-Maesano I. Allergic respiratory diseases and outdoor air pollution. *Rev Mal Respir* 2008;**25**:1013–26.
14. Holguin F. Traffic, outdoor air pollution, and asthma. *Immunol Allergy Clin North Am* 2008;**28**:577–88, viii–ix.
15. Berge M, Munir AK, Dreborg S. Concentrations of cat (Fel d1), dog (Can f1) and mite (Der f1 and Der p1) allergens in the clothing and school environment of Swedish schoolchildren with and without pets at home. *Pediatr Allergy Immunol* 1998;**9**:25–30.
16. Smedje G, Norback D, Edling C. Asthma among secondary schoolchildren in relation to the school environment. *Clin Exp Allergy* 1997;**27**:1270–8.
17. Taskinen T, Meklin T, Nousiainen M, *et al*. Moisture and mould problems in schools and respiratory manifestations in schoolchildren: clinical and skin test findings. *Acta Paediatr* 1997;**86**:1181–7.
18. Kim JL, Elfman L, Mi Y, *et al*. Current asthma and respiratory symptoms among pupils in relation to dietary factors and allergens in the school environment. *Indoor Air* 2005;**15**:170–82.
19. Daisey JM, Angell WJ, Apte MG. Indoor air quality, ventilation and health symptoms in schools: an analysis of existing information. *Indoor Air* 2003;**13**:53–64.
20. Tortolero SR, Bartholomew LK, Tyrrell S, *et al*. Environmental allergens and irritants in schools: a focus on asthma. *J Sch Health* 2002;**72**:33–8.
21. Mi YH, Norbäck D, Tao J, *et al*. Current asthma and respiratory symptoms among pupils in Shanghai, China: influence of building ventilation, nitrogen dioxide, ozone, and formaldehyde in classrooms. *Indoor Air* 2006;**16**:454–64.
22. Fox A, Harley W, Feigley C, *et al*. Increased levels of bacterial markers and CO₂ in occupied school rooms. *J Environ Monit* 2003;**5**:246–52.
23. Hulín M, Caillaud D, Annesi-Maesano I. Indoor air pollution and childhood asthma: variations between urban and rural areas. *Indoor Air* 2010;**20**:502–14.
24. Kim JL, Elfman L, Mi Y, *et al*. Indoor molds, bacteria, microbial volatile organic compounds and plasticizers in schools—associations with asthma and respiratory symptoms in pupils. *Indoor Air* 2007;**17**:153–63.
25. Bessac BF, Jordt SE. Sensory detection and responses to toxic gases: mechanisms, health effects, and countermeasures. *Proc Am Thorac Soc* 2010;**7**:269–77.
26. Stebbings JH Jr, Fogleman DG. Identifying a susceptible subgroup: effects of the Pittsburgh air pollution episode upon schoolchildren. *Am J Epidemiol* 1979;**110**:27–40.
27. Moglia D, Smith A, MacIntosh DL, *et al*. Prevalence and implementation of IAQ programs in U.S. schools. *Environ Health Perspect* 2006;**114**:141–6.

THORAX

Poor air quality in classrooms related to asthma and rhinitis in primary schoolchildren of the French 6 Cities Study

Isabella Annesi-Maesano, Marion Hulin, François Lavaud, Chantal Raheison, Christine Kopferschmitt, Frederic de Blay, Denis André Charpin and Caillaud Denis

Thorax 2012 67: 682-688 originally published online March 21, 2012
doi: 10.1136/thoraxjnl-2011-200391

Updated information and services can be found at:
<http://thorax.bmj.com/content/67/8/682>

These include:

Supplementary Material Supplementary material can be found at:
<http://thorax.bmj.com/content/suppl/2012/03/20/thoraxjnl-2011-200391.DC1>

References This article cites 27 articles, 2 of which you can access for free at:
<http://thorax.bmj.com/content/67/8/682#BIBL>

Open Access This is an open-access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license. See: <http://creativecommons.org/licenses/by-nc/2.0/> and <http://creativecommons.org/licenses/by-nc/2.0/legalcode>.

Email alerting service Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

- [Open access](#) (260)
- [Asthma](#) (1782)
- [Child health](#) (843)
- [Ear, nose and throat/otolaryngology](#) (218)
- [TB and other respiratory infections](#) (1273)
- [Air pollution](#) (110)
- [Environmental issues](#) (253)
- [Screening \(epidemiology\)](#) (366)
- [Screening \(public health\)](#) (366)

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>