

S116 **MICRORNA REGULATION OF TOLL-LIKE RECEPTOR 7 FUNCTION IN SEVERE ASTHMA: RELEVANCE TO VIRAL RESPONSES**

doi:10.1136/thoraxjnl-2012-202678.121

H Rupani, P Dennison, N Jayasekera, T Sanchez-Elsner, P H Howarth. *Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton, Southampton, United Kingdom*

Background and Aims Viral-induced disease exacerbation is common in asthma and studies have identified that both bronchial epithelial cells and alveolar macrophages (AM) from asthmatics have a reduced interferon (IFN) response to rhinovirus infection. The mechanism behind this defect is unclear. As asthmatic peripheral blood mononuclear cells have been reported to have defective toll-like receptor (TLR) 7 function, we investigated the expression of microRNAs (miRNAs) in AM from healthy control (HC) and severe asthma (SA) volunteers with relevance to TLR7-viral interactions. MicroRNAs are non-coding RNAs that down-regulate gene expression by suppressing translation. We identified and focused on 3 miRNAs that could target TLR7. Additionally, we investigated if manipulating the expression of these miRNAs can ameliorate the defective IFN response in AM.

Methods 26 HC and 30 patients with SA (BTS Step 4/5) were recruited for bronchoscopy. AM were isolated from bronchoalveolar lavage using the adherence to plastic technique. Expression of miRNAs and TLR7 was determined by qRT-PCR and western blotting. AM were transfected with a combination of antagomirs, specifically directed against the 3 miRNAs, and then treated with imiquimod (5ug/ml), a TLR7 agonist, or human rhinovirus-16 (HRV16) and IFN- β expression was determined after 24 hours using qRT-PCR and ELISA.

Results Expression of all three miRNAs was significantly increased in SA compared to HC. TLR7 mRNA was found to be significantly reduced in AM from volunteers with SA compared to HC. Western blotting confirmed reduced expression of TLR7 protein in AM from SA compared to HC. Compared to mock transfected AM, AM transfected with the 3 antagomirs showed significantly increased imiquimod-induced IFN- β mRNA and protein expression and significantly increased HRV16-induced IFN- β mRNA production.

Conclusion TLR7 expression is significantly reduced in SA compared to HC. The differential expression of the miRNAs identified may lead to impaired viral sensing by asthmatic AM and contribute to the defective IFN response to rhinovirus. Importantly, TLR7 induced IFN- β production by human AM can be significantly augmented by inhibition of these miRNAs. The identification of these miRNAs and our ability to manipulate their expression in human AM offers the potential for future miRNA-based therapies in asthma.

S117 **EFFECTS OF HYPOXIA ON EOSINOPHIL APOPTOSIS, EFFEROCYTOSIS AND SENSITIVITY TO GLUCOCORTICOSTEROIDS**

doi:10.1136/thoraxjnl-2012-202678.122

L Porter, AS Cowburn, N Farahi, CA Fiddler, AM Condliffe, ER Chilvers. *University of Cambridge, Cambridge, England*

Many tissues including the skin, intestinal epithelium and potentially the airway operate at 'physiological' levels of tissue hypoxia with normal PO_2 values below 3KPa. Both sterile and non-sterile inflammation exacerbates the degree of tissue hypoxia and predicates the need for granulocytes including eosinophils to operate efficiently under hypoxia. In these experiments we have examined the effects of hypoxia on eosinophil longevity and show that a PO_2 below 3KPa severely attenuates the pro-apoptotic effect of dexamethasone.

Human blood eosinophils were prepared from healthy donors using hetastarch-sedimentation and EasySep[®]-immunomagnetic beads, and cultured in RPMI + 10% autologous serum for 6–24 h. Apoptosis and efferocytosis were quantified using standard morphology, AnV/PI staining and myeloperoxidase counter-stain methods. Hypoxic incubation (H) (media PO_2 2.9 ± 0.1 KPa) caused a marked survival response in eosinophils compared to normoxia (N) (% apoptosis at 24 h: N $16.3 \pm 3.0\%$; H $1.2 \pm 0.2\%$, $n=5$), which was of similar magnitude to that observed with IL-5 (N $4.2 \pm 0.3\%$). This hypoxic survival effect was mimicked by the iron chelator/2-oxyglutarate analogue DFO (10 mM) and DMOG (1 mM) and blocked in a concentration-dependent manner by the protein synthesis inhibitor cycloheximide. In preliminary experiments hypoxia also independently reduced the capacity of monocyte-derived macrophages to phagocytose apoptotic eosinophils as seen previously with neutrophils. Most strikingly, hypoxic incubation also reduced the normal pro-apoptotic effect of dexamethasone (1 μ M) (% apoptosis at 24 h: N + Dex $27.6 \pm 3.9\%$, H + Dex $11.2 \pm 2.9\%$). qPCR analysis of the glucocorticosteroid-dependent gene *GILZ* and the hypoxia-HIF-1 α -dependent gene *GLUT1* demonstrated that the above suppression of dex-induced eosinophil apoptosis was not a consequence of the inhibition of steroid-induced transcriptional activity (mRNA fold change at 16 h: *GILZ* N 12.89 ± 3.6 , *GILZ* H 23.9 ± 9.2 ; *GLUT1* H (control) 28.5 ± 7.2 , *GLUT1* H (Dex) 45.5 ± 13.5 , $n=3$).

These data suggest that hypoxia can cause a profound effect on eosinophil longevity and macrophage uptake, and render these cells partially resistant to the pro-apoptotic effects of dexamethasone. This may impede the effective resolution of eosinophilic inflammation *in vivo*.

S118 **IL-13 INDUCED MOUSE AIRWAY INFLAMMATION INDUCES AN INCREASE OF SOLUBLE ADAM33 IN BRONCHOALVEOLAR LAVAGE FLUID, WHICH IS ENZYMATICALLY ACTIVE AND ASSOCIATED WITH BRONCHIAL HYPERRESPONSIVENESS**

doi:10.1136/thoraxjnl-2012-202678.123

^{1,2}ER Davies, ²G Chen, ¹ST Holgate, ¹DE Davies, ²JA Whitsett, ^{1,2}HM Haitchi. ¹*Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton, Southampton, UK*; ²*Division of Pulmonary Biology, Cincinnati children's Hospital Medical Center and University of Cincinnati College of Medicine, Cincinnati, USA*

Rationale The asthma susceptibility gene *ADAM33/Adam33* is associated with bronchial hyperresponsiveness (BHR) in humans and mice. Soluble ADAM33 is increased in bronchoalveolar lavage fluid (BALF) of allergic asthma patients (Lee JY et al, *AJRCCM* 2006 Apr; 173(7):729–35). Its levels correlate with declining FEV₁%, suggesting a role in airway remodelling in asthma. Maternal allergy oregonous IL-13 suppresses *Adam33/ADAM33* mRNA expression but enhances ADAM33 protein processing in human embryonic and juvenile mouse lungs (Haitchi HM et al, *JACI*. 2009 Sep; 124(3):590–7, 597). We hypothesise that conditional expression of IL-13 in mouse lungs induces the enzymatically active, soluble form of ADAM33 in BALF, which is associated with BHR.

Methods IL-13 expression was induced using Doxycycline in CCSP-rtTA/Otet-IL-13 double-transgenic (dTg) mice. Methacholine challenge and lung function measurements were performed and lungs were harvested for mRNA extraction and immunohistochemistry (IHC). BALF was obtained for Western-blotting for ADAM33 and testing of ADAM33 enzymatic activity using a fluorescence resonance energy transfer (FRET) peptide assay.

Results There was a significant increase in BHR to Methacholine in IL-13 expressing double transgenic mice. IHC showed an increase in bronchial smooth muscle in lungs of double transgenic mice. Similar to the RT-qPCR findings in human embryonic and juvenile mouse

lungs, IL-13 suppressed *Adam33* mRNA but no difference in *α-smooth muscle actin* (*αSma*) was evident. Immunoblotting for ADAMA33 in BALF demonstrated a 76kDa band, consistent with the ADAMA33 ectodomain and processed forms at 38/44kDa in dTg animals. ADAMA33 enzymatic activity was also significantly increased.

Conclusion The data suggest that allergic inflammation induced by IL-13 suppresses *Adam33* mRNA expression but induces the release of soluble forms of ADAMA33, yielding enzymatically active forms. The release of soluble forms may play a role in airway remodelling, potentially leading to BHR. We next propose to test the effect of specific ADAMA33 inhibitors on airway remodelling in this allergic mouse model to assess their potential as novel treatments for asthma.

S119 ROLES OF TLR3, TLR4- AND TLR5-7-9 IN INTERFERON INDUCTION IN BRONCHIAL EPITHELIAL CELLS AND PERIPHERAL BLOOD MONONUCLEAR CELLS FROM ASTHMATIC AND NON-ASTHMATIC SUBJECTS

doi:10.1136/thoraxjnl-2012-202678.124

¹A Sykes, ²MR Edwards, ¹J Macintyre, ¹A del Rosario, ¹OM Kon, ¹SL Johnston. ¹Imperial College London and Imperial Healthcare NHS Trust, London, UK; ²Imperial College London, London, UK

Introduction Defective rhinovirus (RV) induced interferon (IFN)- β and IFN- λ production has been reported in primary human bronchial epithelial cells (HBECs) and peripheral blood mononuclear cells (PBMCs) from asthmatics. The mechanisms of defective IFN induction in asthma are unknown. Virus infection can induce IFNs through Toll like Receptors (TLR)3, TLR4 and TLRs7-9 and TLR agonists have been identified as potential therapeutic options for asthma. The role of these TLRs in IFN induction in asthma is unclear.

Objective To investigate IFN responses to TLR stimulation in HBECs and PBMCs from atopic asthmatic and non-asthmatic individuals.

Methods HBECs and PBMCs from atopic asthmatic and non-asthmatic subjects were stimulated with agonists to TLR3, TLR4 & TLRs7-9 and type I and III IFN responses assessed by qPCR and ELISA.

Results TLR3 and TLR7, but not TLR4, 8 or 9, stimulation induced IFN protein and mRNA expression in HBECs and PBMCs. IFNs induced were IFN- β and predominantly type III IFN- λ in HBECs and type I ($-\alpha$ and $-\beta$) with no IFN- λ in PBMCs. TLR function was not defective in asthmatic compared to non-asthmatic subjects.

Conclusions TLR3 & TLR7 were the predominant TLRs involved in IFN induction in HBECs and PBMCs. Defective IFN induction to TLR agonists was not observed in these well controlled asthmatic subjects. TLR3/7 agonists could be effective in inducing IFNs in more severe/less well controlled asthmatics who may have deficient virus induced IFN production.

Evaluation and treatment of Cystic Fibrosis

S120 UPDATE ON THE UK CF GENE THERAPY CONSORTIUM MULTIDOSE, NON-VIRAL, GENE THERAPY TRIAL

doi:10.1136/thoraxjnl-2012-202678.125

¹EWFW Alton, ¹D Ashby, ²C Boyd, ³S Cheng, ⁴S Cunningham, ¹JC Davies¹, ⁵D Gill, ¹U Griesenbach, ¹T Higgins, ⁵S Hyde, ⁶JA Innes, ²G Murray, ²D Porteous. ¹Imperial College, London, UK; ²University of Edinburgh, UK; ³Genzyme Corp, Framingham, USA; ⁴Royal Hospital for Sick Children, Edinburgh, UK; ⁵University of Oxford, UK; ⁶Western General Hospital, Edinburgh, UK

The UK CF GTC has been working for several years to determine the clinical benefit of *CFTR* gene therapy. Our premise was that for such a therapy to achieve clinical benefit, repeated administration

would be required, and that therefore a non-viral approach was needed. We demonstrated in laboratory and preclinical models that GL67A (Genzyme Corp) was the optimal gene transfer agent, and designed a plasmid, pGM169, completely depleted of pro-inflammatory CpG motifs and driven by the novel hCEFI promoter, designed for prolonged expression. In a longitudinal observational study (the Run-in) we measured the variability of multiple outcome measures, both conventional and novel. These data have allowed us to perform power calculations and a) choose our primary outcome (FEV₁), b) secondary efficacy outcomes (lung clearance index, various parameters on CT scan, Quality of life questionnaire [CFQ-R], exercise capacity and activity, and selected sputum and serum inflammatory markers), and c) safety measures (clinical findings, exacerbation rate, gas transfer, sputum culture, serum inflammatory markers, renal and hepatic markers). We have recently completed a single-dose safety and dose ranging study.

In this trial, 130 patients, aged 12 years and above are being randomised in a 1:1 fashion to active treatment or placebo and will receive the nebulised agent at monthly intervals for 12 doses. The group size was determined on the basis of a 6% relative improvement in FEV₁. An adaptive design will be used for additional safety; the first 20 patients will receive 3 doses ahead of the rest of the cohort. Patients will be invited to participate in either one or two substudies, being conducted to explore mechanisms; a) nasal administration followed by nasal potential difference (PD) and brushings for mRNA expression and b) pre and post-treatment bronchoscopies for lower airway PD, gene expression and histology. The double-blinded nature of the trial means that final outcome data will only be available upon completion of the study. The trial was initiated in April 2012; here we will update on recruitment, projected time-lines and progress.

Funded by the MRC and NIHR through the EME programme.

S121 LUNG CLEARANCE INDEX TO EVALUATE THE EFFECT OF IVACAFTOR ON LUNG FUNCTION IN SUBJECTS WITH CF WHO HAVE THE G551D-CFTR MUTATION AND MILD LUNG DISEASE

doi:10.1136/thoraxjnl-2012-202678.126

¹JCD Davies, ²HS Sheridan, ³PSL Lee, ³TS Song, ³AS Stone, ⁴FR Ratjen. ¹Royal Brompton Hospital and Imperial College, London, UK; ²University of Edinburgh, Edinburgh, UK; ³Vertex Pharmaceuticals Incorporated, Cambridge, USA; ⁴The Hospital for Sick Children, Toronto, Canada

Background Ivacaftor has been shown to lead to significant improvement in lung function, exacerbation rate, weight gain and quality of life in adolescents and adults with CF and the *G551D-CFTR* mutation.

Objectives Drugs targeting the basic defect of CF may hold potential for patients with early stage disease, but establishing benefit is more difficult. If FEV₁ is normal, a more sensitive test may be useful. LCI has been shown to become abnormal at an earlier stage of disease than FEV₁ and thus may be a more sensitive outcome measure in this group of patients.

Methods This Phase 2, randomised, double-blind, placebo-controlled, multicenter, crossover study evaluated the effect of ivacaftor on LCI derived from multibreath washout of SF₆ using an Innocor device. Subjects were ≥ 6 years with the *G551D-CFTR* mutation, FEV₁ >90% predicted, and LCI >7.4 (upper limit of normal). Ivacaftor 150 mg or placebo was administered twice daily for two 4-week periods with a 4-week washout in between.

Results Twenty-one subjects were randomised and 20 received a dose of ivacaftor. Seventeen subjects completed both periods. Mean (SD) age was 16.6 (10.9) years. Mean (SD) baseline LCI was 9.0 (1.5). The treatment effect of ivacaftor for adjusted mean change from baseline in LCI at Day 29 was -2.1 ($P=0.0004$). Mean