

A systematic review and meta-analysis of the efficacy and safety of *N*-acetylcysteine in preventing aminoglycoside-induced ototoxicity: implications for the treatment of multidrug-resistant TB

Katharina Kranzer,^{1,2} Wael F Elamin,¹ Helen Cox,³ James A Seddon,⁴ Nathan Ford,⁵ Francis Drobniowski^{1,6}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2015-207245>).

¹National Mycobacterium Reference Laboratory, Public Health England, London, UK

²Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK

³Division of Medical Microbiology and Institute of Infectious Diseases and Molecular Medicine, University of Cape Town, Cape Town, South Africa

⁴Department of Paediatric Infectious Diseases, Imperial College London, St Mary's Hospital, London, UK

⁵Centre for Infectious Disease Epidemiology and Research, University of Cape Town, Cape Town, South Africa

⁶Department of Infectious Diseases, Imperial College, The Hammersmith Campus, London, UK

Correspondence to

Dr Katharina Kranzer, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK; katharina.kranzer@lshtm.ac.uk

Received 7 May 2015

Accepted 18 June 2015

Published Online First

7 September 2015



CrossMark

To cite: Kranzer K, Elamin WF, Cox H, et al. *Thorax* 2015;**70**: 1070–1077.

ABSTRACT

Background Ototoxicity is a severe side effect of aminoglycoside antibiotics. Aminoglycosides are recommended for the treatment of multidrug-resistant TB (MDR-TB). *N*-Acetylcysteine (NAC) appears to protect against drug- and noise-induced hearing loss. This review aimed to determine if coadministering NAC with aminoglycoside affected ototoxicity development, and to assess the safety and tolerability of prolonged NAC administration.

Methods Eligible studies reported on the efficacy of concomitant NAC and aminoglycoside administration for ototoxicity prevention or long-term (≥ 6 weeks) administration of NAC regardless of indication. Pooled estimates were calculated using a fixed-effects model. Heterogeneity was assessed using the I^2 statistic.

Results Three studies reported that NAC reduced ototoxicity in 146 patients with end-stage renal failure receiving aminoglycosides. Pooled relative risk for otoprotection at 4–6 weeks was 0.14 (95% CI 0.05 to 0.45), and the risk difference was –33.3% (95% CI 45.5% to 21.2%). Eighty-three studies (N=9988) described the administration of NAC for > 6 weeks. Abdominal pain, nausea and vomiting, diarrhoea and arthralgia were increased 1.4–2.2 times.

Discussion This review provides evidence for the safety and otoprotective effect of NAC when coadministered with aminoglycoside. It represents a strong justification for a clinical trial to investigate the effect of concomitant NAC treatment in patients receiving aminoglycosides as part of MDR-TB treatment.

INTRODUCTION

Ototoxicity is a potentially severe side effect of aminoglycoside antibiotics. Aminoglycosides induce apoptosis of the inner and outer hair cells—the auditory and vestibular sensory receptors within the cochlea. This apoptosis is mediated by disruption of mitochondrial protein synthesis with the subsequent generation of free radicals.^{1,2} As the sensory epithelium of the mammalian cochlea has little regenerative capacity,³ this apoptosis leads to irreversible loss of hearing and balance.⁴ Hearing loss, which mainly affects high-frequency tones, may progress even after discontinuation of the drug because of the accumulation of free radicals and is irreversible.⁵

Key messages

What is the key question?

- Does coadministration of *N*-acetylcysteine (NAC) with aminoglycosides prevent the development of ototoxicity and is it safe?

What is the bottom line?

- Coadministration of NAC reduces the risk of ototoxicity by 80% and was found to be safe.

Why read on?

- NAC may be an effective strategy to reduce ototoxicity in patients treated with aminoglycoside in the context of multidrug-resistant TB.

Multidrug-resistant (MDR)-TB is defined as resistance to isoniazid and rifampicin, with or without resistance to other anti-TB drugs. The second-line injectable drugs, including the aminoglycosides (amikacin and kanamycin) and the polypeptides (capreomycin), are among the main anti-TB antibiotics used for the treatment of MDR-TB, with a recommended minimum treatment duration of 8 months.⁶ If aminoglycoside-induced hearing loss is detected early, through systematic and regular audiological examination, it may be possible to intervene before the hearing loss reaches the frequencies that might affect communication (mid- and low-frequency ranges). However, audiology assessment is often inadequate in both resource-limited and -rich settings, and even regular assessment may not be timely enough to prevent rapid hearing loss for some patients. In addition, the incidence of, and risk factors for, ototoxicity in patients treated for MDR-TB remain poorly characterised. A recent systematic review identified 35 studies reporting the frequency of ototoxicity in patients receiving MDR-TB treatment, but the majority (86%) of these studies failed to specify the testing and classification methods used. In the five studies that used standardised testing and classification methods, the frequency of ototoxicity ranged from 18% to 62%.⁷

There are limited interventions available to prevent or ameliorate hearing loss in patients



receiving second-line injectable drug treatment for MDR-TB. Streptomycin (an aminoglycoside previously used in retreatment of TB but now rarely used) and capreomycin are thought to be less ototoxic than amikacin or kanamycin,^{8–9} but may be less efficacious.¹⁰ Increasing the dose interval to thrice weekly rather than daily has not been shown to have any impact on ototoxicity.⁸ Although therapeutic drug monitoring is recommended for amikacin and streptomycin, this is not readily available for capreomycin and kanamycin, particularly in low-resource settings where the majority of MDR-TB is managed. The main options that can be used to prevent the progression of hearing loss once it has been detected include stopping the drug, reducing the dose, or increasing the dose interval. However, none of these strategies has been systematically evaluated, and, to date, evidence of any benefit of alteration in dose and interval is lacking. Furthermore these options may reduce treatment efficacy and lower the chance of cure through compromising the regimen.

As the cochlear hair cell damage is caused by reactive oxygen species, it is theoretically possible to mitigate these effects by coadministration of antioxidants.^{11–12} Aspirin, an established antioxidant, has been shown to protect against hearing loss in adults treated with gentamicin.¹³ More recently, several studies in patients undergoing dialysis have shown a protective auditory effect of *N*-acetylcysteine (NAC) when coadministered with either gentamicin or amikacin.^{14–17} NAC, a thiol-containing antioxidant, is a successful and established treatment which ameliorates hepatic and renal toxicity in acetaminophen (paracetamol) overdose and contrast-induced kidney injury.^{18–19} Moreover, NAC has been used in both animals and humans to reduce cisplatin- and noise-induced ototoxicity.^{20–26}

NAC has been available in clinical practice for several decades, and is predominantly used to treat acetaminophen intoxication. It can be administered intravenously, orally or by inhalation. Oral bioavailability is 6–10% because of first-pass metabolism.²⁷ Intravenous NAC carries a small risk of an anaphylaxis-like reaction, including rash, pruritus, angioedema, bronchospasm and, rarely, hypotension.²⁸ NAC given orally is associated with low toxicity, with reported non-life-threatening side effects including nausea, vomiting, rhinorrhoea, pruritus and tachycardia.

To date, studies investigating whether NAC can prevent aminoglycoside-induced ototoxicity have only evaluated the administration of NAC for short durations (10–14 days), and no studies have evaluated the impact of NAC on the polypeptides. However, if NAC were to be used in the context of MDR-TB to reduce aminoglycoside-induced ototoxicity, it would need to be administered for many months. NAC has been used for prolonged periods in patients with cystic fibrosis, COPD and psychiatric disorders.^{29–31} To date, however, studies have not specifically evaluated the safety profile and side effects associated with prolonged NAC use.

In order to assess the potential for NAC use in MDR-TB treatment, this review aimed to determine the effect of NAC on the development of ototoxicity when coadministered with aminoglycosides, as well as the safety and side effect profile of prolonged (>6 weeks) NAC administration.

METHODS

This review was conducted according to the criteria of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses group, and a protocol was developed before the review was conducted.³²

Randomised controlled trials and comparative observational studies were eligible for inclusion if they reported efficacy of concomitant NAC administration on ototoxicity prevention in patients receiving aminoglycoside treatment or if NAC was administered long term (≥6 weeks) regardless of indication. Case reports and case series (<20 patients) were excluded. No date, geographical or language restrictions were applied.

The primary outcome for the efficacy review was ototoxicity (proportion with any hearing loss, proportion with tinnitus and/or vertigo, and degree of hearing loss across different auditory frequencies). The primary outcome measures for the safety review included the number of adverse drug reactions, the number of individuals with an adverse event and/or side effect, and the number of individuals with each specific side effect associated with ≥6 weeks NAC administration. Secondary outcomes included the number of adverse drug reactions resulting in treatment discontinuation and mortality, and the total number of discontinuations and deaths.

Search strategy

A compound search strategy was developed (see online supplementary table S1) to identify all relevant studies regardless of language or publication status. The following electronic databases were searched: Medline (OVID), Embase (OVID), Web of Science, Current Controlled Trials, and the Cochrane database of systematic review. All references were imported into EndNote, and titles and abstracts were examined after duplicates were removed independently by two reviewers (WFE and KK). The full-text articles of all potentially relevant studies were obtained, and the inclusion criteria were applied using a standardised eligibility form. The full text of studies included in 26 previously published reviews investigating the effect of NAC on various chronic conditions was obtained and the studies were screened by two authors (WFE and KK) for inclusion.^{29–31–33–57} Reference lists of all studies identified by the above methods and bibliographies of systematic reviews or meta-analyses were examined. Final agreement on study inclusion was determined through consensus (WFE, KK).

Data extraction and management

Data extraction was performed independently, in duplicate, using a standardised data extraction form. Data regarding efficacy included information on the intervention (number of individuals in the intervention (receiving NAC) and control groups, and dose, frequency, duration and serum concentration of aminoglycoside and NAC), outcomes (number of individuals with evidence of ototoxicity per group) and patient characteristics (age, sex and presence of comorbidities). For the review investigating safety, the underlying condition for which NAC was administered, exclusion criteria, age, sex and comorbidities were recorded, as well as NAC dose, frequency and duration, number of adverse events, number of total deaths and withdrawals attributable to NAC.

Quality of included studies

For randomised trials investigating the efficacy of ototoxicity, the Cochrane risk of bias tool for quality assessment of randomised controlled trials was used, and this information was used to inform an overall assessment of quality using GRADE. Studies reporting safety data were assessed taking into account the design (retrospective/prospective), allocation of intervention (randomised, non-randomised), placebo use, blinding of participants and/or investigators, and monitoring strategy.

Data analysis

Relative risk (RR), risk difference and the frequency of events and corresponding 95% CIs for prevention of ototoxicity and side effects/adverse events were calculated. For relative effect measures, the Haldane method was applied in the event of zero outcomes in one arm; for frequencies, data were transformed before pooling using standard methods.^{58 59} Data were pooled using a fixed-effects model, and heterogeneity assessed using the I^2 statistic.⁶⁰ Pooled frequency estimates, risk ratios, risk difference and corresponding 95% CIs for specific side effect, total withdrawals and death and withdrawals attributable to NAC were calculated. Only placebo-controlled studies and studies in which solely NAC was administered were included to calculate the pooled estimates for side effects. Weighted medians were calculated for dosage and duration of NAC. All data analysis was performed using Stata V.12.0.

RESULTS

From 5941 unique citations identified, 86 studies were included in this review, among which only three studies reported on the efficacy of NAC to prevent ototoxicity in the context of aminoglycoside use, and 83 reported on long-term (≥ 6 weeks) NAC use for other purposes (see online supplementary figure S1). No studies assessing the use of NAC together with aminoglycoside for the treatment of MDR-TB were identified.

Prevention of ototoxicity

Three randomised trials reported on the efficacy of NAC to prevent ototoxicity including a total of 146 patients with end-stage renal disease receiving aminoglycosides for the treatment of bloodstream infections (table 1). Two of these trials were open-label,^{14 16} and one was a randomised, placebo-controlled trial.¹⁷ Patients received 600 mg NAC twice daily for the duration of aminoglycoside treatment,¹⁶ for a total of 14 days¹⁷ or for up to 7 days¹⁴ after completion of aminoglycoside treatment. The aminoglycosides used were amikacin (n=2) and gentamicin (n=1). Two of the studies measured the mean hearing loss (in dB) 1–2 and 4–6 weeks after enrolment and found a significant reduction in aminoglycoside-induced hearing loss at both time points.^{14 16} The pooled RR for otoprotection at 4–6 weeks was 0.14 (95% CI 0.05 to 0.45) and the pooled risk difference was –33.3% (95% CI –45.5% to –21.2%) (figure 1). One study compared transient-evoked otoacoustic emissions (OAEs) and distortion-product OAEs among 23 patients receiving placebo and 23 receiving NAC. This study reported a significant improvement in patients receiving NAC at 1500 and 2000 Hz when measured using transient-evoked OAEs and at 1000 and 800 Hz in terms of distortion-product OAEs.¹⁷ Overall the aminoglycosides caused the greatest hearing loss at high frequencies, and so it was at these frequencies that the most protective effect of NAC was seen.

The risk of bias was high because two of the three trials were open-label and did not include a control group with placebo (table 2). The overall quality of the evidence was rated as low/very low because of risk of bias and indirectness (different patient population).

Long-term NAC use

We identified a total of 83 studies describing the administration of NAC for >6 weeks. NAC was used for psychiatric (N=15), respiratory (N=26) and rheumatological conditions (N=6), blood-borne viruses (N=14), kidney disease (N=6), obstetric and gynaecological conditions (N=5), male infertility (N=2)

Table 1 Studies investigating the effect of NAC in preventing aminoglycoside-induced ototoxicity

Author, year	Condition	AG	N (NAC)	N (P)	NAC dose	Duration of NAC	Duration of AG	Cumulative dose, g	N hearing loss (NAC)	N hearing loss (P)	Risk ratio hearing loss (95% CI)	Mean hearing loss, dB (NAC), early	Mean hearing loss, dB (NAC), late	Mean hearing loss, dB (P), early	Mean hearing loss, dB (P), late
Feldman, 2007 ¹⁴	Haemodialysis, sepsis	Gentamicin	53*	20	20	600 mg twice daily	For the duration of gentamicin therapy until 1 week after completing therapy	14.8±3.8 (NAC) 14.3 (P) ±5.8 (P)	0.68 (NAC) 0.69 (P)	2†	11†	2.0±3.8†	2.1±5.2§	5.8±5.1†	7.0±8.2§
Tokgoz, 2011 ¹⁶	CAPD peritonitis	Amikacin	60	30	30	600 mg twice daily	For the duration of amikacin therapy	1.5 (NAC) 1.25 (P)	1¶	21¶	0.08 (0.01 to 0.55)	–4.7±7.4**	–6.0±8.9††	5.4±8.6**	16.9±8.1††
Kocigit, 2014 ¹⁷	CAPD peritonitis	Amikacin	50§§	23	23	600 mg twice daily	2 weeks	1.5 (NAC) 1.2 (P)	1.5 (NAC) 1.2 (P)	9 (4–20) (NAC) 8 (4–21) (P)					

*In the NAC group, 1 died, 3 had airborne discrepancies, 2 were unable to cooperate; in the placebo group, 2 died, 3 had airborne discrepancies, 1 was unable to cooperate, 1 withdrew consent.

†6 weeks after completing gentamicin therapy.

‡1 week after completing gentamicin therapy, at frequencies 6000, 8000, 12 000 Hz.

§6 weeks after completing gentamicin therapy, at frequencies 6000, 8000, 12 000 Hz.

¶14 weeks after starting amikacin therapy.

**1 week after starting amikacin therapy, at frequency 10 000, 12 000, 14 000, 16 000 Hz.


††4 weeks after starting amikacin therapy, at frequency 10 000, 12 000, 14 000, 16 000 Hz.

‡‡No measurement of pure-tone average hearing threshold, but measurement of transient-evoked otoacoustic emissions and distortion-product otoacoustic emissions.

§§In the NAC group, 1 withdrew consent, 1 dropped out; in the placebo group, 1 died, 1 withdrew consent.

AG, aminoglycoside; CAPD, continuous ambulatory peritoneal dialysis; NAC, N-acetylcysteine; P, placebo.

Figure 1 Effect of *N*-acetylcysteine on aminoglycoside-induced ototoxicity.



and other conditions (N=9) such as non-alcoholic fatty liver disease, diabetes mellitus and Alzheimer’s disease (see online supplementary table S2). The majority of studies were randomised, placebo-controlled trials (N=52); the remainder were randomised, placebo-controlled crossover trials (N=11), randomised crossover trials without placebo (N=2), randomised trials without administration of a placebo (N=17), and a quasi-experimental study with patients choosing to take NAC (N=1). Six studies described the administration of NAC or placebo in combination with metformin, cotrimoxazole, omeprazole, lipoic acid and interferon. A total of 5014 patients received a median of 1200 mg (IQR 600–1800) of NAC per day over a median of 24 weeks (IQR 12–54), and 4974 patients served as a control group. The majority of studies were conducted in Europe (N=44), and 13 studies were conducted in the USA. The age of participants ranged from 5 to 80 years, and severe liver and renal impairment was common. Two studies were conducted exclusively among pregnant woman, with a total of 220 pregnant woman receiving NAC. Specific side effects were reported in only 23 (28%) studies. A statement regarding adverse drug reactions was included in 32 (39%) studies in the results section (see online supplementary tables S3 and S4). The remaining 28 (34%) studies provided no information on adverse drug reactions.

More than half of the studies (n=49) reported the number of patients withdrawn from the study; of those, 34 reported the reason for withdrawal. No deaths were reported as being attributable to NAC administration. There was no difference in the risk of overall withdrawal with a pooled RR of 1.06 (95% CI 0.96 to 1.17, I² 0%) and a pooled risk difference of 0.9% (95% CI 0.6% to 2.5%, I² 0%) in the NAC compared with the control group. Furthermore, there was no increased risk of withdrawal attributable to NAC (pooled RR 0.74 (95% CI 0.59 to 0.93, I² 31%)) and the pooled risk difference was –1.6% (95% CI –2.8% to 0.0%, I² 67%) when comparing NAC with placebo or control group (figure 2). The pooled mortality was 1.1 (95% CI 0.91 to 1.31, I² 0%) across the seven studies reporting data on deaths.

Pooled estimates for specific side effects are presented in table 3. The most commonly reported side effects were abdominal pain, nausea, and vomiting and diarrhoea. The risk of abdominal pain (pooled RR 1.4 (95% CI 1.1 to 2.8)), nausea and vomiting (pooled RR 2.0 (95% CI 1.3 to 3.0)), diarrhoea (pooled RR 1.8 (95% CI 1.0 to 3.2)) and arthralgia (pooled RR 2.2 (95% CI 1.2 to 4.1)) were all significantly increased in patients receiving NAC

compared with placebo. However, the pooled risk differences for all these side effects were relatively small, ranging from 1.6% for diarrhoea to 6.1% nausea. The risks of headache, rash, dizziness, cramps and drowsiness were not significantly increased.

Quality assessment was challenging, as procedures for side effect ascertainment were not reported. Furthermore, the majority of studies failed to report death and discontinuation of treatment because of side effects (see online supplementary table S5).

DISCUSSION

This review identified three randomised trials reporting a protective effect of NAC in preventing aminoglycoside-induced ototoxicity in patients with end-stage renal failure. The short duration of aminoglycoside administration (maximum 3 weeks) and the selected patient population mean that limited inference regarding the applicability of these results to MDR-TB can be made. The overall quality of evidence informing this intervention was rated as low.

The safety of prolonged NAC administration was also addressed in this review, with 83 studies identified in which oral NAC was administered for a minimum of 6 weeks. Specific side effects were only reported in 23 of 83 studies included for review. Pooled RRs for specific adverse side effects showed a 1.4–2.2 times increased risk of abdominal pain, nausea and vomiting, diarrhoea and arthralgia in patients receiving NAC compared with placebo. The proportion of patients developing specific side effects was highly heterogeneous across studies, which is not surprising given the variety of clinical conditions, the wide age ranges, and difference in NAC dosing. The pooled risk difference was highest for nausea and vomiting (6.1%) and lowest for diarrhoea (1.8%). Thirty-two studies commented on side effects and adverse events without providing detailed information on specific adverse side effects. An additional 28 studies administered NAC, but neither reported nor commented on side effects. Most studies did not report data on discontinuation of treatment due to adverse events. However, withdrawal overall and withdrawal attributable to side effects was comparable in patients receiving NAC and placebo or control, where reported. Furthermore, mortality was comparable in the few studies that reported deaths stratified by treatment group.

This review provides evidence for the safety of prolonged NAC administration. However, the reported side effects associated with NAC use are potentially additive to those associated with second-line TB drugs other than the aminoglycosides.^{61 62}

Thorax: first published as 10.1136/thoraxjnl-2015-207245 on 7 September 2015. Downloaded from <http://thorax.bmj.com/> on April 10, 2024 by guest. Protected by copyright.

Table 2 Quality assessment of studies included to assess the effect of NAC on aminoglycoside-induced ototoxicity

Feldman, 2007 ¹⁴		
Methods	Randomised, open-label, controlled, parallel, one centre, duration 50 days, intention to treat analysis	
Participants	53 patients aged 18+ on haemodialysis treated with gentamicin for dialysis catheter-related bacteraemia, excluded if treated with aminoglycosides 3 months before the episode or mechanical occlusion of the external ear or a perforated tympanic membrane	
Interventions	NAC 600 mg twice daily	
Outcomes	Pure-tone audiometry measurements at frequencies 250, 500, 1000, 2000, 3000, 4000, 6000, 8000, 12 000 Hz at 7±3 and 42 ±3 days after completion of gentamicin therapy	
Bias	Author's judgement	Support for judgement
Random sequence generation	Unclear	
Allocation concealment	High risk	No placebo
Blinding of participants and personnel	High risk	No placebo
Incomplete outcome data	Low risk	
Selective reporting	Unclear	Primary outcomes not specifically reported, no protocol available
Other bias	Low risk	
Tokgoz, 2011 ¹⁶		
Methods	Randomised, open-label, controlled, parallel, one centre, duration 28 days, analysis (per protocol or intention to treat) not clarified	
Participants	60 patients on peritoneal dialysis treated with amikacin for their first episode of peritonitis, excluded if tympanic membrane perforated and admitted after office hours	
Interventions	NAC 600 mg twice daily	
Outcomes	Pure-tone audiometry measurements at frequencies 250, 500, 1000, 2000, 3000, 4000, 6000, 8000, 10 000, 12 000, 14 000 and 16 000 Hz at 8±2 days and 28±2 days	
Bias	Author's judgement	Support for judgement
Random sequence generation	Low risk	Patients chose an envelop
Allocation concealment	High risk	No placebo
Blinding of participants and personnel	High risk	No placebo
Incomplete outcome data	Low risk	
Selective reporting	Unclear	Primary outcomes not specifically reported, no protocol available
Other bias	Low risk	
Kocyigit, 2014 ¹⁷		
Methods	Randomised, placebo controlled, parallel, one centre, duration 28 days, analysis (per protocol or intention to treat) not clarified	
Participants	46 patients on peritoneal dialysis treated with amikacin for their first episode of peritonitis, excluded if tympanic membrane perforated and admitted after office hours	
Interventions	NAC 600 mg twice daily or placebo	
Outcomes	Transient-evoked otoacoustic emissions and distortion-product otoacoustic emissions at 1 and 4 weeks	
Bias	Author's judgement	Support for judgement
Random sequence generation	Low risk	Patients chose an envelop
Allocation concealment	Low risk	Placebo administered
Blinding of participants and personnel	Low risk	Measurements of patients were performed by staff who did not know which patient belonged to which group
Incomplete outcome data	Low risk	
Selective reporting	Unclear	Primary outcomes not specifically reported, no protocol available
Other bias	Low risk	

NAC, *N*-acetylcysteine.

Gastrointestinal side effects, such as nausea and vomiting, are associated with thioamides, para-aminosalicylic acid and fluoroquinolones, and, while not life-threatening, severely affect regimen tolerability and therefore potentially default from MDR-TB treatment. Failure to complete MDR-TB treatment is a significant contributor to poor treatment outcomes and generates further resistance.⁶³ In addition, the pill burden associated with MDR-TB treatment is considerable and the impact of adding further medication has to be considered carefully, weighing risks and benefits.

Patients receiving prolonged NAC had a variety of clinical conditions including respiratory, renal, liver, infectious (HIV and hepatitis C), obstetric and psychiatric diseases. The severity of diseases and the frequency of other comorbidities, including

drug and alcohol misuse, were heterogeneous across studies, with some studies including patients with life-threatening conditions such as idiopathic lung fibrosis, systemic sclerosis and end-stage liver disease. The age of included subjects spanned children less than 1 year of age to patients aged over 80 years. Three studies specifically included pregnant women, with a total of 220 receiving NAC. The safe administration of NAC across such a broad spectrum of patients is reassuring when considering its use as adjuvant treatment in patients with MDR-TB. HIV infection, alcohol and drug misuse, and smoking are common in patients with MDR-TB.^{64–66} Furthermore, a considerable proportion of patients treated for MDR-TB experience depression, or develop hepatic and renal impairment as a result of treatment.^{67–69} Thus, the safety of any adjunctive therapy needs to




Figure 2 Risk of withdrawals attributable to *N*-acetylcysteine.

be assessed in the context of both the comorbidities and the side effect profile of MDR-TB treatment.

This review was unable to assess potential drug interactions between NAC and drugs used for the treatment of TB. The trials investigating the ototoxic effects of NAC did not report any drug interactions between NAC and aminoglycosides. However, aminoglycosides and NAC were coadministered for a relatively short duration (2–4 weeks) compared with the 8 months of aminoglycoside that is currently recommended for treatment of MDR-TB. Data on interactions between NAC and other anti-TB drugs are lacking, with only one trial conducted in patients on first-line treatment in Iran⁷⁰; this trial randomised 60 patients being treated with first-line four-drug TB therapy (isoniazid, rifampicin, ethambutol and pyrazinamide) to either receiving 600 mg NAC twice daily for 2 weeks or no additional treatment. The authors hypothesised that NAC would reduce the frequency of drug-induced hepatitis because of its antioxidative properties. The trial reported a significant reduction of alanine aminotransferase and aspartate aminotransferase after 2 weeks of treatment, but drug interactions were not specifically reported.

The strengths of this review include a broad compound search strategy across five different databases and no restrictions with regards to date of publication, language and setting. The safety review included studies administering oral NAC for a minimum of 6 weeks regardless of the disease studied. This permitted the assessment of safety across a broad range of diseases and severity.

This review served to identify a number of important limitations to the existing evidence base. Notably, no studies were identified that investigated the ototoxic potential of NAC in patients receiving aminoglycosides for the treatment of MDR-TB. Furthermore, most studies identified in the safety review failed to provide sufficient information on specific side effects, resulting in a poor quality rating for the purpose of the safety review. The frequency of investigations carried out to assess side effects was only reported in a minority of studies. Thus, the ascertainment of side effects might have been of different quality within and across studies. The review was unable to establish the quality of ascertainment systematically because of lack of information.

Notwithstanding these limitations, the results of this review together with recent findings explaining the mechanism of

Table 3 Pooled estimates of the frequency of specific side effects with risk ratios and risk differences

Side effect	Number of studies	Frequency (NAC)		Frequency (placebo)		Pooled risk ratio		Pooled risk difference	
		Estimate (95% CI)	I ² , p value	Estimate (95% CI)	I ² , p value	Estimate (95% CI)	I ² , p value	Estimate (95% CI)	I ² , p value
Abdominal pain	14	11.8% (8.0% to 15.6%)	93.4%, <0.01	6.1% (3.5% to 8.7%)	88.6%, <0.01	1.4 (1.1 to 1.8)	41.8%, 0.05	2.6% (0.1% to 4.4%)	77.3%, <0.01
Nausea	8	17.6% (9.7% to 25.4%)	91.6%, <0.01	8.1% (3.6% to 12.5%)	61.6%, <0.01	2.0 (1.3 to 3.0)	54.9%, 0.03	6.1% (2.8% to 9.3%)	86.9%, <0.01
Diarrhoea	10	4.2% (1.6% to 6.8%)	66.5%, <0.01	1.8% (0.4% to 3.2%)	38.1%, 0.10	1.8 (1.0 to 3.2)	0%, 0.87	1.6% (0.1% to 3.0%)	27.7%, 0.19
Headache	6	14.4% (6.4% to 22.4%)	81.8%, <0.01	8.6% (5.2% to 12.0%)	0%, 0.78	1.4 (0.8 to 2.4)	0%, 0.47	3.2% (1.4% to 7.9%)	44.2%, 0.11
Arthralgia	7	6.8% (2.6% to 10.9%)	80.4%, <0.01	0.6% (0.4% to 1.5%)	54.3%, 0.04	2.2 (1.2 to 4.1)	0%, 0.67	1.8% (0.4% to 3.2%)	72.3%, <0.01
Rash	7	5.2% (2.1% to 8.2%)	25.8%, 0.23	2.0% (0.2% to 4.3%)	40.3%, 0.12	1.2 (0.6 to 2.4)	0%, 0.56	1.2% (-3.1% to 5.4%)	0%, 0.59
Dizziness	4	6.1% (-0.5% to 12.7%)	78.7%, <0.01	2.9% (-0.5% to 6.3%)	45.2%, 0.14	1.0 (0.5 to 1.8)	20.5%, 0.29	-0.1% (-0.2% to 1.7%)	27.7%, 0.25
Cramps	3	3.4% (-1.7% to 8.5%)	81.8%, <0.01	4.3% (1.8% to 10.5%)	76.5%, 0.01	1.2 (0.5 to 3.1)	39.9%, 0.19	0.7% (-2.4% to 3.8%)	54.3%, 0.11
Drowsiness	3	19.4% (9.5% to 37.8%)	81.6%, 0<0.01	12.5% (5.4% to 19.6%)	0%, 0.66	1.3 (0.62 to 2.7)	0%, 0.42	3.2% (-0.6% to 12.0%)	19.9%, 0.29

NAC, N-acetylcysteine.

aminoglycoside-induced ototoxicity and the established antioxidative properties of NAC provide a strong justification for a clinical trial to investigate the effect of concomitant NAC treatment in patients receiving aminoglycosides as part of MDR-TB treatment. The roll-out of the Xpert MTB/RIF (a test to detect TB and rifampicin resistance directly from sputum) in countries with limited laboratory capacity is likely to increase the number of MDR-TB diagnoses and consequently those receiving aminoglycoside therapy, many in areas with limited capacity to monitor ototoxicity.⁷¹ While the desired aim for MDR-TB treatment is to develop shortened, more tolerable and aminoglycoside-sparing regimens,⁷² this may be some way from reality. Published studies of novel combinations are promising.⁷³ Nevertheless, a promising 9-month regimen, piloted in Bangladesh which showed good results in the absence of fluoroquinolone resistance,⁷⁴ and which is now being evaluated in the STREAM trial, has aminoglycosides as a key part of the regimen.⁷⁵ While we await the incorporation of bedaquiline and delamanid, two newly registered second-line anti-TB drugs, into recommended second-line treatment regimens, it is likely that aminoglycosides will remain a key pillar of MDR-TB treatment, albeit potentially for shortened periods.⁷⁵

Acknowledgements Isatou N'jie and Timo Pilgram from the Knowledge and Library Services, Barts Health NHS trust for helping with retrieval of references.

Contributors WFE, FD and KK conceptualised the idea. KK developed the protocol and conducted the literature searches. KK and WFE screened the titles and abstracts and the full-text manuscripts for eligibility, and performed the data extraction. The meta-analysis was performed by KK and NF. KK wrote the paper with input from WFE, HC, JAS, NF and FD. All authors read and approved the final version of the manuscript.

Funding This study was funded by internal funds (National Mycobacterium Reference Laboratory, Public Health England).

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All authors had full access to all data and can take responsibility for the integrity of the data and the accuracy of the data analysis.

REFERENCES

- Segal JA, Skolnick P. Polyamine-like actions of aminoglycosides and aminoglycoside derivatives at NMDA receptors. *Eur J Pharmacol* 1998;347:311–17.
- Chang J, Yang JY, Choi J, et al. Calcium imaging in gentamicin ototoxicity: increased intracellular calcium relates to oxidative stress and late apoptosis. *Int J Pediatr Otorhinolaryngol* 2011;75:1616–22.
- Santaolalla F, Salvador C, Martinez A, et al. Inner ear hair cell regeneration: A look from the past to the future. *Neural Regen Res* 2013;8:2284–9.
- Selimoglu E. Aminoglycoside-induced ototoxicity. *Curr Pharm Des* 2007;13:119–26.
- Black FO, Pesznecker S, Stallings V. Permanent gentamicin vestibulotoxicity. *Otol Neurotol* 2004;25:559–69.
- WHO. *Guidelines for the programmatic management of drug-resistant tuberculosis 2011 update*. Geneva, Switzerland: World Health Organization, 2011.
- Seddon JA, Godfrey-Faussett P, Jacobs K, et al. Hearing loss in patients on treatment for drug-resistant tuberculosis. *Eur Respir J* 2012;40:1277–86.
- Peloquin CA, Berning SE, Nitta AT, et al. Aminoglycoside toxicity: daily versus thrice-weekly dosing for treatment of mycobacterial diseases. *Clin Infect Dis* 2004;38:1538–44.
- Sturdy A, Goodman A, Jose RJ, et al. Multidrug-resistant tuberculosis (MDR-TB) treatment in the UK: a study of injectable use and toxicity in practice. *J Antimicrob Chemother* 2011;66:1815–20.
- Ahuja SD, Ashkin D, Avendano M, et al. Multidrug resistant pulmonary tuberculosis treatment regimens and patient outcomes: an individual patient data meta-analysis of 9,153 patients. *PLoS Med* 2012;9:e1001300.
- Darrat I, Ahmad N, Seidman K, et al. Auditory research involving antioxidants. *Curr Opin Otolaryngol Head Neck Surg* 2007;15:358–63.
- Sinswat P, Wu WJ, Sha SH, et al. Protection from ototoxicity of intraperitoneal gentamicin in guinea pig. *Kidney Int* 2000;58:2525–32.
- Sha SH, Qiu JH, Schacht J. Aspirin to prevent gentamicin-induced hearing loss. *N Engl J Med* 2006;354:1856–7.
- Feldman L, Efrati S, Eviatar E, et al. Gentamicin-induced ototoxicity in hemodialysis patients is ameliorated by N-acetylcysteine. *Kidney Int* 2007;72:359–63.

- 15 Feldman L, Sherman RA, Weissgarten J. N-acetylcysteine use for amelioration of aminoglycoside-induced ototoxicity in dialysis patients. *Semin Dial* 2012;25:491–4.
- 16 Tokgoz B, Ucar C, Kocyigit I, et al. Protective effect of N-acetylcysteine from drug-induced ototoxicity in uraemic patients with CAPD peritonitis. *Nephrol Dial Transplant* 2011;26:4073–8.
- 17 Kocyigit I, Vural A, Unal A, et al. Preventing amikacin related ototoxicity with N-acetylcysteine in patients undergoing peritoneal dialysis. *Eur Arch Otorhinolaryngol* Published Online First: 30 Jul 2014. doi:10.1007/s00405-014-3207-z
- 18 Yarema MC, Johnson DW, Berlin RJ, et al. Comparison of the 20-hour intravenous and 72-hour oral acetylcysteine protocols for the treatment of acute acetaminophen poisoning. *Ann Emerg Med* 2009;54:606–14.
- 19 Weisbord SD, Gallagher M, Kaufman J, et al. Prevention of contrast-induced AKI: a review of published trials and the design of the prevention of serious adverse events following angiography (PRESERVE) trial. *Clin J Am Soc Nephrol* 2013;8:1618–31.
- 20 Bielefeld EC, Kopke RD, Jackson RL, et al. Noise protection with N-acetyl-L-cysteine (NAC) using a variety of noise exposures, NAC doses, and routes of administration. *Acta Oto-Laryngologica* 2007;127:914–19.
- 21 Coleman J, Huang XY, Liu JZ, et al. Random study on the effectiveness of salicylate/ N-acetylcysteine for prevention of noise-induced hearing loss. *Noise Health* 2010;12:159–65.
- 22 Ewert DL, Lu JZ, Li W, et al. Antioxidant treatment reduces blast-induced cochlear damage and hearing loss. *Hear Res* 2012;285:29–39.
- 23 Lorito G, Giordano P, Petruccioli J, et al. Different strategies in treating noise-induced hearing loss with N-acetylcysteine. *Med Sci Monit* 2008;14:BR159–64.
- 24 Lin CY, Wu JL, Shih TS, et al. N-Acetyl-cysteine against noise-induced temporary threshold shift in male workers. *Hear Res* 2010;269:42–7.
- 25 Yoo J, Hamilton SJ, Angel D, et al. Cisplatin ototoxicity using transtympanic L-N-acetylcysteine: a pilot randomized study in head and neck cancer patients. *Laryngoscope* 2014;124:E87–94.
- 26 Riga MG, Chelis L, Kakolyris S, et al. Transtympanic injections of N-acetylcysteine for the prevention of cisplatin-induced ototoxicity: a feasible method with promising efficacy. *Am J Clin Oncol* 2013;36:1–6.
- 27 Samuni Y, Goldstein S, Dean OM, et al. The chemistry and biological activities of N-acetylcysteine. *Biochim Biophys Acta* 2013;1830:4117–29.
- 28 Sandilands EA, Bateman DN. Adverse reactions associated with acetylcysteine. *Clin Toxicol (Phila)* 2009;47:81–8.
- 29 Cheng K, Smyth RL, Motley J, et al. Randomized controlled trials in cystic fibrosis (1966–1997) categorized by time, design, and intervention. *Pediatr Pulmonol* 2000;29:1–7.
- 30 Poole P, Black PN. Mucolytic agents for chronic bronchitis or chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2010;(2):CD001287.
- 31 Asevedo E, Mendes AC, Berk M, et al. Systematic review of N-acetylcysteine in the treatment of addictions. *Revista Brasileira De Psiquiatria* 2014;36:168–75.
- 32 Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 33 van Dalen EC, Caron HN, Dickinson HO, et al. Cardioprotective interventions for cancer patients receiving anthracyclines. *Cochrane Database Syst Rev* 2011;(6):CD003917.
- 34 Rothbart R, Amos T, Siegfried N, et al. Pharmacotherapy for trichotillomania. *Cochrane Database Syst Rev* 2013;11:CD007662.
- 35 Albers JW, Chaudhry V, Cavaletti G, et al. Interventions for preventing neuropathy caused by cisplatin and related compounds. *Cochrane Database Syst Rev* 2014;3: CD005228.
- 36 Showell MG, Brown J, Clarke J, et al. Antioxidants for female subfertility. *Cochrane Database Syst Rev* 2013;8:CD007807.
- 37 Poole P, Black PN, Cates CJ. Mucolytic agents for chronic bronchitis or chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012;8:CD001287.
- 38 Atkins CP, Loke YK, Wilson AM. Outcomes in idiopathic pulmonary fibrosis: a meta-analysis from placebo controlled trials. *Respir Med* 2014;108:376–87.
- 39 Sutherland ER, Crapo JD, Bowler RP. N-acetylcysteine and exacerbations of chronic obstructive pulmonary disease. [Review] [38 refs]. *COPD* 2006;3:195–202.
- 40 Boogaard R, de Jongste JC, Merkus PJFM. Pharmacotherapy of impaired mucociliary clearance in non-CF pediatric lung disease. A review of the literature. [Review] [93 refs]. *Pediatr Pulmonol* 2007;42:989–1001.
- 41 Block KI, Koch AC, Mead MN, et al. Impact of antioxidant supplementation on chemotherapeutic efficacy: a systematic review of the evidence from randomized controlled trials. *Cancer Treat Rev* 2007;33:407–18.
- 42 Brown KK, Wells AU. Recent clinical trials in idiopathic pulmonary fibrosis and the BUILD-1 study. *Eur Respir J* 2008;17:116–22.
- 43 Yancy WS Jr, McCrory DC, Coeytaux RR, et al. Efficacy and tolerability of treatments for chronic cough: a systematic review and meta-analysis. *Chest* 2013;144:1827–38.
- 44 Coombes JS, Fassett RG. Antioxidant therapy in hemodialysis patients: a systematic review. *Kidney Int* 2012;81:233–46.
- 45 Berk M, Malhi GS, Gray LJ, et al. The promise of N-acetylcysteine in neuropsychiatry. *Trends Pharmacol Sci* 2013;34:167–77.
- 46 Grandjean EM, Berthet P, Ruffmann R, et al. Efficacy of oral long-term N-acetylcysteine in chronic bronchopulmonary disease: a meta-analysis of published double-blind, placebo-controlled clinical trials. *Clinical Ther* 2000;22:209–21.
- 47 Duijvestijn YCM, Brand PLP. Systematic review of N-acetylcysteine in cystic fibrosis. *Acta Paediatrica* 1999;88:38–41.
- 48 Davies L, Calverley PMA. The evidence for the use of oral mucolytic agents in chronic obstructive pulmonary disease (COPD). *Br Med Bull* 2010;93:217–27.
- 49 Mecocci P, Polidori MC. Antioxidant clinical trials in mild cognitive impairment and Alzheimer's disease. *Biochim Biophys Acta* 2012;5:631–8.
- 50 Musso G, Gambino R, Cassader M, et al. A meta-analysis of randomized trials for the treatment of nonalcoholic fatty liver disease. *Hepatology* 2010;52:79–104.
- 51 Qureshi NA, Al-Bedah AM. Mood disorders and complementary and alternative medicine: a literature review. *Neuropsychiatr Dis Treat* 2013;9:639–58.
- 52 Poole C. Low P-values or narrow confidence intervals: which are more durable?. *Epidemiology* 2001;12:291–4.
- 53 Dean OM, Bush AI, Copolov DL, et al. Effects of N-acetyl cysteine on cognitive function in bipolar disorder. *Psychiatry Clin Neurosci* 2012;66:514–17.
- 54 Schloss JM, Colosimo M, Airey C, et al. Nutraceuticals and chemotherapy induced peripheral neuropathy (CIPN): A systematic review. *Clin Nutr* 2013;32:888–93.
- 55 Stey C, Steurer J, Bachmann S, et al. The effect of oral N-acetylcysteine in chronic bronchitis: a quantitative systematic review. *Eur Respir J* 2000;16:253–62.
- 56 Sommer IE, van Westrhenen R, Begemann MJH, et al. Efficacy of anti-inflammatory agents to improve symptoms in patients with schizophrenia: an update. *Schizophr Bull* 2014;40:181–91.
- 57 Shen Y, Cai W, Lei S, et al. Effect of high/low dose N-acetylcysteine on chronic obstructive pulmonary disease: a systematic review and meta-analysis. *COPD* 2014;11:351–8.
- 58 Freeman MF, Tukey JW. Transformations related to the angular and the square root. *Ann Inst Stat Math* 1950;21:607–11.
- 59 Miller J. The inverse of the Freeman-Tukey double arcsine transformation. *Am Stat* 1978;32:138.
- 60 Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- 61 Isaakidis P, Varghese B, Mansoor H, et al. Adverse events among HIV/MDR-TB co-infected patients receiving antiretroviral and second line anti-TB treatment in Mumbai, India. *PLoS ONE* 2012;7:e40781.
- 62 Nathanson E, Gupta R, Huamani P, et al. Adverse events in the treatment of multidrug-resistant tuberculosis: results from the DOTS-Plus initiative. *Int J Tuberc Lung Dis* 2004;8:1382–4.
- 63 Tozcek A, Cox H, du Cros P, et al. Strategies for reducing treatment default in drug-resistant tuberculosis: systematic review and meta-analysis. *Int J Tuberc Lung Dis* 2013;17:299–307.
- 64 Flor de Lima B, Tavares M. Risk factors for extensively drug-resistant tuberculosis: a review. *Clin Respir J* 2014;8:11–23.
- 65 Cavanaugh JS, Kazenny BY, Nguyen ML, et al. Outcomes and follow-up of patients treated for multidrug-resistant tuberculosis in Orel, Russia, 2002–2005. *Int J Tuberc Lung Dis* 2012;16:1069–74.
- 66 Isaakidis P, Das M, Kumar AM, et al. Alarming levels of drug-resistant tuberculosis in HIV-infected patients in metropolitan Mumbai, India. *PLoS ONE* 2014;9:e110461.
- 67 Das M, Isaakidis P, Van den Bergh R, et al. HIV, multidrug-resistant TB and depressive symptoms: when three conditions collide. *Glob Health Action* 2014;7:24912.
- 68 Baghaei P, Tabarsi P, Dorriz D, et al. Adverse effects of multidrug-resistant tuberculosis treatment with a standardized regimen: a report from Iran. *Am J Ther* 2011;18:e29–34.
- 69 Wu S, Zhang Y, Sun F, et al. Adverse events associated with the treatment of multidrug-resistant tuberculosis: a systematic review and meta-analysis. *Am J Ther* Published Online First: 26 Nov 2013. doi:10.1097/01.mjt.0000433951.09030.5a
- 70 Baniyadi S, Filik L. Protective effect of N-acetyl cysteine on antituberculosis drug-induced hepatotoxicity. *Eur J Gastroenterol Hepatol* 2011;23:193; author reply.
- 71 Creswell J, Codlin AJ, Andre E, et al. Results from early programmatic implementation of Xpert MTB/RIF testing in nine countries. *BMC Infect Dis* 2014;14:2.
- 72 Brigid G, Nyang'wa BT, du Cros P, et al. Principles for designing future regimens for multidrug-resistant tuberculosis. *Bull World Health Organ* 2014;92:68–74.
- 73 Diacon AH, Dawson R, von Groote-Bidlingmaier F, et al. 14-day bactericidal activity of PA-824, bedaquiline, pyrazinamide, and moxifloxacin combinations: a randomised trial. *Lancet* 2012;380:986–93.
- 74 Aung KJ, Van Deun A, Declercq E, et al. Successful '9-month Bangladesh regimen' for multidrug-resistant tuberculosis among over 500 consecutive patients. *Int J Tuberc Lung Dis* 2014;18:1180–7.
- 75 Nunn AJ, Rusen ID, Van Deun A, et al. Evaluation of a standardized treatment regimen of anti-tuberculosis drugs for patients with multi-drug-resistant tuberculosis (STREAM): study protocol for a randomized controlled trial. *Trials* 2014;15:353.

Supplement, table 1: Search strategy

Set	Topic	Medline	Embase
Set 1	N-acetylcysteine	ACETYLCYSTEINE	ACETYLCYESTINE or ACEYTLCYSTEINE DERIVATE
Set 2		n-acetylcysteine or acetylsteine	Aceytlcysteine
Set 3	Ototoxicity	HEARING LOSS	HEARING IMPAIRMEN
Set 4		COCHLEA	OTOTOXICITY
Set 5		ototoxicity or cochleotoxic or ototoxic	ototoxicity
Set 6		(hearing adj1 loss)	(hearing adj1 loss)
Set 7		(hearing ajd1 impairment)	(hearing ajd1 impairment)
Set 8	Long term administration	RADOMIZED CONTROLLED TRIAL	RADOMIZED CONTROLLED TRIAL
Set 9		CLINICAL TRIAL	CLINICAL TRIAL
Set 10		OBSERVATIONAL STUDY	OBSERVATIONAL STUDY
Set 11		LONGITUDINAL STUDY	LONGITUDINAL STUDY
Set 12		COHORT STUDY	COHORT ANALYSIS
Set 13		(observational adj1 study)	(observational adj1 study)
Set 14		(longitudinal adj1 study)	(longitudinal adj1 study)
Set 15		(clinical adj1 trial)	(clinical adj1 trial)
Set 16		combine set1 and set 2 with “or”	
Set 17		combine set3-7 with “or”	
Set 18		combine set8-15 with “or”	
Set 19		combine set16-18 with “and”	
Set 20		Limit set19 to human	

Words in capital letters were searched as MESH terms and as free text terms.

Web of Science

Combination of “acetylcysteine” “ototoxic” “vertigo” “longitudinal” “randomized controlled” “clinics trial”

Supplement Table 2: Studies reporting on long-term NAC use

Author, year	Country	Disease or Condition	NAC dose (mg)	Frequency of dosing	Duration (weeks)	N (NAC)	N (placebo)	Age - NAC	N men - NAC	Age - Placebo	N men - Placebo
Psychiatric disease											
Berk, 2012[1]	Australia	Bipolar disorder	1000	2	12	76	73	47.1	16	44.4	32
Berk,2008[2]	Australia	Bi-polar disorder	1000	2	24	38	37	44.6	15	46.6	15
Berk, 2014[3]	Australia	Major depressive disorder	1000	2	12	135	134	49.9	46	50.5	54
Farokhnia, 2013[4]	Iran	Schizophrenia	1000	2	8	23	23	32.2	11	33.4	13
Carmeli, 2012 ^a [5]	Switzerland	Schizophrenia	2000	1	8	11	11	31.9	9	31.9	9
Lavoie, 2008 ^a [6]	Switzerland	Schizophrenia	1000	2	8	7	7	31.9	5	31.9	5
Berk, 2008[7]	Australia	Schizophrenia	1000	2	24	69	71	37.2	48	36.1	50
Afshar, 2012[8]	Iran	Obsessive compulsive disorder	600, 1200, 2400		12	24	24	30.6	6	31.3	6
Ghanizadeh, 2013[9]	Iran	Autistic disorder	600	2	8	20	20	8.8	13	7.9	12
Bloch,2013[10]	USA	Trichotillomania	1200	2	12	20	19	14	3	13.1	2
Grant, 2009 ^b [11]	USA	Trichotillomania	1200 (6wks), 2400 (6wks)	1	12	25	25	32.7	4	35.8	
Grant, 2007[12]	USA	Pathological gambling	600	2-3	6	6	7				
LaRow, 2013[13]	USA	Cocaine Dependence	600-1200	2	8	78	38	43.5	58	43.3	25
Gray, 2012[14]	USA	Cannabis dependency	1200	2	8	58	58	18.9	39	18.8	45
Grant,2010[15]	USA	Methamphetamine dependence	600, 1200, 1800, 2400		8	14	17	37.2	8	36.1	14
Respiratory disease											
Zheng, 2014[16]	China	COPD	600	2	52	504	502	66.2	415	66.4	409
Tse, 2013[17]	Hong Kong	COPD	600	2	52	58	62	71	54	70.8	58
de Backer, 2013 ^a [18]	Belgium	COPD	600	3	12	12	12	65	9	65	9
Patil, 2011[19]	India	COPD	600	1,2	8	54	23				
Stav, 2009 ^a [20]	Israel	COPD	1200	1	6	24	24	66		66	
Schermer, 2009[21]	Netherlands	COPD	600	1	24	96	96	59.2	75	59.6	65
Decramer, 2005[22]	Europe	COPD	600	1	156	256	267	62	204	62	210
de Benedetto, 2005[23]	Italy	COPD	600	2	8	32	23	66.2		66.3	
van Overveld, 2005 ^{a,b} [24]	Poland	COPD	600	1	10	20	20		16		16
Kasielski, 2001[25]	Poland	COPD	600	1	52	22	22	61	10	60	11
Pela, 1999 ^b [26]	Italy	COPD	600	1	24	85	84	66	60	66	68
Lukas, 2005[27]	Germany	Chronic bronchitis	600	2	12	15	17	53.6	9	58.0	8

Supplement Table 2: Studies reporting on long-term NAC use – con't

Author, year	Country	Disease or Condition	NAC dose (mg)	Frequency of dosing	Duration (weeks)	N (NAC)	N (placebo)	Age - NAC	N men - NAC	Age - Placebo	N men - Placebo
Hansen, 1994[28]	Denmark	Chronic bronchitis	600	2	22	75	78	51.1	30	51.7	36
Rasmussen, 1988[29]	Sweden	Chronic bronchitis	300	2	24	59	57	58.8	31	58.9	35
Poder, 1984 ^b [30]	Hungary	Chronic bronchitis	10mg/kg	2-3	12	27	18	0.5-3		0.5-3	
McGavin, 1985[31]	UK	Chronic bronchitis	200	3	20	85	96	64.3	75	62	80
Stafanger, 1989 ^a [32]	Denmark	Cystic fibrosis with <i>P. aeruginosa</i> infection	200 (<30kg), 400 (>30 kg)	3	12	52	52	15.8	17	15.8	17
Stafanger, 1988 ^a [33]	Denmark	Cystic fibrosis	200/400	3/2	12	22	22	9.5	23	9.5	23
Ratjen, 1985[34]	Germany	Cystic fibrosis	200	3	10	12	12	13.9	10	13.9	10
Mitchell, 1982 ^a [35]	New Zealand	Cystic fibrosis	200	3	12	20	20	10.8	10	10.8	10
Stafanger, 1988 ^a [33]	Denmark	Primary ciliary dyskinesia	200/400	3/2	12	8	8	29.7	6	29.7	6
Martinez, 2014[36]	USA	Idiopathic pulmonary fibrosis	600	3	60	133	131	68.3	107	67.2	98
Demedts, 2005[37]	Europe	Idiopathic pulmonary fibrosis	600	3	52	92	90	62	69	64	75
Ghanei, 2008[38]	Iran	Chronic lung disease due to mustard gas exposure	600	2	16	72	72	44.9	32	46.7	37
Van Zandwijk, 2000 ^b [39]	Multicentre Europe	Non-small-cell lung cancer, pT1-2, N0-1, T2N0 or cancer of the larynx	600	2	104	642	641	61	556	60	559
Van Schooten, 2002[40]	Netherlands	Smoker	600	2	24	21	20	42	6	44	8
Blood born viruses											
Milazzo, 2010 ^c [41]	Italy	HIV, lipoatrophy	2000	1	48	20	20	45	7	44	9
Spada, 2002[42]	Brazil	HIV	600	1	24	10	10				
Breitkreutz, 2000[43]	Germany	HIV, not on ART	600	6, 4, 2, 1	28	16	13		8		8
Breitkreutz, 2000[43]	Germany	HIV, ART	600	6, 4, 2, 1	28	21	16		11		10
de Rosa, 2000[44]	USA	HIV	8000	1	8	41	42	38	41	38	42
Walmsley, 1998 ^d [45]	Canada	HIV, PCP	3000	2	8	96	102	38.1	83	38.6	100
Look, 1998 ^{a,b} [46]	Germany	HIV	1200	1	12	24	24	36.5	17	36.5	17
Grant, 2000[47]	Spain/Italy	HCV	600	3	24	73	74	39.1	51	40.9	53
Neri, 2000 ^b [48]	Italy	HCV	1200	2	64	38	39				
Look, 1999 ^{b,e} [49]	Germany	HCV	1800	1	24	8	8	35.7	6	38.7	3
Ideo, 1999 ^{b,e,f} [50]	Italy	HCV	1200	1	24	58	62	48.3	33	46.9	37
Cimino, 1998 ^{b,f} [51]	Italy	HCV	1200	1	12	12	13	30-62	8	23-57	8
Tripi, 1998 ^b [52]	Italy	HCV	600	2	24	14	7	53.8	6	46.5	5
Bernhard, 1998[53]	Switzerland	HCV	600	3	24	19	17				

Supplement Table 2: Studies reporting on long-term NAC use – con't

Author, year	Country	Disease or Condition	NAC dose (mg)	Frequency of dosing	Duration (weeks)	N (NAC)	N (placebo)	Age - NAC	N men - NAC	Age - Placebo	N men - Placebo
Kidney disease											
Purwanto, 2012[54]	Indonesia	Peritoneal dialysis	600	2	8	16	16	45.8	10	42.5	11
Hashemi, 2012 ^b [55]	Iran	Proteinuria, diabetes mellitus type 2	600	2	8	35	35	60.2	19	63.4	20
Renke, 2010/2008 ^a [56-57]	Poland	Non diabetic chronic kidney disease	600	2	8	20	20	39.4	12	39.4	12
Nascimento, 2009[58]	Brazil	Peritoneal dialysis	600	2	8	12	10	57	5	54	4
Hsu, 2009 ^b [59]	Taiwan	Hemodialysis	200	3	12	38	227	57.9	20	61.3	107
Tepel, 2003[60]	Germany	Hemodialysis	600	2	58	64	70	63	33	62	43
Obstetric and gynaecology conditions											
Popora, 2013 ^b [61]	Italy	Endometriosis	600	3	12	45	47	32.9	0	32.5	0
Hashim, 2009 ⁱ [62]	Egypt	Polycystic Ovary Syndrome	600	3	6	95	97	27.3	0	26.8	0
Elnashar, 2007 ^j [63]	Egypt	Polycystic Ovary Syndrome	600	3	6	30	31	26.7	0	27.3	0
Shahin, 2009 [64]	Egypt	Preterm labour	600	1	>12	140	140	26.5	0	25.9	0
Amin, 2008 ^b [65]	Egypt	Recurrent pregnancy Loss	600	1	20	80	86	26.2	0	25.2	0
Male infertility											
Safarinejad, 2009[66]	Iran	Male infertility	600	1	26	118	118	32	118	31	118
Ciftci, 2009[67]	Turkey	Male infertility	600	1	12	118	60	33.1	60	32.8	60
Rheumatological conditions											
Lai, 2012[68]	USA	Lupus erythematosus	600, 1200, 2400	3	12	18	9				
van Dielen, 2003[69]	Netherlands	Reflex sympathetic dystrophy	600	3	54	67	64	48	34	52	39
Perez, 2003 ^k [70]	Netherlands	Regional pain syndrome	600	3	17	74	71	50	29	49	20
Yalcin, 2002 ^b [71]	Turkey	Blepharitis	100	3	8	43	36	42.9	8	43.7	4
Currie, 1988 ^{a,l} [72]	UK	Young syndrome	200	3	8	8	8	37	8	37	8
Furst, 1978[73]	USA	Systemic sclerosis	10000		8	11	11	55.1	1	60.1	2
Others											
Kasperczyk, 2014 ^b [74]	Poland	Lead exposure	200	1, 2, 3	12	120	49				
Dabirmoghaddam, 2013 ^m [75]	Iran	Laryngopharyngeal Reflux	600	1	12	30	30				
Khoshbaten, 2010 ⁿ [76]	Iran	Non-alcoholic fatty liver	600	2	12	15	15	40.1	6	46.8	5
Martina, 2008[77]	Italy	Diabetes mellitus II	600	2	24	12	12	62.5	12	67	12
Pace, 2003[78]	USA	Sickle cell disease	200, 400, 800	3	28	16	5		8		3
Adair, 2001[79]	USA	Alzheimer disease	50 mg/kg/day	3	24	25	22				

Supplement Table 2: Studies reporting on long-term NAC use – con't

Author, year	Country	Disease or Condition	NAC dose (mg)	Frequency of dosing	Duration (weeks)	N (NAC)	N (placebo)	Age - NAC	N men - NAC	Age - Placebo	N men - Placebo
Estensen, 1999[80]	USA	Adenomatous colonic polyps	400	2	12	34	30				
de Flora, 1997[81]	Italy	Chronic degenerative disease (other than chronic respiratory disease)	600	2	24	133	129	69	60	68	48
Ardissino, 1997[82]	Italy	Unstable angina	600	3	16	45	46	58	32	58	32

^a Cross-over trial

^b No placebo in the control group

^c Patients were given NAC plus lipoic acid

^d All patients were treated for PCP with co-trimoxazole

^e All patients received IFN, patients in the NAC also received sodium selenite

^f Patients who did not respond to IFN treatment at 3 months were given NAC

^g No RCT, patients were given an option to take NAC or not

^h Drug administration 3 times per week

ⁱ All patients received metformin, no placebo in the control group

^j Patients in the comparison group received metformin (not placebo)

^k Comparison group DMSO cream

^l Patients in the control group took ambroxol or bromhexine or carbocisteine

^m All patients received omeprazole

ⁿ Patients in the control group took Vitamin C

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Total Withdrawn (NAC)	Total Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Berk, 2012[1]	76	73	Systemic medical disorder, respiratory insufficiency, pregnancy, breastfeeding	49% smokers, 47% alcohol use, 21% alcohol abuse	No side effects reported	21	18	1	0		Not reported	Not reported
Berk, 2008[2]	38	37	Respiratory disease, PUD, pregnancy, breastfeeding	79% alcohol use, 45% smokers	Side effects in text	14	13	Not reported	Not reported		Not reported	Not reported
Berk, 2014[3]	135	134	PUD, pregnancy, breastfeeding	22% CVS, 19% GI, 19% smokers, 60% alcohol use	Side effects in text	38	27	2	1		Not reported	Not reported
Farokhnia, 2013[4]	23	23	Serious medical or neurological disorders, alcohol or substance abuse, pregnancy, lactation, hepatic and kidney disease		Side effects reported	2	2	0	0		Not reported	Not reported
Carmeli, 2012 ^a [5]	11	11			No side effects reported	2	0	0	0		Not reported	Not reported
Lavoie, 2008 ^a [6]	7	7			No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Berk, 2008[7]	69	71	Abnormal renal, hepatic, thyroid or haematological findings, pregnancy	16% substance abuse, 53% alcohol use	Side effects in text	15	14	Not reported	Not reported	"There were no significant effects of NAC on safety parameters or reported adverse events"	Not reported	Not reported
Afshar, 2012[8]	24	24	Substance abuse, pregnancy, breastfeeding, convulsive disorder, suicidal		Side effects in text	5	4	3	0		Not reported	Not reported
Ghanizadeh, 2013[9]	20	20	Psychotic disorder, substance abuse, liver disease, seizures, hypertension, cardiac disease, unstable asthma, kidney disease		Side effects reported	3	6	1	0		Not reported	Not reported

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Total Withdrawn (NAC)	Total Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Bloch, 2013[10]	20	19	Bipolar, psychotic, substance abuse, development disorder, mental retardation, pregnancy		Side effects in text	4	0	1	0		Not reported	Not reported
Grant, 2009[11]	25	25	Unstable medical disease, abnormal laboratory tests, pregnancy, breastfeeding	60% psychiatric co-morbidity	Side effects reported	3	3	Not reported	Not reported		Not reported	Not reported
Grant, 2007[12]	6	7	Abnormal physical examination		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
LaRow, 2013[13]	78	38	Substance abuse, pregnancy, breastfeeding, serious medical condition, asthma	36% alcohol abuse,	Side effects reported	38	17	3	0		Not reported	Not reported
Gray, 2012[14]	58	58	Co-morbid substance abuse, unstable psychiatric disease, pregnancy	14% psychiatric co-morbidity	Side effects in text	21	25	Not reported	Not reported	"There were no FDA-defined serious adverse events and there were no significant differences between the two treatment groups in the occurrence of any adverse events (38 adverse events in 24 participants receiving NAC, 46 adverse events in 27 participants receiving placebo)"	Not reported	Not reported
Grant, 2010[15]	14	17	Medical disease, pregnancy, suicidal, bipolar disorder, dementia, psychotic disorder, abnormal liver function tests, substance abuse		Side effects in text	5	9	Not reported	Not reported	"Rates of side effects did not significantly differ between groups (57.1% NAC vs 41.5% placebo)"	Not reported	Not reported

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Withdrawn (NAC)	Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Zheng,, 2014[16]	504	502	Asthma, long term oxygen	18% smokers	Side effects reported	124	119	32	24		Not reported	Not reported
Tse, 2013[17]	58	62	Co-existing pulmonary disease, severe dyspnoea		Side effects in text	4	5	0	0	"No major adverse effects occurred in either group. There was no increase in incidence of minor adverse effects with NAC (3/58, 5%) vs placebo (5/62, 8%)"	2	1
de Backer, 2013[18]	12	12	Exacerbation during the last 8 weeks, PUD, steroids, pregnancy, breastfeeding		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Patil, 2011[19]	54	23	Decompensated cardiovascular, endocrine, hepatic or renal function, parenchymal lung pathology, active infection due to M tuberculosis, gastric or duodenal ulcer		Side effects in text	Not reported	Not reported	Not reported	Not reported	"Nausea and stomatitis were the most common adverse reactions"	Not reported	Not reported
Stav, 2009[20]	24	24	Asthma, long term oxygen		Side effects in text	Not reported	Not reported	Not reported	Not reported	"Apart from mild epigastric discomfort that was reported by a few patients in the treated group, no other complaints or findings were recorded"	Not reported	Not reported
Schermer, 2009[21]	96	96	Asthma, allergic rhinitis, eczema		No side effects reported	44	40	4	4		1	3

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Withdrawn (NAC)	Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Decramer, 2005[22]	256	267	Long term, PUD, congestive heart failure, oral steroids, cystic fibrosis, bronchiectasis, past history of TB		Side effects in text	70	99	19	26	"1428 adverse events in NAC group, 1381 adverse events in placebo group, no adverse events were thought to be drug-related"	Not reported	Not reported
de Benedetto, 2005[23]	32	23	Neoplasma, DIP, pulmonary disease		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
van Overveld, 2005[24]	20	20	Atopy, asthma, hepatic or renal failure, cystic fibrosis, CVS		No side effects reported	not reported	not reported	Not reported	Not reported		Not reported	Not reported
Kasielski, 2001[25]	22	22	Alcohol or substance abuse		Side effects in text	not reported	not reported	Not reported	Not reported	"The number of adverse events was low: two in the NAC group and three in the placebo group"	Not reported	Not reported
Pela, 1999[26]	85	84	Lung cancer, cardiomyopathy, metabolic disease, chronic renal disease	28% smokers	Side effects in text	2	3	1	0	"NAC was well tolerated. There was no difference in side effects reported in both groups"	0	1
Lukas, 2005[27]	17	15			No side effects reported	8	10	Not reported	Not reported		Not reported	Not reported
Hansen, 1994[28]	75	78	Eosinophilia, positive skin test to allergens, long term antibiotic treatment		Side effects in text	16	8	Not reported	Not reported	"There were no serious adverse events during the study"	Not reported	Not reported
Rasmussen, 1988[29]	59	57	Pregnancy, antibiotics		Side effects reported	10	7	10	7		Not reported	Not reported
Poder, 1984[30]	27	18			Side effects in text	Not reported	Not reported	Not reported	Not reported	"No side effects were observed"	Not reported	Not reported
McGavin, 1985[31]	85	96	Bronchiectasis, insulin dependent DM, PUD, pregnancy	27% smokers	Side effects reported	13	20	1	2		3	2

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Withdrawn (NAC)	Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Stafanger, 1989 ¹ [32]	52	52	PUD, liver of kidney disease, pregnancy		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Stafanger, 1988[33]	22	22	PUD, liver of kidney disease, pregnancy		Side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Ratjen, 1985[34]	12	12	Atopy, bronchodilators		Side effects in text	3	1	Not reported	Not reported	"No side effects of active therapy were noted"	Not reported	Not reported
Mitchell, 1982[35]	20	20			Side effects in text	2	2	0	0	"No side effects were noticed for either placebo or NAC"	Not reported	Not reported
Stafanger, 1988[33]	8	8	PUD, liver of kidney disease, pregnancy		Side effects reported	9	12	1	1		Not reported	Not reported
Martinez, 2014[36]	133	131	Non-idiopathic fibrotic lung disease, coexisting medical disease, on the waiting list of a lung transplant	23% CVS, 19% diabetes, 61% GERD	Side effects reported	23	20	1	4		Not reported	Not reported
Demedts, 2005[37]	92	90	Prednisolone dose >0.5mg/kg		Side effects reported	16	16	2	2		7	8
Ghanei, 2008[38]	72	72	Pneumonia, history of TB, smoking, substance abuse		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Van Zandwijk, 2000[39]	642	641	Recurrent disease, synchronous multiple tumours, previous malignant disease, abnormal LFTs, abnormal renal function, DM, HTN, PUD	93% smokers	Side effects reported	115	Not applicable	Not reported	Not reported	"No comparison of side effects to the non-treatment groups (the control group did not receive any placebo), GI and skin side effects were similar in the groups receiving NAC, NAC+Retinyl and Retinyl alone."	167	147
Van Schooten, 2002[40]	21	20			No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Withdrawn (NAC)	Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Milazzo, 2010[41]	20	20	Neoplastic disease, alcohol abuse	Median CD4 count 490 (NAC) and 524 (placebo)	Side effects in text	Not reported	Not reported	Not reported	Not reported	"Two patients reported insomnia one in the NAC and one in the control group."	Not reported	Not reported
Spada, 2002[42]	10	10			No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Breitkreutz, 2000[43]	21	16	CD4 >200, endocrine disease, liver cirrhosis, serum creatine >1.5 mg/dl, cardiorespiratory insufficiency, substance or alcohol abuse		No side effects reported	3	2	0	0		Not reported	Not reported
Breitkreutz, 2000[43]	16	13	CD4 200-500, endocrine disease, liver cirrhosis, serum creatine >1.5 mg/dl, cardiorespiratory insufficiency, alcohol or drug abuse		No side effects reported	2	0	0	0		Not reported	Not reported
de Rosa, 2000[44]	41	42	CD4>500	Mean CD4 count 203 (NAC), 160 (placebo)	Side effects in text	10	12	Not reported	Not reported	"No evidence of toxicity associated with NAC administration was found"	Not reported	Not reported
Walmsley, 1998 [45]	96	102	CD4>200	Concurrent cotrimoxazole use, mean CD4 count 148 (NAC) and 160 (control)	Side effects reported	23	17	11	Not reported		Not reported	Not reported
Look, 1998[46]	24	24	CD4<200, opportunistic infection, abnormal laboratory findings		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Grant, 2000[47]	73	74	HIV/HBV co-infection		No side effects reported	not reported	not reported	Not reported	Not reported		Not reported	Not reported

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Withdrawn (NAC)	Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Neri, 2000[48]	39	38	Renal insufficiency, cancer, respiratory distress, alcohol abuse, obesity, DM		Side effects in text	Not reported	Not reported	Not reported	Not reported	"No adverse reaction to, nor side effects of, treatment, necessitated suspension of IFN or NAC"	Not reported	Not reported
Look, 1999 [49]	8	8	Cirrhosis, prior IFN treatment, renal disorder, HIV/HBV coinfection, other causes of chronic liver disease, contraindication against IFN-therapy		Side effects in text	Not reported	Not reported	0	0	"The medication was well tolerated by all patients and no patient stopped therapy due to side effects"	Not reported	Not reported
Ideo, 1999[50]	58	62	Decompensated cirrhosis, steroid therapy, HIV co-infection, drug addiction, psychosis, malignancy		Side effects reported	9	10	9	10		Not reported	Not reported
Cimino, 1998[51]	12	13			No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Tripi, 1998 [52]	14	7			No side effects reported	0	0	0	0	"No serious side effects were observed"	Not reported	Not reported
Bernhard, 1998[53]	19	17			No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Purwanto, 2012[54]	16	16	Stage V diabetic nephropathy, steroids, malignancy, obstructive uropathy		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Hashemi, 2012[55]	35	35	Creatinine > 1.8mg/dl		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Renke, 2010/2008 [56-57]	20	20	Steroids or immunosuppressive therapy		Side effects in text	1	1	Not reported	Not reported	"NAC therapy was well tolerated by all patients. Adverse effects were not reported"	Not reported	Not reported

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Withdrawn (NAC)	Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Nascimento, 2009[58]	12	10	Chronic inflammatory disease, DM, active infection, hepatitis B or C		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Hsu, 2009[59]	38	227	Vitamin C, malignancy, active infections, haematological disorders, recent blood transfusion or surgery, renal transplantation	44% diabetic nephropathy	No side effects reported	14	not reported	Not reported	Not reported		Not reported	Not reported
Tepel, 2003[60]	64	70		31% DM	Side effects in text	Not reported	Not reported	Not reported	Not reported	"Five patients (8%) reported GI discomfort during treatment with NAC. No major side effects were observed"	14	14
Popora, 201[61]	45	47	Steroids		Side effects in text	Not reported	Not reported	Not reported	Not reported	"NAC was well tolerated by all patients and no adverse reactions were reported"	Not reported	Not reported
Hashim, 2009[62]	95	97	Other causes of infertility, DM, smoking, alcohol use, age>40		No side effects reported	0	0	0	0		Not reported	Not reported
Elnashar, 2007[63]	30	31	History of pelvic pelvic surgery, infertility other than anovulation, endocrine disorders		No side effects reported	2	1	2	1		Not reported	Not reported
Shahin, 2009[64]	140	140	>35 years, <20 years, threatened abortion in the current pregnancy	Pregnant (mean gestational age 17wks)	Side effects reported	16	0	16	10		Not reported	Not reported
Amin, 2008[65]	80	86	Consanguineous marriage, uterine anatomic abnormality, positive antibodies for antiphospholipid syndrome, endocrine abnormality	Pregnant (treatment started once pregnancy confirmed)	Side effects in text	2	0	Not reported	Not reported		Not reported	Not reported

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Withdrawn (NAC)	Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Safarinejad, 2009[66]	118	118	History of cancer, genital disease, STI, smokers, hepatobiliary disease, renal disease, neurological or psychiatric disease		Side effects in text	13	12	0	0		Not reported	Not reported
Ciftci, 2009[67]	60	60	Varicocele, leukospermia, hormonal abnormalities, obstruction		Side effects in text	0	0	0	00	"None of the patients in the present study reported any side effects with use of the drug"	Not reported	Not reported
Lai, 2012[68]	18	9	Pregnancy, breastfeeding, chronic infection, serious co-morbidities (e.g. diabetes)		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
van Dieten, 2003[69]	67	64			No side effects reported	5	3	Not reported	Not reported		Not reported	Not reported
Perez, 2003[70]	74	71	More than one limb involved, surgery, pregnancy	37% smokers	Side effects in text	18	15	Not reported	Not reported	"The most prominent side effects were sulphur like taste and stomach reaction"	Not reported	Not reported
Yalcin, 2002[71]	43	36			Side effects in text	1	0	1	0	"In one patient oral NAC was discontinued because of diarrhoea. Other side effects of the drug included minor nausea in one patient and minor nasal leak in another"	Not reported	Not reported
Currie, 1988[72]	8	8			No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Furst, 1978[73]	11	11	Malignant hypertension, acute renal failure		Side effects in text	4	4	3	1		Not reported	Not reported

Supplement table 3: Co-morbidities, side effects, withdrawals and death in studies reporting long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Exclusion criteria	Co-morbidities	Side effects reported	Withdrawn (NAC)	Withdrawn (placebo)	Attributable Withdraw (NAC)	Attributable Withdraw (Placebo)	Side effects	Death (NAC)	Death (placebo)
Kasperczyk, 2014[74]	120	49		55% smokers	No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Dabirmoghadam, 2013[75]	30	30	<12, positive history of drug reaction to NAC, history of previous reflux treatment, laryngeal cancer		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Khoshbaten, 2010[76]	15	15			No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Martina, 2008[77]	12	12	Women, smokers, secondary hypertension, cancer, hepatic, pulmonary, renal disease, psychiatric disorder		Side effects in text	1	1	Not reported	Not reported	"No adverse events were noted during the treatment"	Not reported	Not reported
Pace, 2003[78]	16	5	<15, pregnancy, history of stroke, HIV		No side effects reported	Not reported	Not reported	Not reported	Not reported		Not reported	Not reported
Adair, 2001[79]	25	22	Dementia, alcohol or substance abuse, major depressive disorder, Hachinski ischemic score >4		Side effects reported	2	2	0	0		Not reported	Not reported
Estensen, 1999[80]	34	30			Side effects in text	1	1	1	0	"Only one patient reported an adverse reaction (headaches)"	Not reported	Not reported
de Flora, 1997[81]	133	129	Chronic respiratory disease	11% smokers	Side effects reported	7	7	1	1		Not reported	Not reported
Ardissino, 1997[82]	45	46	>75, coronary artery bypass, valvular heart disease, congenital heart disease, symptomatic cerebrovascular disease, anaemia, fever, infections, hypertension, thyrotoxicosis	30% smokers, 5% diabetics, 53% hypertension	Side effects reported	8	4	Not reported	Not reported		Not reported	Not reported

Supplement table 4: Specific side effects in studies reporting on long-term NAC use

[illegible]

Supplement table 4: Specific side effects in studies reporting on long-term NAC use, con't

Author, year	Total (NAC)	Total (placebo)	Abdominal pain (NAC)	Abdominal pain (Placebo)	Nausea vomiting (NAC)	Nausea vomiting (Placebo)	Diarrhoea (NAC)	Diarrhoea (Placebo)	Headache (NAC)	Headache (Placebo)	Arthralgia (NAC)	Arthralgia (Placebo)	Rash (NAC)	Rash (Placebo)	Dizziness (NAC)	Dizziness (Placebo)	Cramps (NAC)	Cramps (Placebo)	Drowsiness (NAC)	Drowsiness (Placebo)
Pela, 1999[26]	85	84	2	2			1	0					0	1						
Rasmussen, 1988[29]	59	57	10	6									5	4						
Shahin, 2009[64]	140	140	28	0	26	0														
Stafanger, 1989[32]	52	52	1	1									1	0						
Walmsley, 1998[45]	96	102			4	1							20	25						
Zheng, 2014[16]	504	502	15	17			5	3			2	0			4	9				
Ardissino, 1997[82]	45	46	5	0			4	0	1	2										
LaRowe, 2013[13]	78	33	31	19					12	3	6	2	5	5					4	3

Supplement table 5: Risk of bias assessment for studies reporting on long-term NAC use

Author, year	Sequence generation	Allocation concealment	Blinding-participant	Blinding-investigator	Complete outcome data - side effect	Complete outcome data - withdraw	complete outcome data - death	Selective outcome reporting
Berk, 2012[1]	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear
Berk,2008[2]	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear
Berk, 2014[3]	Unclear	Unclear	Low risk	Low risk	Unclear	Low risk	Unclear	Unclear
Farokhnia, 2013[4]	Unclear	Unclear	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk
Carmeli, 2012 [5]	Unclear	Unclear	Low risk	Low risk	High risk	Unclear	Unclear	Unclear
Lavoie, 2008 [6]	Unclear	Unclear	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear
Berk, 2008[7]	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	Unclear
Afshar, 2012[8]	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear
Ghanizadeh, 2013[9]	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear
Bloch,2013[10]	Unclear	Low risk	Low risk	Low risk	Unclear	Low risk	Unclear	Unclear
Grant, 2009 [11]	Low risk	Low risk	Low risk	Unclear	Low risk	Unclear	Unclear	Unclear
Grant, 2007[12]	High risk	High risk	High risk	High risk	Unclear	Unclear	Unclear	Unclear
LaRow, 2013[13]	Unclear	Unclear	Unclear	Unclear	Low risk	Low risk	Unclear	Unclear
Gray, 2012[14]	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear
Grant,2010[15]	Unclear	Unclear	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear
Zheng, 2014[16]	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear
Tse, 2013[17]	Unclear	Low risk	Low risk	Low risk	Unclear	Unclear	Low risk	Unclear
de Backer, 2013 [18]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Patil, 2011[19]	Unclear	Unclear	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear
Stav, 2009 ^a [20]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Schermer, 2009[21]	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk	Low risk	Unclear
Decramer, 2005[22]	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk	Unclear	Low risk
de Benedetto, 2005[23]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
van Overveld, 2005[24]	High risk	High risk	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Kasielski, 2001[25]	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear	Unclear
Pela, 1999 [26]	Unclear	Unclear	Unclear	Unclear	Low risk	Low risk	Unclear	Unclear

Supplement table 5: Risk of bias assessment for studies reporting on long-term NAC use, con't


Author, year	Sequence generation	Allocation concealment	Blinding-participant	Blinding-investigator	Complete outcome data - side effect	Complete outcome data - withdraw	Complete outcome data - death	Selective outcome reporting
Lukas, 2005[27]	Unclear	Unclear	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear
Hansen, 1994[28]	Low risk	Unclear	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear
Rasmussen, 1988[29]	Low risk	Unclear	Unclear	Unclear	Low risk	Low risk	Unclear	Unclear
Poder, 1984[30]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
McGavin, 1985[31]	Low risk	Unclear	Unclear	Unclear	Low risk	Low risk	Unclear	Unclear
Stafanger, 1989[32]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Stafanger, 1988[33]	Unclear	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear
Ratjen, 1985[34]	Low risk	Unclear	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear
Mitchell, 1982[35]	Unclear	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Stafanger, 1988[33]	Unclear	Unclear	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear
Martinez, 2014[36]	Low risk	Unclear	Unclear	Unclear	Low risk	Low risk	Low risk	Low risk
Demedts, 2005[37]	Low risk	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear	Low risk
Ghanei, 2008[38]	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear
Van Zandwijk, 2000[39]	Low risk	Unclear	Unclear	Unclear	Low risk	Unclear	Low risk	Unclear
Van Schooten, 2002[40]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Milazzo, 2010[41]	High risk	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear	Unclear
Spada, 2002[42]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Breitkreutz, 2000[43]	Low risk	Unclear	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear
Breitkreutz, 2000[43]	Unclear	Low risk	Low risk	Low risk	Unclear	Low risk	Low risk	Unclear
de Rosa, 2000[44]	Low risk	Low risk	High risk	High risk	Low risk	Low risk	Unclear	Unclear
Walmsley, 1998 [45]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Look, 1998[46]	Unclear	Unclear	Low risk	Unclear	High risk	Unclear	Unclear	Unclear
Neri, 2000[48]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Look, 1999 [49]	Unclear	Unclear	High risk	High risk	Unclear	Unclear	Unclear	Unclear
Ideo, 1999[50]	Unclear	Unclear	High risk	High risk	Low risk	Unclear	Unclear	Unclear
Cimino, 1998[51]	High risk	High risk	High risk	Unclear	Unclear	Unclear	Unclear	Unclear

Supplement table 5: Risk of bias assessment for studies reporting on long-term NAC use, con't

Author, year	Sequence generation	Allocation concealment	Blinding-participant	Blinding-investigator	complete outcome data - side effect	complete outcome data - withdraw	complete outcome data - death	Selective outcome reporting
Tripi, 1998 [52]	Low risk	Unclear	High risk	Unclear	Unclear	Unclear	Unclear	Unclear
Bernhard, 1998[53]	Unclear	Unclear	Unclear	Unclear	High risk	High risk	Unclear	Unclear
Purwanto, 2012[54]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Hashemi, 2012[55]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Renke, 2010/2008 [56-57]	Low risk	Low risk	High risk	High risk	Unclear	Low risk	Unclear	Unclear
Nascimento, 2009[58]	High risk	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear
Hsu, 2009[59]	High risk	High risk	High risk	High risk	Unclear	Low risk	Unclear	Unclear
Tepel, 2003[60]	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	Low risk	
Popora, 201[61]	High risk	High risk	High risk	Unclear	Unclear	Unclear	Unclear	Unclear
Hashim, 2009[62]	Low risk	Low risk	High risk	High risk	Unclear	Unclear	Unclear	Unclear
Elnashar, 2007[63]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Shahin, 2009[64]	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear
Amin, 2008[65]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Safarinejad, 2009	Low risk	Unclear	Unclear	Unclear	Unclear	Low risk	Unclear	Unclear
Safarinejad, 2009[66]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Ciftci, 2009[67]	Unclear	Low risk	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear
Lai, 2012[68]	Low risk	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
van Dieten, 2003[69]	Low risk	Low risk	Low risk	Low risk	Unclear	Low risk	Unclear	Unclear
Perez, 2003[70]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Yalcin, 2002[71]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Currie, 1988[72]	Low risk	Unclear	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear
Kasperczyk, 2014[74]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Dabirmoghaddam, 2013[75]	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	Unclear	Unclear
Khoshbaten, 2010[76]	Unclear	Low risk	High risk	High risk	Unclear	Unclear	Unclear	Unclear

Supplement table 5: Risk of bias assessment for studies reporting on long-term NAC use, con't

Author, year	Sequence generation	Allocation concealment	Blinding-participant	Blinding-investigator	Complete outcome data - side effect	Complete outcome data - withdraw	Complete outcome data - death	Selective outcome reporting
Martina, 2008[77]	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Pace, 2003[78]	Unclear	Unclear	Low risk	Unclear	Unclear	Low risk	Unclear	Unclear
Adair, 2001[79]	Unclear	Unclear	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear
Estensen, 1999[80]	Unclear	Unclear	Low risk	Low risk	Unclear	Low risk	Unclear	Unclear
de Flora, 1997[81]	Low risk	Unclear	Low risk	Unclear	Low risk	Unclear	Unclear	Unclear
Ardissino, 1997[82]	Unclear	Unclear	Low risk	Unclear	Low risk	Unclear	Low risk	Unclear



Supplement figure 1: Selection process for the inclusion of studies

References

1. Berk M, Dean OM, Cotton SM, et al. Maintenance N-acetyl cysteine treatment for bipolar disorder: A double-blind randomized placebo controlled trial. *BMC Medicine* 2012; 10.
2. Berk M, Copolov DL, Dean O, et al. N-Acetyl Cysteine for Depressive Symptoms in Bipolar Disorder-A Double-Blind Randomized Placebo-Controlled Trial. *Biological Psychiatry* 2008; 64: 468-75.
3. Berk M, Dean OM, Cotton SM, et al. The Efficacy of Adjunctive N-Acetylcysteine in Major Depressive Disorder: A Double-Blind, Randomized, Placebo-Controlled Trial. *Journal of Clinical Psychiatry* 2014; 75: 628-U95.
4. Farokhnia M, Azarkolah A, Adinehfar F, et al. N-acetylcysteine as an adjunct to risperidone for treatment of negative symptoms in patients with chronic schizophrenia: A randomized, double-blind, placebo-controlled study. *Clinical Neuropharmacology* 2013; 36: 185-92.
5. Carmeli C, Knyazeva MG, Cuenod M, Do KQ. Glutathione precursor N-acetyl-cysteine modulates EEG synchronization in schizophrenia patients: A double-blind, randomized, placebo-controlled trial. *PLoS ONE* 2012; 7.
6. Lavoie S, Murray MM, Deppen P, et al. Glutathione precursor, N-acetyl-cysteine, improves mismatch negativity in schizophrenia patients. *Neuropsychopharmacology* 2008; 33: 2187-99.
7. Berk M, Copolov D, Dean O, et al. N-Acetyl Cysteine as a Glutathione Precursor for Schizophrenia-A Double-Blind, Randomized, Placebo-Controlled Trial. *Biological Psychiatry* 2008; 64: 361-8.
8. Afshar H, Roohafza H, Mohammad-Beigi H, et al. N-acetylcysteine add-on treatment in refractory obsessive-compulsive disorder: A randomized, double-blind, placebo-controlled trial. *Journal of Clinical Psychopharmacology* 2012; 32: 797-803.
9. Ghanizadeh A, Moghimi-Sarani E. A randomized double blind placebo controlled clinical trial of N-Acetylcysteine added to risperidone for treating autistic disorders. *BMC Psychiatry* 2013; 13.
10. Bloch MH, Panza KE, Grant JE, Pittenger C, Leckman JF. N-acetylcysteine in the treatment of pediatric trichotillomania: A randomized, double-blind, placebo-controlled add-on trial. *Journal of the American Academy of Child and Adolescent Psychiatry* 2013; 52: 231-40.
11. Grant JE, Odlaug BL, Kim SW, Suck WK. N-acetylcysteine, a glutamate modulator, in the treatment of trichotillomania: a double-blind, placebo-controlled study. *Archives of General Psychiatry* 2009; 66: 756-63.
12. Grant JE, Kim SW, Odlaug BL. N-Acetyl Cysteine, a Glutamate-Modulating Agent, in the Treatment of Pathological Gambling: A Pilot Study. *Biological Psychiatry* 2007; 62: 652-7.
13. LaRowe SD, Kalivas PW, Nicholas JS, Randall PK, Mardikian PN, Malcolm RJ. A double-blind placebo-controlled trial of N-acetylcysteine in the treatment of cocaine dependence. *American Journal on Addictions* 2013; 22: 443-52.
14. Gray KM, Carpenter MJ, Baker NL, et al. A double-blind randomized controlled trial of N-acetylcysteine in cannabis-dependent adolescents. *American Journal of Psychiatry* 2012; 169: 805-12.
15. Grant JE, Odlaug BL, Kim SW. A double-blind, placebo-controlled study of N-acetyl cysteine plus naltrexone for methamphetamine dependence. *European Neuropsychopharmacology* 2010; 20: 823-8.
16. Zheng JP, Wen FQ, Bai CX, et al. Twice daily N-acetylcysteine 600 mg for exacerbations of chronic obstructive pulmonary disease (PANTHEON): a randomised, double-blind placebo-controlled trial. *Lancet Respiratory Medicine* 2014; 2: 187-94.
17. Tse HN, Raiteri L, Wong KY, et al. High-dose N-acetylcysteine in stable COPD: The 1-year, double-blind, randomized, placebo-controlled HIACE study. *Chest* 2013; 144: 106-18.
18. De Backer J, Vos W, Van Holsbeke C, et al. Effect of high-dose N-acetylcysteine on airway geometry, inflammation, and oxidative stress in COPD patients. *International Journal of COPD* 2013; 8.
19. Patil AB, Kale AB, Singhal SS, Khan TA. Study of malondialdehyde as an indicator of oxidative stress and its modulation by N-acetylcysteine in chronic obstructive pulmonary disease. *Journal of Clinical and Diagnostic Research* 2011; 5: 48-51.
20. Stav D, Raz M. Effect of N-acetylcysteine on air trapping in COPD: A randomized placebo-controlled study. *Chest* 2009; 136: 381-6.
21. Schermer T, Chavannes N, Dekhuijzen R, et al. Fluticasone and N-acetylcysteine in primary care patients with COPD or chronic bronchitis. *Respiratory medicine* 2009; 103: 542-51.
22. Decramer M, Rutten-van Molken M, Dekhuijzen PNR. Effects of N-acetylcysteine on outcomes in chronic obstructive pulmonary disease (Bronchitis Randomized on NAC Cost-Utility Study, BRONCUS): a randomised placebo-controlled trial (vol 365, pg 1552, 2005). *Lancet* 2005; 366: 984.
23. De Benedetto F, Aceto A, Dragani B, et al. Long-term oral n-acetylcysteine reduces exhaled hydrogen peroxide in stable COPD. *Pulmonary Pharmacology and Therapeutics* 2005; 18: 41-7.

24. Van Overveld FJ, Demkow U, Gorecka D, De Backer WA, Zielinski J. New developments in the treatment of COPD: Comparing the effects of inhaled corticosteroids and N-acetylcysteine. *Journal of Physiology and Pharmacology* 2005; 56: 135-42.
25. Kasielski M, Nowak D. Long-term administration of N-acetylcysteine decreases hydrogen peroxide exhalation in subjects with chronic obstructive pulmonary disease. *Respiratory medicine* 2001; 95: 448-56.
26. Pela R, Calcagni AM, Subiaco S, Isidori P, Tubaldi A, Sanguinetti CM. N-acetylcysteine reduces the exacerbation rate in patients with moderate to severe COPD. *Respiration* 1999; 66: 495-500.
27. Lukas R, Scharling B, Schultze-Werninghaus G, Gillissen A. Administration of N-acetylcysteine and Vitamin C to augment antioxidant protection in patients with chronic bronchitis. [German]. *Deutsche medizinische Wochenschrift* 2005; 130: 563-7.
28. Hansen NCG, Skriver A, Brorsen-Riis L, et al. Orally administered N-acetylcysteine may improve general well-being in patients with mild chronic bronchitis. *Respiratory medicine* 1994; 88: 531-5.
29. Rasmussen JB, Glennow C. Reduction in days of illness after long-term treatment with N-acetylcysteine controlled-release tablets in patients with chronic bronchitis. *European Respiratory Journal* 1988; 1: 351-5.
30. Poder G, Puskas J, Kelemen J. Acetylcysteine in chronic obstructive bronchitis. [German]. *Therapiewoche* 1984; 34: 7047-52.
31. Anonymous, McGavin CR, Prescott RJ, Nariman S, Macfarlane JT. Oral N-acetylcysteine and exacerbation rates in patients with chronic bronchitis and severe airways obstruction. *Thorax* 1985; 40: 832-5.
32. Stafanger G, Koch C. N-acetylcysteine in cystic fibrosis and *Pseudomonas aeruginosa* infection: clinical score, spirometry and ciliary motility. *European Respiratory Journal* 1989; 2: 234-7.
33. Stafanger G, Garne S, Howitz P, Morkassel E, Koch C. The clinical effect and the effect on the ciliary motility of oral N-acetylcysteine in patients with cystic fibrosis and primary ciliary dyskinesia. *European Respiratory Journal* 1988; 1: 161-7.
34. Ratjen F, Wonne R, Posselt HG, Stover B, Hofmann D, Bender SW. A double-blind placebo controlled trial with oral ambroxol and N-acetylcysteine for mucolytic treatment in cystic fibrosis. *European Journal of Pediatrics* 1985; 144: 374-8.
35. Mitchell EA, Elliott RB. Controlled trial of oral N-acetylcysteine in cystic fibrosis. *Australian Paediatric Journal* 1982; 18: 40-2.
36. Martinez FJ, de Andrade JA, Anstrom KJ, King TE, Raghu G, Idiopathic Pulm Fibrosis C. Randomized Trial of Acetylcysteine in Idiopathic Pulmonary Fibrosis. *New England Journal of Medicine* 2014; 370: 2093-101.
37. Demedts M, Behr J, Buhl R, et al. High-dose acetylcysteine in idiopathic pulmonary fibrosis. *New England Journal of Medicine* 2005; 353: 2229-42.
38. Ghanei M, Shohrati M, Jafari M, Ghaderi S, Alaeddini F, Aslani J. N-acetylcysteine improves the clinical conditions of mustard gas-exposed patients with normal pulmonary function test. *Basic and Clinical Pharmacology and Toxicology* 2008; 103: 428-32.
39. van Zandwijk N, Dalesio O, Pastorino U, de Vries N, van Tinteren H. EUROSCAN, a randomized trial of vitamin A and N-acetylcysteine in patients with head and neck cancer or lung cancer. For the European Organization for Research and Treatment of Cancer Head and Neck and Lung Cancer Cooperative Groups. *Journal of the National Cancer Institute* 2000; 92: 977-86.
40. Van Schooten FJ, Nia AB, De Flora S, et al. Effects of oral administration of N-Acetyl-L-cysteine: A multi-biomarker study in smokers. *Cancer Epidemiology Biomarkers and Prevention* 2002; 11: 167-75.
41. Milazzo L, Menzaghi B, Caramma I, et al. Effect of antioxidants on mitochondrial function in HIV-1-related lipoatrophy: A pilot study. *AIDS Research and Human Retroviruses* 2010; 26: 1207-14.
42. Spada C, Treitinger A, Reis M, et al. The effect of N-acetylcysteine supplementation upon viral load, CD4, CD8, total lymphocyte count and hematocrit in individuals undergoing antiretroviral treatment. *Clinical Chemistry & Laboratory Medicine* 2002; 40: 452-5.
43. Breitzkreutz R, Pittack N, Nebe CT, et al. Improvement of immune functions in HIV infection by sulfur supplementation: Two randomized trials. *Journal of Molecular Medicine* 2000; 78: 55-62.
44. De Rosa SC, Zaretsky MD, Dubs JG, et al. N-acetylcysteine replenishes glutathione in HIV infection. *European Journal of Clinical Investigation* 2000; 30: 915-29.
45. Walmsley SL, Khorasheh S, Singer J, Djurdjev O. A randomized trial of N-acetylcysteine for prevention of trimethoprim- sulfamethoxazole hypersensitivity reactions in pneumocystis carinii Pneumonia prophylaxis (CTN 057). *Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology* 1998; 19: 498-505.
46. Look MP, Rockstroh JK, Rao GS, et al. Sodium selenite and N-acetylcysteine in antiretroviral-naïve HIV-1-infected patients: a randomized, controlled pilot study. *European Journal of Clinical Investigation* 1998; 28: 389-97.

47. Grant PR, Black A, Garcia N, Prieto J, Garson JA. Combination therapy with interferon-alpha plus N-acetyl cysteine for chronic hepatitis C: A placebo controlled double-blind multicentre study. *Journal of Medical Virology* 2000; 61: 439-42.
48. Neri S, Ierna D, Antoci S, Campanile E, D'Amico RA, Noto R. Association of alpha-interferon and acetyl cysteine in patients with chronic C hepatitis. *Panminerva medica* 2000; 42: 187-92.
49. Look MP, Gerard A, Rao GS, et al. Interferon/antioxidant combination therapy for chronic hepatitis C--a controlled pilot trial. *Antiviral Research* 1999; 43: 113-22.
50. Ideo G, Bellobuono A, Tempini S, et al. Antioxidant drugs combined with alpha-interferon in chronic hepatitis C not responsive to alpha-interferon alone: A randomized, multicentre study. *European Journal of Gastroenterology & Hepatology* 1999; 11: 1203-7.
51. Cimino L, Belisario MA, Intrieri M, et al. Effect of N-acetyl-cysteine on lymphomonocyte glutathione and response to interferon treatment in C-virus chronic hepatitis. *Italian Journal of Gastroenterology and Hepatology* 1998; 30: 189-93.
52. Tripi S, Di Gaetano C, Soresi M, et al. Acetylcysteine therapy for chronic hepatitis C. Are its effects synergistic with interferon alpha? A pilot study. *Clinical Drug Investigation* 1998; 16: 297-302.
53. Bernhard MC, Junker E, Hettinger A, Lauterburg BH. Time course of total cysteine, glutathione and homocysteine in plasma of patients with chronic hepatitis C treated with interferon-alpha with and without supplementation with N-acetylcysteine. *Journal of Hepatology* 1998; 28: 751-5.
54. Purwanto B, Prasetyo DH. Effect of oral N-acetylcysteine treatment on immune system in continuous ambulatory peritoneal dialysis patients. *Acta medica Indonesiana* 2012; 44: 140-4.
55. Hashemi SR, Noshad H, Tabrizi A, et al. Angiotensin receptor blocker and N-Acetyl Cysteine for reduction of proteinuria in patients with type 2 diabetes mellitus. *Iranian Journal of Kidney Diseases* 2012; 6: 39-43.
56. Renke M, Tylicki L, Rutkowski P, et al. The effect of N-acetylcysteine on proteinuria and markers of tubular injury in non-diabetic patients with chronic kidney disease: A placebo-controlled, randomized, open, cross-over study. *Kidney and Blood Pressure Research* 2008; 31: 404-10.
57. Renke M, Tylicki L, Rutkowski P, et al. The effect of N-acetylcysteine on blood pressure and markers of cardiovascular risk in non-diabetic patients with chronic kidney disease: A placebo-controlled, randomized, cross-over study. *Medical Science Monitor* 2010; 16: 13-8.
58. Nascimento MM, Suliman ME, Silva M, et al. Effect of oral N-acetylcysteine treatment on plasma inflammatory and oxidative stress markers in peritoneal dialysis patients: A placebo-controlled study. *Peritoneal Dialysis International* 2010; 30: 336-42.
59. Hsue PY, Hunt PW, Wu Y, et al. Association of abacavir and impaired endothelial function in treated and suppressed HIV-infected patients. *AIDS* 2009; 23(15): 2021-7.
60. Tepel M, van der Giet M, Statz M, Jankowski J, Zidek W. The antioxidant acetylcysteine reduces cardiovascular events in patients with end-stage renal failure: a randomized, controlled trial. *Circulation* 2003; 107(7): 992-5.
61. Porpora MG, Brunelli R, Costa G, et al. A promise in the treatment of endometriosis: An observational cohort study on ovarian endometrioma reduction by N-acetylcysteine. *Evidence based Complementary and Alternative Medicine* 2013; 240702.
62. Abu Hashim H, Anwar K, Abd El-Fatah R, et al. N-acetyl cysteine plus clomiphene citrate versus metformin and clomiphene citrate in treatment of clomiphene-resistant polycystic ovary syndrome: a randomized controlled trial. *Journal of Women's Health* 2013; 19: 2043-8.
63. Elnashar A, Fahmy M, Mansour A, Ibrahim K. N-acetyl cysteine vs. metformin in treatment of clomiphene citrate-resistant polycystic ovary syndrome: a prospective randomized controlled study. *Fertil Steril* 2007; 88(2): 406-9.
64. Shahin AY, Hassanin IMA, Ismail AM, Kruessel JS, Hirchenhain J. Effect of oral N-acetyl cysteine on recurrent preterm labor following treatment for bacterial vaginosis. *International Journal of Gynecology and Obstetrics* 2009; 104: 44-8.
65. Amin AF, Shaaban OM, Bediawy MA. N-acetyl cysteine for treatment of recurrent unexplained pregnancy loss. *Reproductive BioMedicine Online* 2008; 17: 722-6.
66. Safarinejad MR, Safarinejad S. Efficacy of Selenium and/or N-Acetyl-Cysteine for Improving Semen Parameters in Infertile Men: A Double-Blind, Placebo Controlled, Randomized Study. *Journal of Urology* 2009; 181: 741-51.
67. Ciftci H, Verit A, Savas M, Yeni E, Erel O. Effects of N-acetylcysteine on Semen Parameters and Oxidative/Antioxidant Status. *Urology* 2009; 74: 73-6.

68. Lai ZW, Hanczko R, Bonilla E, et al. N-Acetylcysteine Reduces Disease Activity by Blocking Mammalian Target of Rapamycin in T Cells From Systemic Lupus Erythematosus Patients A Randomized, Double-Blind, Placebo-Controlled Trial. *Arthritis and Rheumatism* 2012; 64: 2937-46.
69. Van Dielen HEM, Perez RSGM, Van Tulder MW, et al. Cost effectiveness and cost utility of acetylcysteine versus dimethyl sulfoxide for reflex sympathetic dystrophy. *Pharmacoeconomics* 2003; 21: 139-48.
70. Perez RSGM, Zuurmond WWA, Bezemer PD, et al. The treatment of complex regional pain syndrome type I with free radical scavengers: a randomized controlled study. *Pain* 2003; 102: 297-307.
71. Yalcin E, Altin F, Cinhuseyinoglu F, Arslan MO. N-acetylcysteine in chronic blepharitis. *Cornea* 2002; 21: 164-8.
72. Currie DC, Greenstone M, Pavia D, et al. Efficacy of 'mucoregulatory' agents in Young's syndrome. *Thorax* 1988; 43: 480-1.
73. Furst DE, Clements PJ, Harris R, Ross M, Levy J, Paulus HE. Measurement of clinical change in progressive systemic sclerosis: a 1 year double-blind placebo-controlled trial of N-acetylcysteine. *Annals of the Rheumatic Diseases* 1979; 38: 356-61.
74. Kasperczyk S, Dobrakowski M, Kasperczyk A, et al. Effect of treatment with N-acetylcysteine on non-enzymatic antioxidant reserves and lipid peroxidation in workers exposed to lead. *Annals of Agricultural and Environmental Medicine* 2014; 21: 272-7.
75. Dabirmoghaddam P, Amali A, Langroudi MM, Fard MRS, Hejazi M, Razavi MS. The effect of N-acetyl Cysteine on laryngopharyngeal reflux. *Acta Medica Iranica* 2013; 51: 757-64.
76. Khoshbaten M, Aliasgarzadeh A, Masnadi K, et al. N-acetylcysteine improves liver function in patients with non-alcoholic fatty liver disease. *Hepatitis Monthly* 2010; 10: 12-6.
77. Martina V, Masha A, Gigliardi VR, et al. Long-term n-acetylcysteine and l-arginine administration reduces endothelial activation and systolic blood pressure in hypertensive patients with type 2 diabetes. *Diabetes Care* 2008; 31: 940-4.
78. Pace BS, Shartava A, Pack-Mabien A, Mulekar M, Ardia A, Goodman SR. Effects of N-acetylcysteine on dense cell formation in sickle cell disease. *American Journal of Hematology* 2003; 73: 26-32.
79. Adair JC, Knoefel JE, Morgan N. Controlled trial of N-acetylcysteine for patients with probable Alzheimer's disease. *Neurology* 2001; 57: 1515-7.
80. Estensen RD, Levy M, Klopp SJ, et al. N-Acetylcysteine suppression of the proliferative index in the colon of patients with previous adenomatous colonic polyps. *Cancer Letters* 1999; 147: 109-14.
81. De Flora S, Grassi C, Carati L. Attenuation of influenza-like symptomatology and improvement of cell-mediated immunity with long-term N-acetylcysteine treatment. *European Respiratory Journal* 1997; 10: 1535-41.
82. Ardissino D, Merlini PA, Savonitto S, et al. Effect of transdermal nitroglycerin or N-acetylcysteine, or both, in the long-term treatment of unstable angina pectoris. *Journal of the American College of Cardiology* 1997; 29: 941-7.