

CORRESPONDENCE

Response to: What characteristics of primary care and patients are associated with early death in patients with lung cancer in the UK?

We appreciate the paper by O'Dowd *et al*¹ as an interesting contribution to the literature concerning the primary care diagnosis of lung cancer. By concentrating on those patients dying soon after diagnosis, this paper is addressing the issue of access to care of patients with lung cancer who are already gravely ill. The disappointingly low chest X-ray (CXR) referral rates (we calculate obtained in only 40.2% and 50.4% of the patients attending with medium/high frequency and dying before and after 90 days, respectively) may represent irregular recording of CXR requests in The Health Improvement Network (THIN). Electronic research databases store their data in several fields, and CXR requests do not always appear in the main clinical field, sometimes being relegated to 'hidden text'. In a study of paper and electronic records, Stapley *et al* found a primary care CXR was obtained in 164/247 (66%) of lung cancer cases;² a second study examined 3184 cases in the GPRD from 2007 to 2009 inclusive and found a CXR in 1947 (61%), though we believed there was too much missing CXR data to submit the latter paper for publication. Thus, the CXR referral rates in the present study do appear unusually low, especially given that data from practices reporting very low referral rates were excluded.

The more important consideration is whether having a 'liberal' primary care CXR policy might facilitate the diagnosis of early-stage and potentially curable lung

cancer. We agree with the authors that the association between higher CXR rates at practice level and increased lung cancer mortality likely reflects better identification of lung cancer as a cause of death as a result of having a CXR. We would be interested to know whether the incidence of lung cancer was higher in practices that ordered CXRs more frequently and whether practice-level CXR rates might yet correlate with higher rates of long-term survival in lung cancer.

The study suggests reliance on primary care to identify advanced, let alone early, cases of symptomatic lung cancer is a failing strategy in isolation despite the high frequency with which these patients interact with primary care. Besides the use of a risk calculator, as suggested by the authors, other interventions include empowering patients to request a CXR, as we attempted in the original El Cid study,³ lowering the threshold suggested by the National Institute of Health and Care Excellence (NICE) for ordering a CXR in primary care⁴ or proactively screening patients with COPD for NICE lung cancer criteria.⁵

Trevor K Rogers,¹ William Hamilton,² Angela Tod,³ Richard Neal⁴

¹Chest Clinic, Doncaster Royal Infirmary, Doncaster, South Yorkshire, UK

²Department of Primary Care Diagnostics, University of Exeter, Exeter, UK

³School of Nursing, Midwifery and Social Work/Central Manchester NHS Foundation Trust, Manchester, UK

⁴Department of Primary Care Medicine, University of Bangor, Wrexham, UK

Correspondence to Dr Trevor K Rogers, Chest Clinic, Doncaster Royal Infirmary, Armthorpe Road, Doncaster, South Yorkshire DN2 5LT, UK; Trevor.rogers@dbh.nhs.uk

Correction notice This article has been corrected since it was publishing Online First. 'CPRD' has been corrected to 'GPRD' for 'General Practice Research Database'. The title has also been amended to read 'Response to: What characteristics of primary care and patients are associated with early death in patients with lung cancer in the UK?'

Contributors TKR drafted the letter after receiving suggestions from AT and WH, who provided data from two of his group's studies. RN then made further comments. All authors approved the article before submission.

Competing interests None.

Provenance and peer review Not commissioned; internally peer reviewed.



CrossMark

To cite Rogers TK, Hamilton W, Tod A, *et al*. *Thorax* 2015;**70**:184.

Received 5 November 2014

Accepted 2 December 2014

Published Online First 22 December 2014



► <http://dx.doi.org/10.1136/thoraxjnl-2014-205692>

► <http://dx.doi.org/10.1136/thoraxjnl-2014-206638>

Thorax 2015;**70**:184.

doi:10.1136/thoraxjnl-2014-206514

REFERENCES

- O'Dowd EL, McKeever TM, Baldwin DR, *et al*. What characteristics of primary care and patients are associated with early death in patients with lung cancer in the UK? *Thorax* 2015;**70**: 161–8.
- Stapley S, Sharp D, Hamilton W. Negative chest X-rays in primary care patients with lung cancer. *Br J Gen Pract* 2006;**56**:570–3.
- Athey VL, Suckling RJ, Tod AM, *et al*. Early diagnosis of lung cancer: evaluation of a community-based social marketing intervention. *Thorax* 2012;**67**: 412–17.
- Hurt CN, Roberts K, Rogers TK, *et al*. A feasibility study examining the effect on lung cancer diagnosis of offering a chest X-ray to higher-risk patients with chest symptoms: protocol for a randomized controlled trial. *Trials* 2013;**14**:405.
- Zarmouh A, Bowen D, Boon M, *et al*. Prospective assessment of incidental respiratory symptoms in COPD (PAIRS-COPD), feasibility study. *European Respiratory Society Annual Congress*. September 2014. https://http://www.ersnetsecure.org/public/prg_congress_abstract?ww_i_presentation=66694



THORAX

Response to: What characteristics of primary care and patients are associated with early death in patients with lung cancer in the UK?

Trevor K Rogers, William Hamilton, Angela Tod and Richard Neal

Thorax 2015 70: 184 originally published online December 22, 2014
doi: 10.1136/thoraxjnl-2014-206514

Updated information and services can be found at:
<http://thorax.bmj.com/content/70/2/184>

References

These include:

This article cites 3 articles, 2 of which you can access for free at:
<http://thorax.bmj.com/content/70/2/184#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>