

AUDIT, RESEARCH AND GUIDELINE UPDATE

# Developmental determinants in non-communicable chronic diseases and ageing

J Bousquet, <sup>1,2,3,4</sup> J M Anto, <sup>4,5,6,7,8</sup> K Berkouk, <sup>9</sup> P Gergen, <sup>10</sup> J Pinto Antunes, <sup>11</sup> P Augé, <sup>3,12</sup> T Camuzat, <sup>3,13</sup> J Bringer, <sup>3,14</sup> J Mercier, <sup>3,15,16</sup> N Best, <sup>3,17</sup> R Bourret, <sup>3,18</sup> M Akdis, <sup>4,19</sup> S H Arshad, <sup>20</sup> A Bedbrook, <sup>3</sup> C Berr, <sup>3,21</sup> A Bush, <sup>22</sup> G Cavalli, <sup>3,23</sup> M A Charles, <sup>24</sup> F Clavel-Chapelon, <sup>25</sup> M Gillman, <sup>26,27</sup> D R Gold, <sup>28</sup> M Goldberg, <sup>29</sup> J W Holloway, <sup>30</sup> P Iozzo, <sup>31</sup> S Jacquemin, <sup>3,32</sup> C Jeandel, <sup>3,33</sup> F Kauffmann, <sup>4,34</sup> T Keil, <sup>4,35,36</sup> G H Koppelman, <sup>4,37</sup> S Krauss-Etschmann, <sup>38</sup> D Kuh, <sup>39</sup> S Lehmann, <sup>3,40</sup> K C Lodrup Carlsen, <sup>4,41</sup> D Maier, <sup>4,42</sup> M Méchali, <sup>23</sup> E Melén, <sup>4,43</sup> J P Moatti, <sup>44</sup> I Momas, <sup>4,45,46</sup> P Nérin, <sup>3,47</sup> D S Postma, <sup>4,48</sup> K Ritchie, <sup>3,49</sup> J M Robine, <sup>3,50,51,52</sup> B Samolinski, <sup>53</sup> V Siroux, <sup>4,54,55,56</sup> P E Slagboom, <sup>57,58</sup> H A Smit, <sup>4,59</sup> J Sunyer, <sup>4,5,6,7,8</sup> R Valenta, <sup>4,60</sup> P Van de Perre, <sup>61</sup> J M Verdier, <sup>3,62,63,64</sup> M Vrijheid, <sup>5,7,8</sup> M Wickman, <sup>4,43</sup> P Yiallouros, <sup>65</sup> M Zins<sup>66</sup>

► Additional material is published online only. To view please visit the journal online (http://dx.doi.org/10.1136/ thoraxjnl-2014-206304). **ABSTRACT** 

TC representing Mr C Bourquin, President of the Région Languedoc Roussillon (C Bourquin, deceased author)

For numbered affiliations see end of article.

#### Correspondence to

Professor Jean Bousquet, 273 av d'Occitanie, Montpellier 34090, France; fabienne.portejoie@gmail.com

Received 14 October 2014 Revised 22 October 2014 Accepted 24 October 2014 Published Online First 23 January 2015 Prenatal and peri-natal events play a fundamental role in health, development of diseases and ageing (Developmental Origins of Health and Disease (DOHaD)). Research on the determinants of active and healthy ageing is a priority to: (i) inform strategies for reducing societal and individual costs of an ageing population and (ii) develop effective novel prevention strategies. It is important to compare the trajectories of respiratory diseases with those of other chronic diseases.

Prenatal and peri-natal events play a fundamental role in health, development of diseases and ageing (Developmental Origins of Health and Disease (DOHaD)). Research on the determinants of active and healthy ageing (AHA) is a priority to: (i) inform strategies for reducing societal and individual costs of an ageing population and (ii) develop effective novel prevention strategies.

The European Union (EU) leads a global effort to understand the early determinants of ageing. The Polish Presidency of the EU Council (2011) targeted chronic respiratory diseases in children to promote AHA. The developmental determinants of chronic diseases in ageing were reinforced during the Cyprus Presidency of the EU Council (2012). Several projects of the EU Sixth and Seventh Framework Programme for Research and Technological Development (FP6 and FP7) were funded to understand the mechanisms of prenatal and early life events on the development of chronic diseases. One of the action plans of the European Innovation Partnership on AHA is devoted to integrated care for chronic diseases, and chronic respiratory diseases have been selected as the model for chronic diseases.

As part of this research initiative, a meeting in Montpellier on 2 and 3 December 2013 attempted to better understand the early life events that may impact ageing in health and disease, with a particular focus on respiratory diseases. The results of

the FP6 and FP7 projects on population-based cohorts in adults and old age were reviewed to propose novel research, policies and value creation. The full report is in the online supplement and on the MACVIA-LR website (macvia.cr-languedocroussillon.fr).

## IMPORTANCE OF LIFE COURSE MODELS TO UNDERSTAND HEALTHY AGEING AND DISEASE

Biological ageing is the progressive deterioration of function that occurs in the postmaturity phase and can be assessed at the individual, physiological systems and cellular levels.<sup>2</sup> Cross cohort comparisons of British birth cohorts show that social and biological factors from early life onwards can affect ageing, in particular respiratory function and disease.

In order to detect health-promoting factors, the early life events of diseases such as COPD need to be understood (European COST action BM1201: Developmental Origins of Chronic Respiratory Disease<sup>3</sup>). Susceptibility to COPD is associated with markers of foetal growth and early childhood disadvantage. Lung function at birth is determined by in utero processes.

Future research needs to develop a life course model of ageing that integrates the rather separate research on specific diseases or clinical conditions, functional ageing and well-being and should include socio-economic characteristics.

# BIRTH COHORTS FOCUSING ON CHRONIC RESPIRATORY DISEASES IN THE GENERAL POPULATION

Asthma and allergic diseases begin early in life. Birth cohort studies are relevant to investigate environmental and lifestyle determinants of asthma and IgE-associated diseases or the absence of such diseases. Over 30 community-based birth cohorts



**To cite:** Bousquet J, Anto JM, Berkouk K, *et al. Thorax* 2015;**70**:595–597.



focusing on asthma and allergies have started in Europe and many have been pooled.<sup>4</sup> The Network of Excellence GA<sup>2</sup>LEN (FP6, contract No FOOD-CT-2004-506378) has initiated birth cohort harmonisation. Two concerted actions, CHICOS (FP7 No. 241604) and ENRIECO (FP7 grant agreement No 226285), have built a network of more than 70 birth cohorts across Europe. Mechanisms of the Development of ALLergy (MeDALL, FP7 No. 261357) attempts to generate novel knowledge on the mechanisms of allergy onset and proposes early diagnosis, prevention and targets for therapy based on a new standardised follow-up of birth cohorts in Europe.

Numerous cohort studies have shown that early decrements in spirometry persist into late middle age, underscoring the crucial importance of early life influences. Optimal lung development depends on normal airway function at birth as well as normal growth during childhood up to the plateau of spirometric function around 25 years of age. After this, lung function declines as a normal event.

The geographical and temporal diversities of birth cohorts in MeDALL provide an excellent opportunity to study the effects of living conditions in different places since these are likely to be major determinants in the causation of the wave of chronic diseases currently observed.

An NIAID, NHLBI and MeDALL joint workshop (September 2012)<sup>5</sup> identified over 60 birth cohorts focusing on asthma and allergy worldwide in order to (i) document the knowledge obtained, (ii) identify the knowledge gaps and inconsistencies and (iii) develop strategies for moving forward. Following the workshop in 2012, an online database containing information about existing cohorts was created to facilitate collaboration (AsthmaBirthCohorts.niaid.nih.gov).

Some population-based birth cohorts (PIAMA, BAMSE, ECA, MAS) have information on chronic diseases and their risk factors and can assess chronic diseases and AHA across the life cycle.

## NUTRITION, OBESITY AND DIABETES ACROSS THE LIFE COURSE

Chronic diseases often start early in life and their processes expand over the life course. DORIAN (Developmental ORigins of healthy and unhealthy AgeiNg: The Role of Maternal Obesity, FP7) aims at linking studies of early developmental processes with those of ageing from a life course perspective.

A better knowledge on the developmental origin of obesity and type-2 diabetes may lead to preventive measures in order to reduce the global epidemic of these two diseases.

Nutrition is vital for health and disease prevention. The Supplementation with Antioxidant Vitamins and Minerals (SU. VI.MAX) study was a controlled trial (12 741 persons) followed up for 8 years (1994–2002) to test the efficacy of dietary manipulations in lowering the incidence of chronic diseases and cancer. No supplementation was administered to participants in SU.VI.MAX 2. Food intake and nutritional factors in infancy and childhood have also been studied in several of the ongoing European birth cohorts, which allows for unique longitudinal association analyses on various chronic diseases.

Nutrition is vital to understand the development of chronic diseases and the promotion of healthy ageing.

#### ADULT COHORTS IN CHRONIC DISEASES

The French prospective cohort (E3N), the EPIC-France cohort, is composed of women who were under a health insurance plan, the *Mutuelle Générale de l'Education Nationale*, for schoolteachers and coworkers. In 1990, about 100 000 women (40–65 years) were recruited. Epidemiological research in nutrition, hormones and chronic diseases (asthma, diabetes or osteoporosis) studied the risk factors for developing these diseases.

Two large European epidemiological cohorts, recruited in the early 1990s and followed up for 20 years, have been carried out in respiratory health: (i) the European Community Respiratory Health Survey (ECRHS; http://www.ecrhs.org/), including 18 668 individuals aged 20–44 years at baseline and (ii) the Epidemiological study of the Genetics and Environment of Asthma, bronchial hyperresponsiveness and atopy (EGEA; http://egeanet.vjf.inserm.fr), a French case-control and family study of adults and children at baseline, including 2120 individuals aged 7–70 years at baseline. Combined analyses between these cohorts have been performed.

COPACETIC, a cohort of the lung cancer screening trial with over 2200 individuals with lung function and CT scans, identified genes associated with lung function decline in interaction with smoking, CT scan-based emphysema and coronary calcifications.

LifeLines (The Netherlands) is a three-generation cohort of 165 000 individuals followed annually by questionnaire and, every 5 years, by extensive functional screening of the five prime research areas: (1) metabolic/hormonal, (2) heart/vessel/kidney, (3) lung/respiratory/allergy, (4) psychiatric and (5) musculoskeletal.

Adult cohorts include participants that have been followed up for a long period of time, and may be used to link the development of factor profiles of chronic diseases from early life.

#### **OLD AGE COHORTS**

Neuropsychiatric disorders of old-age (eg, depression, Alzheimer's disease) may be determined by earlier risk exposure. The DEVELAGE consortium (FP7) aims to characterise shared molecular pathways between the early developmental processes in the brain and brain ageing.

This example underlines the limits of old age cohorts in the study of chronic neuropsychiatric disorder in the elderly.

Integrated research on DEvelopmental determinants of Ageing and Longevity (IDEAL; http://www.ideal-ageing.eu, FP7) examines the role of epigenetic regulation and transmission to next generations. A unique human cohort and animal studies are studied to discover novel longevity pathways and the links between development and ageing.

The Three-City Study (3C Study) is a population-based longitudinal study of the relation between vascular diseases and dementia in persons aged 65 years and older. A total of 9294 participants of both sexes were recruited from three French cities.

CONSTANCES, a large general-purpose epidemiological population-based cohort, aims to provide public health information. The cohort is a representative sample of 200 000 French adults, aged 18–69 years at inception.

Cohorts in middle and old age adults can be intertwined with earlier cohorts to understand the mechanisms of AHA and diseases in ageing.

Recommendations for future research are reported in the online supplement and the MACVIA-LR website on (i) integration of omics in epidemiological studies across the life cycle, (ii) performance of longitudinal studies assessing chronic diseases and (iii) value of European birth cohorts to study chronic diseases and AHA. Cohorts with an initial focus on chronic respiratory diseases could be expanded to understand the trajectories of chronic diseases and AHA.

Cohorts across the life cycle are essential to understand AHA. Lifelong birth cohorts give an overview of the problem and are complemented by cohorts carried out at different ages. Birth cohorts initially focusing on chronic respiratory diseases offer highly valuable additive information to understand chronic diseases and AHA.

#### **Author affiliations**

- <sup>1</sup>University Hospital, Montpellier, France
- <sup>2</sup>Inserm U 1168, Paris, France
- <sup>3</sup>Contre les Maladies Chroniques pour un Vieillissement Actif en Languedoc Roussillon, Site de Référence de l'EIP on AHA, Montpellier, France
- <sup>4</sup>MeDALL, Mechanisms of the Development of Allergy, FP7, Amsterdam, Groningen, the Netherlands
- <sup>5</sup>Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain
- <sup>6</sup>IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain
- <sup>7</sup>Universitat Pompeu Fabra (UPF), Barcelona, Spain
- <sup>8</sup>CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain
- <sup>9</sup>Deputy Head of Unit for Medical Research and the Challenge of Ageing, DG Research & Innovation, European Commission, Brussels, Belgium
- <sup>10</sup>National Institute of Allergy and Infectious Diseases, Bethesda, Maryland, USA <sup>11</sup>European Commission, Directorate General for Health and Consumers, Brussels,
- <sup>12</sup>University Montpellier 1, France
- <sup>13</sup>Région Languedoc Roussillon, France
- <sup>14</sup>Montpellier Medical School, France
- <sup>15</sup>Department of Physiology, Montpellier University Hospital, France
- <sup>16</sup>University Montpellier 1, France
- <sup>17</sup>Nimes University Hospital, France
- <sup>18</sup>Montpellier University Hospital, France
- <sup>19</sup>Swiss Institute of Allergy and Asthma Research (SIAF), Davos and University of Zurich, Switzerland
- <sup>20</sup>David Hide Asthma and Allergy Research Centre, Isle of Wight, UK
- <sup>21</sup>Inserm, Research Unit U1061, University Montpellier I, Montpellier, France
- <sup>22</sup>Department of Paediatric Respiratory Medicine, Royal Brompton Hospital and National Heart and Lung Institute, Imperial College, London, UK
- <sup>23</sup>Institute of Human Genetics, CNRS, Montpellier, France
- <sup>24</sup>Equipe 10 UMR Inserm-Université Paris-Sud (Centre de recherche en Epidémiologie et Santé des Populations, CESP), Villejuif, France <sup>25</sup>Nutrition, Hormones and Women's Health Team, INSERM UMR-S 1018, Paris-
- South University, Villejuif, France
- <sup>26</sup>Obesity Prevention Program, Department of Population Medicine, Harvard Medical School/Harvard Pilgrim Health Care Institute, Boston, Massachusetts, USA
- <sup>27</sup>Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts,
- <sup>28</sup>Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, and Harvard School of Public Health, UK
- <sup>29</sup>Population-Based Epidemiological Cohorts, INSERM-UVSQ UMS 011, Villejuif,
- <sup>30</sup>Human Development & Health, Faculty of Medicine, University of Southampton, Southampton, UK
- <sup>31</sup>Institute of Clinical Physiology, National Research Council (CNR), Pisa, Italy <sup>32</sup>Horiba, Montpellier, France
- <sup>33</sup>Department of Geriatrics, University Hospital, Montpellier, France
- <sup>34</sup>CESP—Team of Respiratory and Environmental Epidemiology INSERM UMR-S1018, University Paris-Sud, Villejuif, France
- <sup>35</sup>Institute of Social Medicine, Epidemiology and Health Economics, Charité— Universitätsmedizin, Berlin, Germany
- <sup>36</sup>Institute for Clinical Epidemiology and Biometry, Julius Maximilian University of Wuerzburg, Germany
- <sup>37</sup>Department of Pediatric Pulmonology and Pediatric Allergology, University of Groningen, University Medical Center Groningen, Beatrix Children's Hospital, Groningen Research Institute for Asthma and COPD, Groningen, The Netherlands

- <sup>38</sup>Comprehensive Pneumology Center, Ludwig Maximilians University and Helmholtz Zentrum Muenchen, Member of the German Research Center for Lung Research, Großhadern, Germany
- <sup>39</sup>MRC Unit for Lifelong Health and Ageing at UCL, London, UK
- <sup>40</sup>Institut de Médecine Régénératrice et de Biothérapie (I.M.R.B.), University Hospital, INSERM U1040, Montpellier, France
- <sup>41</sup>Department of Paediatrics, University of Oslo and Oslo University Hospital, Oslo, Norway
- <sup>42</sup>Biomax Informatics AG, Planegg, Germany
- <sup>43</sup>Institute of Environmental Medicine, Karolinska Institutet and Sachs' Children's Hospital, Stockholm, Sweden
- 44Aix-Marseille University (AMU), Research Unit 912 AMU/INSERM/IRD Social and Economic Sciences Applied to Health (SESSTIM), France
- <sup>45</sup>Department of Public health and biostatistics, Descartes University, Paris, France <sup>46</sup>Municipal Department of social action, childhood, and health, Paris, France <sup>47</sup>SATT AxLR, Montpellier, France
- <sup>48</sup>Department of Pulmonology, University Medical Center Groningen, University of Groningen, GRIAC Research Institute, Groningen, The Netherlands
- <sup>49</sup>Inserm U1061 Neuropsychiatry, Montpellier and Faculty of Medicine, Imperial College London, London, UK
- <sup>50</sup>Inserm Research Unit 988, Paris, France
- <sup>51</sup>Inserm Research Unit 710, Montpellier, France
- <sup>52</sup>Ecole Pratique des Hautes Etudes (EPHE), Paris, France
- 53Department of Prevention of Environmental Hazards and Allergology, Medical University of Warsaw, Warsaw, Poland
- <sup>54</sup>Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, University Grenoble Alpes, IAB, Grenoble, France
- <sup>55</sup>Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, INSERM, IAB, Grenoble, France
- <sup>56</sup>Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, CHU de Grenoble, IAB, Grenoble, France
- <sup>57</sup>Section of Molecular Epidemiology, Leiden University Medical Center, Leiden, the Netherlands
- <sup>58</sup>Consortium for Healthy Aging, Leiden University Medical Center, Leiden, the Netherlands
- <sup>59</sup>Julius Center of Health Sciences and Primary Care, University Medical Center Utrecht, University of Utrecht, Utrecht, the Netherlands
- <sup>60</sup>Division of Immunopathology, Department of Pathophysiology and Allergy Research, Center for Pathophysiology, Infectiology and Immunology, Medical University of Vienna, Vienna, Austria
- <sup>61</sup>University Hospital and INSERM U 1058, Montpellier, France
- <sup>62</sup>EPHE, Section des Sciences de la Vie et de la Terre, Paris, France
- <sup>63</sup>UMR S 710, University Montpellier 2, Montpellier, Paris, France
- <sup>64</sup>Institut Transdisciplinaire d'Etudes du Vieillissement, Montpellier, France
- <sup>65</sup>Cyprus International Institute for Environmental & Public Health in Association with Harvard School of Public Health, Cyprus University of Technology, Limassol, Cyprus <sup>66</sup>Director of Population-Based Epidemiological Cohorts, INSERM-UVSQ UMS 011,
- Villejuif, France

Correction notice This article has been corrected since it was published Online First. The statement 'PG and JPA did not participate in the discussion' on the title page has been removed. It has now been corrected and included in the 'Contributors' statement.

Contributors All authors participated in the meeting. All authors contributed to the discussion, except for KB and JPA.

**Funding** The meeting was supported by an unrestricted educational grant from the Région Languedoc Roussillon awarded by Mr. C Bourquin (President).

**Competing interests** None declared.

Provenance and peer review Not commissioned; internally peer reviewed.

#### REFERENCES

- Samolinski B, Fronczak A, Wlodarczyk A, et al. Council of the European Union conclusions on chronic respiratory diseases in children. Lancet 2012;379:e45-6.
- Kuh D, Cooper R, Hardy R, et al. A life course approach to healthy ageing. Oxford: Oxford University Press, 2014.
- Krauss-Etschmann S, Bush A, Bellusci S, et al. Of flies, mice and men: a systematic approach to understanding the early life origins of chronic lung disease. Thorax 2013;68:380-4.
- Bousquet J, Anto J, Sunyer J, et al.; MeDALL Study Group; CHICOS Study Group; ENRIECO Study Group; GA2LEN Study Group. Pooling birth cohorts in allergy and asthma: European Union-funded initiatives—a MeDALL, CHICOS, ENRIECO, and GA(2)LEN joint paper. Int Arch Allergy Immunol 2013;161:1-10.
- Bousquet J, Gern JE, Martinez FD, et al. Birth cohorts in asthma and allergic diseases: Report of a NIAID/NHLBI/MeDALL joint workshop. J Allergy Clin Immunol 2014;133:1535-46.

## Developmental determinants in non-communicable chronic diseases and ageing

THORAXJNL-2014-206304

European Innovation Partnership on Active and Healthy Ageing: DG Sanco and DG CNECT
Framework Programme 7, DG Research and Innovation
National Institute of Allergy and Infectious Diseases (NIH)

Proposal following the recommendations of the EU Council of the Polish (2011) and Cyprus (2012) Presidencies

Montpellier, 2-3 December 2013

J Bousquet (1-4), JM Anto (4-8), K Berkouk (9)\*, P Gergen (10)\*, J Pinto Antunes (11), P Augé (3, 12), T Camuzat (3, 13)\*\*, J Bringer (3, 14), J Mercier (3, 15, 16), N Best (3, 17), R Bourret (3, 18), M Akdis (4, 19), SH Arshad (20), A Bedbrook (3), C Berr (3, 21), A Bush (22), G Cavalli (3, 23), MA Charles (24), F Clavel-Chapelon (25), M Gillman (26), DR Gold (27), M Goldberg (28), JW Holloway (29), P Iozzo (30), S Jacquemin (3, 31), C Jeandel (3, 32), F Kauffmann (4, 61), T Keil (4, 33), GH Koppelman (4, 34), S Krauss-Etschmann (35), D Kuh (36), S Lehmann (3, 37), KC Lodrup Carlsen (4, 38), D Maier (4, 39), M Méchali (23), E Melén (4, 40), JP Moatti (41), I Momas (4, 42, 43), P Nérin (3, 44), DS Postma (4, 45), K Ritchie (3, 46), JM Robine (3, 47-49), B Samolinski (50), V Siroux (4, 51), PE Slagboom (52), HA Smit (4, 53), J Sunyer (4-8), R Valenta (4, 54), P Van de Perre (55), JM Verdier (3, 56-58), M Vrijheid (5, 7, 8), M Wickman (4, 40), P Yiallouros (59), M Zins (60).

- \*: did not participate in the discussion
- \*\*: Representing Mr. C Bourquin, President of the Région Languedoc Roussillon
- 1. University hospital, Montpellier, France
- 2. Inserm U1018, Villejuif, France
- 3. MACVIA-LR. Contre les Maladies Chroniques pour un Vieillissement Actif en Languedoc Roussillon, Site de Référence de l'EIP on AHA
- 4. MeDALL, Mechanisms of the Development of Allergy, FP7
- 5. Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain
- 6. IMIM (Hospital del Mar Medical Research Institute, Barcelona, Spain
- 7. Universitat Pompeu Fabra (UPF), Barcelona, Spain
- 8. CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain
- 9. Deputy Head of Unit for Medical Research and the Challenge of Ageing, DG Research & Innovation, European Commission, Brussels
- 10. National Institute of Allergy and Infectious Diseases, Bethesda, MD, USA
- 11. European Commission, Directorate General for Health and Consumers, Belgium
- 12. President, University Montpellier 1, France
- 13. Assitant Director General, Région Languedoc Roussillon, France
- 14. Dean, Montpellier Medical School, France
- 15. Department of Physiology, Montpellier University Hospital, France
- 16. Vice President for Research, University Montpellier 1, France
- 17. Assitant Director General, Nimes University Hospital, France
- 18. Directeur Général Adjoint, Montpellier University Hospital, France
- 19. Swiss Institute of Allergy and Asthma Research (SIAF), Davos and University of Zurich, Switzerland.

- 20. David Hide Asthma and Allergy Research Centre, Isle of Wight, United Kingdom
- 21. Inserm, Research Unit U1061, Montpellier, and University Montpellier I, France 22. Department of Paediatric Respiratory Medicine, Royal Brompton Hospital and National Heart and Lung Institute, Imperial College, London, UK.
- 23. Institute of Human Genetics, CNRS, Montpellier, France
- 24. Equipe 10 UMR Inserm-Université Paris-Sud (Centre de recherche en Epidémiologie et Santé des Populations, CESP), Villejuif, France25. Nutrition, Hormones and Women's Health Team, INSERM UMR-S 1018, Paris-South University, Villejuif, France 26. Obesity Prevention Program, Department of Population Medicine, Harvard Medical School/Harvard Pilgrim Health Care Institute, Boston, MA, USA, Department of Nutrition, Harvard School of Public Health, Boston, M, USA
- 27. Channing Division of Network Medicine, Brigham and Women's Hospital, Department of Medicine, Harvard Medical School, and Harvard School of Public Health
- 28. Population-Based Epidemiological Cohorts, INSERM-UVSQ UMS 011, Villejuif, France
- 29. Human Development & Health, Faculty of Medicine, University of Southampton, Southampton, UK.
- 30. Institute of Clinical Physiology, National Research Council (CNR), Pisa, Italy.
- 31. Horiba, Montpellier, France
- 32. Department of Geriatrics, Montpellier University hospital, Montpellier, France
- 33. Institute of Social Medicine, Epidemiology and Health Economics, Charité Universitätsmedizin Berlin, Berlin, Germany; Institute for Clinical Epidemiology and Biometry, Julius Maximilian University of Wuerzburg, Germany
- 34. University of Groningen, University Medical Center Groningen, Department of Pediatric Pulmonology and Pediatric Allergology, Beatrix Children's Hospital, Groningen Research Institute for Asthma and COPD, Groningen, The Netherlands.
- 35. Comprehensive Pneumology Center, Ludwig Maximilians University and Helmholtz Zentrum Muenchen, Großhadern, Germany; Member of the German Research Center for Lung Research
- 36. MRC Unit for Lifelong Health and Ageing at UCL, London, England
- 37. Institut de Médecine Régénératrice et de Biothérapie (I.M.R.B.), University hospital, INSERM U1040, Montpellier, France.
- 38. University of Oslo and Oslo University Hospital, Department of Paediatrics, Oslo, Norway.
- 39. Biomax Informatics AG, Planegg, Germany
- 40. Institute of Environmental Medicine, Karolinska Institutet and Sachs' Children's Hospital, Stockholm, Sweden
- 41. Professor of Economics, Aix-Marseille University (AMU), Research Unit 912 AMU/INSERM/IRD Social and Economic Sciences Applied to Health (SESSTIM).
- 42. Department of Public health and biostatistics, Paris Descartes University, EA 4064
- 43. Paris municipal Department of social action, childhood, and health, Paris, France
- 44. SATT AxLR, Montpellier, France
- 45. University of Groningen, Department of Pulmonology, University Medical Center Groningen, GRIAC Research Institute, Groningen, The Netherlands
- 46. Inserm U1061 Neuropsychiatry, Montpellier and Faculty of Medicine, Imperial College London, UK
- 47. Inserm Research Unit 988, Paris, France
- 48. Inserm Research Unit 710, Montpellier, France
- 49. Ecole Pratique des Hautes Etudes (EPHE), Paris, France
- 50. Department of Prevention of Environmental Hazards and Allergology, Medical University of Warsaw, Poland
- 51. Univ. Grenoble Alpes, IAB, Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, F-38000 Grenoble, France; INSERM, IAB, Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, Grenoble, France; CHU de Grenoble, IAB, Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, Grenoble, France
- 52. Section of Molecular Epidemiology, Leiden University Medical Center, Leiden, and Netherlands Consortium for Healthy Aging, Leiden University Medical Center, Leiden, The Netherlands
- 53. Julius Center of Health Sciences and Primary Care, University Medical Center Utrecht, University of Utrecht, Utrecht, the Netherlands
- 54. Division of Immunopathology, Department of Pathophysiology and Allergy Research, Center for Pathophysiology, Infectiology and Immunology, Medical University of Vienna, Vienna, Austria
- 55. Univesity hospital and INSERM U 1058, Montpellier, France
- 56. EPHE, Section des Sciences de la Vie et de la Terre, Paris, France
- 57. UMR S 710, University Montpellier 2, Montpellier, Paris
- 58. Institut Transdisciplinaire d'Etudes du Vieillissement, Montpellier, France

- 59. Cyprus International Institute for Environmental & Public Health in Association with Harvard School of Public Health, Cyprus University of Technology, Limassol, Cyprus
- 60. Director of Population-Based Epidemiological Cohorts, INSERM-UVSQ UMS 011, Villejuif, France
- 61. CESP Team of Respiratory and Environmental Epidemiology INSERM UMR-S1018, Univ Paris-Sud, Villejuif, France

#### **Abbreviations**

AHA: Active and healthy ageing

AMICS: Asthma Multicentre Infant Cohort Study

BAMSE: Barn Allergi Milj. Stockholm Epidemiologi Projektet CHICOS: Developing a Child Cohort Research Strategy for

Europe

COPACETIC: COPD Pathology: Adressing Critical gaps, Early Treatment & diagnosis and Innovative Concepts

COPD: Chronic obstructive pulmonary disease

COST: European Cooperation in Science and Technology

DARC:Study: Danish Allergy Research Centre study

DOHaD: Developmental Origins of Health and Disease

E3N: Étude Épidémiologique auprès de femmes de la MGEN (Mutuelle Générale de l'Éducation Nationale)

ECA: Environment and Childhood Asthma

EIP on AHA: European Innovation Partnership on Active and Healthy Ageing

ENRIECO: Environmental Health Risks in European Birth Cohorts

ENRIECO: Environmental Health Risks in European Birth Cohorts EPIC: European Prospective Investigation into Cancer and Nutrition

EU: European Union

FP: Framework Program for Research and Technological Development

GA<sup>2</sup>LEN: Global Allergy and Asthma European Network GINI-B: German Infant Nutritional Intervention-Study KOALA: Kind, Ouder en gezondheid, Aandacht voor Leefstijl en Aanleg

LISA: Influences of Lifestyle-related Factors on the Immunesystem and the development of allergies

MACVIA-LR: Contre les MAladies Chroniques pour un VIeillissement Actif en Languedoc Roussillon (EIP on AHA Reference site)

MAS: Multi-centre Allergy Study

MeDALL:Mechanisms of the Development of ALLergy

NHLBI: National Heart Lung and Blood Institute

NIAID: National Institute of Allergy and Infectious Diseases

NIH: National Institutes of Health

PARIS: Pollution and Asthma Risk: an Infant Study

PIAMA-NHS: The Prevention and Incidence of Asthma and Mite Allergy -Natural History Study

### Introduction

Poor health is largely shaped by chronic diseases which share common risk and socio-economic factors, biological mechanisms and cluster in co-morbidities. These are intertwined with ageing and represent a major cause of frailty [1]. Pre-and peri-natal events play a fundamental role in health, the development of chronic diseases and ageing (Developmental Origins of Health and Disease: DOHaD) [2]. Links between environmental and behavioural factors with epigenetic mechanisms [3-5] and conserved genes [6, 7] have been proposed. Ageing is regulated by several pathways including telomer dysfunction related to the senescence of tissues and cells [8]. Like other organ systems, the immune and metabolic systems are highly vulnerable to environmental changes, and is also highly influenced by internal molecules such as hormones [9] and ageing [10]. Research on the determinants of healthy ageing has become a priority to inform strategies for reducing societal and individual costs of an ageing population and to develop effective novel prevention strategies to promote active and healthy ageing (AHA) [11].

#### 1- European Union activities

The European Union (EU) is at the fore front of the global effort to better understand early determinants of ageing.

The 2010 Council of the European Union prioritised the understanding, prevention and management of chronic diseases [12]. The Polish Presidency of the EU Council (2011) targeted chronic respiratory diseases in children to promote their early recognition, prevention and management to ultimately impact AHA [13]. The developmental determinants of chronic diseases in ageing have been reinforced during the Cyprus Presidency of the EU Council "Healthy ageing across the lifecycle" (2012) [14].

Several projects of the EU Seventh Framework Programme for Research and Technological Development (FP7) have attempted to understand the mechanisms of pre-natal and early life events on the development of chronic diseases and ageing. Epidemiologic studies across the life cycle combined with appropriate omics data may help to define pathways and biomarkers of healthy ageing. Research and innovation are key contributors for achieving the goals of the Europe 2020 strategy: growth, jobs, competitiveness, quality of life and social inclusion. The new EU framework for research, called Horizon 2020, will be one of the main tools to implement the Innovation Union policy. The joint strategic EC-COST workshop "Relationship between genome and epigenome", held 14-15 February 2013 in Brussels, addressed the links between genetic, epigenetic and non-genetic factors throughout the lifespan and across generations, their role in disease susceptibility and disease progression and the associated challenges of data handling/storage and interpretation. The outcomes of the workshop set the vision for future policies and research agendas at European level

European Innovation Partnerships (EIP) aim to enhance EU competitiveness and tackle societal challenges [15]. Active and Healthy Ageing (AHA) is a major societal challenge, common to all European countries and all populations. The EIP on AHA has 6 action plans, one of them devoted to integrated care for chronic diseases.

## 2- Objectives of the meeting

The Région Languedoc Roussillon is deeply involved in chronic diseases and AHA through MACVIA-LR (Reference site of the EIP on AHA) [16, 17], and the Président, Mr C Bourquin, has given an unrestricted grant to support this meeting.

Following the EU efforts over the last 10 years in cohort investigations across the life cycle, a meeting was convened in Montpellier, December 2 and 3, 2013 in order to better understand early life events that may impact ageing in health and disease. The results of the FP6 and FP7 project on pre-natal and early life events in chronic diseases were confronted to population-based cohorts in adults and old age to propose novel research, policies and value creation. The programme and abstracts of the meeting are presented in the online supplement.

## 3- Reducing health inequalities in France as an example of priorities in the EU

Non-communicable diseases represent a WHO, UN and EU priority [18]. Life course epidemiology research is a key priority of the French Institute of Public Health (ISP), the institute covering the public health disciplines (biostatistics, epidemiology and social sciences applied to health) of the Health and Life Sciences Research Alliance (AVIESAN). These priorities are in line with the WHO Commission on Social Determinants of Health [19]. The "French paradox" gives a particular importance to the recommendations of this report for a better understanding of health inequalities and for promoting and evaluating public policies aimed at reducing them. While France ranks high on some health indicators (2<sup>nd</sup> longest life expectancy for females, 7th for males, high quality of its health system, etc...), it is also the one country in Western Europe with mortality and morbidity inequalities [20, 21]. A better understanding of the complex causal chains which translate social differences in health inequalities and higher exposure to morbidity and premature mortality, and how they affect populations at various ages, is therefore a key priority for French public health research [11]. Promoting life-cycle epidemiology and its interdisciplinary dialogue with both social sciences and biomedical sciences is clearly a major need for reaching this goal.

### 4- Importance of life course models to understand healthy ageing and disease

Biological ageing is the progressive deterioration of function that occurs in the post maturity phase, and can be assessed at the individual, physiological systems and cellular levels [22, 23]. In depth studies and cross cohort comparisons of British birth cohort and life course studies are providing growing evidence that social and biological factors from early life onwards can affect the peak function achieved at maturity and the rate of functional decline at all these levels; and this has consequences for wellbeing, quality and length of life [24, 25].

In order to detect health-promoting factors, it would be helpful to understand the inverse, ie factors that promote disease. Multiple independent approaches are currently researching the developmental origins of NCDs. Understanding the early life determinants of diseases such as COPD is relevant and the focus of the European COST action BM1201: Developmental Origins of Chronic Respiratory Disease [26]. COPD is a significant cause of morbidity and mortality. Susceptibility to COPD is associated with markers of foetal growth and markers of early childhood disadvantage. Lung function at birth is determined by *in utero* processes (e.g. low birth weight babies or mothers who smoked during pregnancy). Many genes that are associated with spirometry in adult life are also associated with lung function in childhood and infancy [27].

Future research needs to develop a life course model of ageing that integrates the rather separate research on specific diseases or clinical conditions, functional ageing and wellbeing and include socio-economic characteristics.

## 5- Birth cohorts in the general population

Asthma and allergic diseases, the most common disorders in children, begin early in life, but no effective preventive interventions exist to date. Birth cohort studies are a relevant research strategy to investigate the environmental and lifestyle determinants of asthma and atopic disease as well as the absence of such diseases. They could therefore act in health promoting identification. The prioritised research agenda of WHO for the prevention and control of chronic respiratory diseases also includes follow up of cohorts in developing countries to better understand disease onset, progression or disappearance [28].

More than 30 community-based birth cohorts focusing on asthma and allergies have started in Europe over the past 25 years. By 2014/2015, the first two European birth cohorts on allergy and respiratory diseases from the UK (Isle of Wight 1989) [29-31] and Germany (MAS 1990) [32-38] are reaching 25 years of age, followed by others that are soon approaching 18-20 years, such as BAMSE (Sweden) [39-41], ECA (Norway) [42-46], PIAMA (NL) [47-50], GiniPLUS, LISA (both Germany) [51, 52], AMICS-Menorca (Spain) [53, 54] and DARC (Denmark) [55]. Among the younger European birth cohorts are KOALA (NL) [56, 57], INMA (Spain) [58, 59], PARIS (France) [60, 61], EDEN [62] and others with special interest in environmental risk factors for chronic diseases.

In Europe, the Network of Excellence GA<sup>2</sup>LEN (FP6, contract N° FOOD-CT-2004-506378) [63-65] has initiated the harmonisation of birth cohorts [66]. ENRIECO (FP7 grant agreement N° 226285) [67, 68], another initiative built on the GA<sup>2</sup>LEN experience [69], assessed the effects of outdoor and indoor air pollution, and compared pooled analysis with de-centralised coordinated meta-analysis. The two Concerted Actions, CHICOS (FP7 grant agreement No. 241604) and ENRIECO, have built a network of more than 70 birth cohorts across Europe that are prospectively studying more than 500,000 mothers, fathers and children.

Following the experience of these projects, MeDALL (Mechanisms of the Development of ALLergy, FP7 grant agreement No. 261357) [70, 71] was conceived to generate novel knowledge on the mechanisms of initiation of allergy and to propose early diagnosis, prevention and targets for therapy

based on existing information and a new standardised follow up of birth cohorts in Europe. Using the DataSHaPER methodology [72, 73], historical pooled data were harmonised from 14 birth cohorts, representing over 40,000 participants, covering 130 variables and 125 individual data generation events (follow-ups) requiring almost 3,000 mapping decisions [74].

The MeDALL study found that comorbidities are of major importance for chronic respiratory and allergic diseases in childhood, and that IgE sensitisation is adding its effects on top of them [75]. Such a finding could not have been observed when individual cohorts were analysed due to an insufficient number of subjects. A standardised follow-up questionnaire, interoperable with the historical one, made it possible to follow over 11,000 children, many of them up to the age of 18 years [76]. The harmonisation and knowledge management strategies developed by MeDALL, combining retrospective and prospective data, offer a high statistical power to assess trajectories of lifestyle and environmental exposures and to study multidimensional trajectories of chronic respiratory and allergic symptoms using "unsupervised" statistical techniques. They can now be applied to provide efficient opportunities for collaborative studies across the life cycle. An open data approach can extend the benefits of such collaborations to the full society.

Numerous cohort studies have shown that early decrements in spirometry persist into late middle age, underscoring the crucial importance of early life influences [77, 78]. Optimal lung development depends on normal airway function and birth at birth, and normal growth during childhood up to the plateau of spirometric function around 25 years of age. After this, lung function declines as a normal event.

The geographical and temporal diversities of birth cohorts in MeDALL provide an excellent opportunity to study the effects of living conditions in different places, since these are likely to be major determinants in the causation of the wave of chronic diseases currently observed..

An NIAID, NHLBI, MeDALL joint workshop (September 2012) has provided the opportunity to consider more ambitious initiatives identifying over 60 birth cohorts focusing on asthma and allergy worldwide which have been initiated in the last 30 years in order to (1) document the knowledge that asthma/allergy birth cohorts have provided, (2) identify the knowledge gaps and inconsistencies and (3) develop strategies for moving forward, including potential new study designs and the harmonisation of existing asthma birth cohort data. This database will be updated as more cohorts are identified.

Following the workshop in 2012, an online database containing information about existing cohorts was created to facilitate collaboration (AsthmaBirthCohorts.niaid.nih.gov).

The Project Viva is a U.S. pre-birth cohort study of prenatal and early-life determinants of growth, adiposity, cardiovascular risk factors, cognition, and respiratory disease and intermediate metabolic, inflammatory and epigenetic marks that may be on pathways that influence these outcomes [79-81]. While Project Viva investigators have focused on prenatal maternal diet and dietary supplements as primary exposures, they have ascertained many additional co-exposures and potentially modifying factors both in early-life and in later childhood [82].

In the PIAMA birth cohort study, there is yearly information on height and weight as well as on lifestyle and environmental factors [83]. Starting at age 8, chronic disease parameters were included (eg cholesterol, HbA1c, random glucose, blood pressure, lung function and hyperresponsiveness), and at age 16, also intima media thickness [84]. Similar clinical and epidemiological data are available in other European birth cohorts (e.g. BAMSE, ECA and MAS). These additional measurements make it

possible to study the levels and development of cardio-metabolic endpoints from age 8 onwards through puberty and the associations with early life risk factors.

Some population-based birth cohorts have information on chronic diseases and their risk factors. They can be of great interest to assess chronic diseases and healthy ageing across the life cycle.

## 6- Nutrition, obesity and diabetes across the life course

The global epidemic of obesity was soon followed by an epidemic of type 2 diabetes in adults and it is starting to appear in earlier ages more recently. These chronic diseases have genetic and developmental origins and physiopathological processes expanding over the life course. Although diabetes is usually diagnosed after 50 years of age, obesity has developed across the life cycle. Foetal and early life events are of great importance [85, 86]. As examples, in France, thinness rather than obesity in childhood and adolescence is the most common pathway to type 2 diabetes in middle age and older adults [87]. Obesity in pregnancy promotes accelerated foetal growth and gestational diabetes that increase the offspring susceptibility to childhood obesity and type 2 diabetes at an earlier age.

DORIAN (Developmental ORigins of healthy and unhealthy AgeiNg: The Role of Maternal Obesity, FP7 NO 278603) aims at linking studies of early developmental processes with those of ageing from a life course perspective. As a consequence of maternal obesity, the offspring will present early disease features, predisposing to the development of cardiovascular disease, type 2 diabetes and cognitive impairment. The project focuses on the involvement of insulin resistance and glucocorticoid overexposure, oxidative stress or inflammation, telomere shortening, and epigenetic changes.

A better knowledge in the developmental origin of obesity and type-2 diabetes may lead to preventive measures in order to reduce the global epidemic of these two diseases.

Nutrition is vital for health and disease. The Supplementation with Antioxidant Vitamins and Minerals (SU.VI.MAX) Study was a randomised double-blind, placebo-controlled trial including 12,741 persons followed-up for 8 years (1994–2002) to test the efficacy of dietary manipulations in lowering the incidence of cancer, ischemic heart disease, and overall mortality [88, 89]. At the end of the trial phase, participants were invited to participate in an extended follow-up on a voluntary basis. 6,850 subjects were included in the observational SU.VI.MAX 2 Study (2007–2009) for a new clinical examination focusing on cognitive and functional capacities. In addition, participants completed a self-administered questionnaire providing information about health and life-style [90]. No supplementation was administered to participants in SU.VI.MAX 2. Food intake and nutritional factors in infancy and childhood have also been studied in several of the ongoing European birth cohorts, which allows for unique longitudinal association analyses on various chronic diseases.

Nutrition is vital to understand the development of chronic diseases and the promotion of healthy ageing.

#### 7- Adult cohorts in chronic diseases

The French prospective cohort (E3N) [91], the EPIC-France cohort [92, 93], is composed of women who were under a health insurance plan, MGEN (*Mutuelle Générale de l'Education Nationale*), for schoolteachers and co-workers. In 1990, about 100,000 women aged 40-65 years were recruited. Blood samples were collected from approximately 25% of the participants, and saliva samples from almost 50%. About 75,000 of the E3N women answered the diet history questionnaire sent in 1993. Analytical epidemiological research in nutrition, hormones and chronic diseases (e.g. asthma [16],

diabetes [94] or osteoporosis [95]) has been carried out to understand the risk factors for developing cancer and other major diseases.

Two large European epidemiological cohorts, recruited in the early 1990s and followed up for 20 years, have been carried out in respiratory health. The European Community Respiratory Health Survey (ECRHS, <a href="http://www.ecrhs.org/">http://www.ecrhs.org/</a>, FP3) is an international population-based study of asthma and allergy, including 18,668 individuals aged 20-44 yr at baseline [96, 97]. The Epidemiological study of the Genetics and Environment of Asthma, bronchial hyperresponsivness and atopy (EGEA, <a href="http://egeanet.vjf.inserm.fr">http://egeanet.vjf.inserm.fr</a>) is a French case—control and family study of adults and children at baseline, including 2,120 individuals aged 7-70 yr at baseline [98, 99]. Combined analyses between these cohorts have been made [100]. The studies can appraise the trajectories of participants concerning respiratory diseases (onset, persistence, disappearance).

COPACETIC (FP7 N° 201379) is a cohort of the lung cancer screening trial with over 2,200 individuals with lung function and CT scans. The study identified genes associated with lung function decline in interaction with smoking, CT scan based emphysema and coronary calcifications [101].

LifeLines (The Netherlands) is a three-generation cohort of 165,000 individuals followed annually by questionnaire and, each 5 years, by extensive functional screening of the 5 prime research areas (1) metabolic/hormonal, (2) heart/vessel/kidney, (3) lung/respiratory/allergy, (4) psychiatric and (5) musculoskeletal [102].

Adults cohorts have followed up participants for a long period of time and may be used to link development of factor profiles of chronic diseases from early life.

## 8- Old age cohorts

Neuropsychiatric disorders of old-age may be determined by earlier risk exposure. Late-life depression is modulated by exposure to childhood adversity interacting with genetic vulnerability, and the principal risk factors for Alzheimer's disease appear to occur principally in early adulthood [103, 104]. Current knowledge is insufficient to identify the transition of normal brain ageing into Alzheimer-like brain damage. The aim of the DEVELAGE consortium is to characterise shared molecular pathways between early developmental processes in the brain and brain ageing.

This example underlines the limits of old age cohorts in the study of chronic neuropsychiatric disorder in the elderly.

Integrated research on DEvelopmental determinants of Ageing and Longevity (IDEAL; <a href="www.ideal-ageing.eu">www.ideal-ageing.eu</a>, FP7) examines the role of epigenetic regulation and transmission to next generations. A unique human cohort and animal studies (including fruitflies and Xenopus) are being studied in order to discover novel longevity pathways and to determine whether they affect the link between development and ageing [105, 106]. To find set-points of epigenetic control, genome-scale analyses and studies of specific gene systems such as histone methylation of homeobox genes at different embryonic stages are compared. Epigenetic responses are linked with phenotypic consequences of early developmental conditions with biomarkers of the ageing rate.

The Three-City Study (3C Study) is a population-based longitudinal study of the relation between vascular diseases and dementia in persons aged 65 years and older [107, 108, 109]. A total of 9,294 participants of both sexes were recruited from three French cities: Bordeaux (South-West), Dijon (North-East) and Montpellier (South-East). The 3C Study started in 1999 and produced information on life habits, nutrition, hypertension, depression, genetics, other risk factors of dementia and neuroimaging.

CONSTANCES is a large general-purpose epidemiologic population-based cohort in epidemiologic research to provide public health information. The cohort is a representative sample of 200,000 French adults aged 18-69 years at inception. Data are prospectively collected from different sources: annual self-administered questionnaire, periodic health examinations, linkage to health and social national administrative databases. Subjects over 45 years will have a comprehensive work-up of functional physical and cognitive capacities. The data include social and demographic characteristics, social status, life events, behaviours, and occupational factors.

Cohorts in middle and old age adults can be intertwined with earlier cohorts to understand the mechanisms of healthy ageing or diseases.

## 9- From science to policies and value creation

#### K Berkouk

The European Innovation Partnership for Active and healthy Ageing reflects a growing awareness that better care in an ageing society calls for innovative ways to address the needs of older people and improved understanding of the mechanisms of ageing.

#### 10- Recommendations

Recommendations for future research were developed in 3 breakout sessions.

### Breakout Session 1: How to integrate omics in epidemiologic studies across the life cycle

Population-based studies enriched by patient populations are of interest to be used for sampling of omics. Tissues and cells (Peripheral blood cells, T and B cells, airway epithelial cells can be studied in many biological fluids and biopsies [110], but most cohorts only have blood and/or urine samples as they are easy to collect, process, and store, and also carry the strongest implications for intervention at the population level. Recently, several cohorts (VIVA, PIAMA) have implemented collection of nasal epithelial cells to be able to address the role of airway epithelial mechanisms.

To integrate omics across the life cycle, multiple cohorts with information on risk factors and diseases should be investigated at different stages of the life (infants, children, adults, old age). The time of study and trajectories embedding acute and chronic effects of risk factors and disease co-morbidities need to be characterised. As biomarkers can change over time, susceptibility to reverse causality and residual confounding is a potential limitation of these studies.

Omic studies need to be based on research questions. The sample size and availability of biological samples should be carefully determined *a priori*.

Candidate biomarkers and pathways have raised remarkable interest in clinical and epidemiological research [111]. They can improve the understanding molecular mechanisms of diseases, predict models of complex diseases, refine disease phenotypes and guide treatment responses [112, 113]. Much of the research to identify new biomarkers has evolved from studies has focused on genetics and genomics.

- For genomics, whole genome DNA sequencing is now favoured, but many cohorts have historical data from genome-wide association studies (GWAs) [114], which may be used.
- Several types of epigenomic studies can be performed to complement and expand genomic data, including DNA methylation, short and long noncoding RNAs [115], chromatin marks [116] or chromatin contact maps [7, 117].
- Transcriptomic profiling using RNA sequencing or up-to-date microarray platforms can give valuable insights about gene expression patterns related disease and exposure. Studies on

- epigenetics and transcriptomics, may need to be investigated in a specific cell type, given the tissue specificity [118].
- Proteomic analysis is still limited by the sensitivity of the current technology and its ability to expand the information provided by epigenomic analysis may need to be investigated in a specific cell type before adopting it in large-scale applications. Using current proteomics technology, protein wide analysis of cells and tissues is not yet feasible with, coverage of all protein isoforms and post-translational modifications within the large dynamic concentration range of proteins. Therefore, targeted proteins need to be selected for further study. Future initiatives, such as the Human Proteome Project, may change this concept [119].
- Integration of genetic, epigenetic, transcriptomic and proteomic data (integrative genomics) may assist in determining causal networks that lead to disease development [120].
- For all these studies, bioinformatic applications are of vital importance and must be carefully planned right from the beginning of the project strategy.

## Breakout Session 2: Longitudinal studies assessing chronic diseases Breakout Session 2: Longitudinal studies assessing chronic diseases and ageing

D Postma (NL)

It is now widely recognised that ageing is largely determined during the life course, including the prebirth period. One of the main outcomes of the discussion was that aging can be defined as a collapse of compensatory mechanisms. This session contributed to shed some light on several crucial questions by turning traditional projects into innovative ones.

As a basics the following reasons for studying healthy ageing were mentioned: (1) political agenda (consumer costs reduction); (2) biological pathway identification for early intervention and late preventive strategies; (3) personal well being.

The research questions to do so need to be well defined including (1) biological ageing, (2) frailty, (3) aggregated measures of ageing in elderly cohorts and (4) active and healthy ageing.

Several types of cohorts should be studied: including population-based, disease-oriented, case-control studies and GP databases. In these studies, environmental changes and concurrent findings should be confronted with genomics. In long-term studies, the methodology to assess questions, diagnosis and coding of disease may have changed and should be considered. Retrospective and prospective studies do often not provide similar results: so also birth cohorts with prospective longitudinal design are necessary.

Some specific questions need to be answered:

- Can we disentangle the respective contribution of normal advance in age and chronic diseases to the health consequences of ageing?
- What are the most satisfying models of the life course (critical periods, accumulation, pathways)?
- Should we consider different models according to specific chronic diseases (COPD, neurodegenerative diseases, musculoskeletal disorders...)?
- Can we develop beter methods for cross-cohort and cross-data-type integrative analysis?

### Breakout Session 3: Impact of European birth cohorts on chronic diseases and healthy ageing

MeDALL has created a new cross-European birth cohort including now over 35,000 children and their families with clinical data on lung function, body mass index, skin prick tests, house dust samples prospectively collected since early childhood, blood pressure and biomaterial including cord blood, breast milk, serum/plasma, saliva, urine, and stool samples. The prospective regularly collected subjective data include pregnancy (smoking, stress, etc), allergic and respiratory symptoms, doctor

diagnoses, respiratory infections, vaccinations, physical activity, dietary habits, socio-economic characteristics, family lifestyle, home environment, parental and siblings' medical history, etc.

This **unique health data source** will help to address relevant research questions examining:

- The development of respiratory and allergic diseases from childhood to adulthood and old age (sensitisation, disease, lung function decline, COPD).
- The effect of allergy/asthma in young age on lung function decline and COPD in older adults.
- Childhood determinants (e.g. perinatal factors, obesity, dietary habits, physical inactivity, lower social status, stress and environmental exposures) for lung function decline in early adulthood (later leading to chronic resp. diseases)
- The observed links between allergy/asthma and cancer, autoimmune disorders, other inflammatory diseases, infections, heart disease, mental illnesses, etc.
- The links between early life events and chronic diseases using a common questionnaire for chronic diseases, in particular in the 20-25 year old participants of the cohorts.
- The links between early life events, chronic diseases in adulthood and healthy ageing using the harmonisation method of MeDALL.
- The application of large-scale genetic, epigenetic, transcriptomic and proteomic profiling related to onset and progress of chronic diseases.
- Factors associated with lack of disease.

#### References

- Bousquet, J., et al., Systems medicine and integrated care to combat chronic noncommunicable diseases. Genome Med, 2011. 3(7): p. 43.
- Barker, D.J., The fetal and infant origins of adult disease. BMJ, 1990. 301(6761): p. 1111.
- Bousquet, J., et al., Epigenetic inheritance of fetal genes in allergic asthma. Allergy, 2004. 59(2): p. 138-47.
- Koppelman, G.H. and M.C. Nawijn, Recent advances in the epigenetics and genomics of asthma. Curr Opin Allergy Clin Immunol, 2011. 11(5): p. 414-9.
- 5. Lockett, G.A., et al., *Epigenomics and allergic disease*. Epigenomics, 2013. **5**(6): p. 685-99.
- 6. Bousquet, J., H. Yssel, and A.M. Vignola, *Is allergic asthma associated with delayed fetal maturation or the persistence of conserved fetal genes?* Allergy, 2000. **55**(12): p. 1194-7.
- Sexton, T., et al., Three-dimensional folding and functional organization principles of the Drosophila genome. Cell, 2012. 148(3): p. 458-72.
- Newgard, C.B. and N.E. Sharpless, Coming of age: molecular drivers of aging and therapeutic opportunities. J Clin Invest, 2013. 123(3): p. 946-50.
- Akdis, M. and C.A. Akdis, *IgE class switching and cellular memory*. Nat Immunol, 2012. 13(4): p. 312-4.
- Montecino-Rodriguez, E., B. Berent-Maoz, and K. Dorshkind, *Causes, consequences, and reversal of immune system aging*. J Clin Invest, 2013. 123(3): p. 958-65.
- Moatti, J.P., [Social determinants of health inequalities: a public health research priority]. Rev Epidemiol Sante Publique, 2013. 61 Suppl 3: p. S123-5.

- 12. Council conclusions "Innovative approaches for chronic diseases in public health and healthcare systems". COUNCIL OF THE EUROPEAN UNION. 3053rd EMPLOYMENT, SOCIAL POLICY HEALTH and CONSUMER AFFAIRS Council meeting Brussels, 7 December 2010. 2010.
- 13. Samolinski, B., et al., Council of the European Union conclusions on chronic respiratory diseases in children. Lancet, 2012. **379**(9822): p. e45-6.
- 14. Bousquet, J., et al., Impact of early diagnosis and control of chronic respiratory diseases on active and healthy ageing. A debate at the European Union Parliament. Allergy, 2013. **68**(5): p. 555-61.
- European Innovation Partnership on Active and Healthy Ageing (EIP on AHA). <a href="http://ec.europa.eu/health/ageing/innovation/indexen.htm">http://ec.europa.eu/health/ageing/innovation/indexen.htm</a>, 2012.
- 16. Sanchez, M., et al., Temporal asthma patterns using repeated questionnaires over 13 years in a large French cohort of women. PLoS One, 2013. **8**(5): p. e65090.
- 17. European Innovation Partnership on Active & Healthy Ageing: 32 star cities and regions across Europe rewarded for their innovative ideas.

  <a href="http://ec.europa.eu/health/ageing/events/ev\_201307">http://ec.europa.eu/health/ageing/events/ev\_201307</a>
  <a href="http://ec.europa.eu/health/ageing/events/ev\_201307">http://ec.eu/health/ageing/events/ev\_201307</a>
  <a href="http://ec.europa.eu/health/ageing/events/ev\_201307">http://ec.eu/health/ageing/events/ev\_201307</a>
  <a href="http://ec.europa.eu/health/ageing/events/ev
- Beaglehole, R., et al., UN High-Level Meeting on Non-Communicable Diseases: addressing four questions. Lancet, 2011. 378(9789): p. 449-55.
- 19. Marmot, M., et al., WHO European review of social determinants of health and the health divide. Lancet, 2012. **380**(9846): p. 1011-29.
- 20. Jagger, C., et al., Inequalities in healthy life years in the 25 countries of the European Union in 2005: a cross-national meta-regression analysis. Lancet, 2008. 372(9656): p. 2124-31.

- Robine, J.M., et al., The joint action on healthy life years (JA: EHLEIS). Arch Public Health, 2013.
   71(1): p. 2.
- Kuh, D., et al., A life course approach to healthy ageing. 2014: Oxford University Press.
- Hurst, L., et al., Lifetime socioeconomic inequalities in physical and cognitive aging. Am J Public Health, 2013. 103(9): p. 1641-8.
- 24. Power, C., D. Kuh, and S. Morton, From developmental origins of adult disease to life course research on adult disease and aging: insights from birth cohort studies. Annu Rev Public Health, 2013. 34: p. 7-28.
- Wills, A.K., et al., Life course trajectories of systolic blood pressure using longitudinal data from eight UK cohorts. PLoS Med, 2011. 8(6): p. e1000440.
- 26. Krauss-Etschmann, S., et al., Of flies, mice and men: a systematic approach to understanding the early life origins of chronic lung disease. Thorax, 2013. **68**(4): p. 380-4.
- Collins, S.A., et al., HHIP, HDAC4, NCR3 and RARB polymorphisms affect fetal, childhood and adult lung function. Eur Respir J, 2013. 41(3): p. 756-7.
- Bousquet, J., et al., Prioritised research agenda for prevention and control of chronic respiratory diseases. Eur Respir J, 2010. 36(5): p. 995-1001.
- Arshad, S.H., et al., The effect of parental allergy on childhood allergic diseases depends on the sex of the child. J Allergy Clin Immunol, 2012. 130(2): p. 427-34 e6.
- Soto-Ramirez, N., et al., Breastfeeding is associated with increased lung function at 18 years of age: a cohort study. Eur Respir J, 2012. 39(4): p. 985-91
- 31. Yousefi, M., et al., The methylation of the LEPR/LEPROT genotype at the promoter and body regions influence concentrations of leptin in girls and BMI at age 18 years if their mother smoked during pregnancy. Int J Mol Epidemiol Genet, 2013. 4(2): p. 86-100.
- Nickel, R., et al., How should a birth cohort study be organised? Experience from the German MAS cohort study. Paediatr Respir Rev, 2002. 3(3): p. 169-76.
- 33. Illi, S., et al., *The natural course of atopic dermatitis from birth to age 7 years and the association with asthma.* J Allergy Clin Immunol, 2004. **113**(5): p. 925-31.
- 34. Nickel, R., et al., Variability of total serum immunoglobulin E levels from birth to the age of 10 years. A prospective evaluation in a large birth cohort (German Multicenter Allergy Study). Clin Exp Allergy, 2005. 35(5): p. 619-23.
- 35. Illi, S., et al., Perennial allergen sensitisation early in life and chronic asthma in children: a birth cohort study. Lancet, 2006. **368**(9537): p. 763-70.
- 36. Karaolis-Danckert, N., et al., How pre- and postnatal risk factors modify the effect of rapid weight gain in infancy and early childhood on subsequent fat mass development: results from the Multicenter Allergy Study 90. Am J Clin Nutr, 2008. 87(5): p. 1356-64.
- 37. Gruber, C., et al., *History of respiratory infections* in the first 12 yr among children from a birth

- *cohort.* Pediatr Allergy Immunol, 2008. **19**(6): p. 505-12.
- Grabenhenrich, L., et al., Early life determinants of asthma from birth to age 20: a German birth cohort study. J Allergy Clin Immunol, 2014. in press.
- 39. Wickman, M., et al., Quantification of IgE antibodies simplifies the classification of allergic diseases in 4-year-old children. A report from the prospective birth cohort study--BAMSE. Pediatr Allergy Immunol, 2003. **14**(6): p. 441-7.
- 40. Wickman, M., et al., *Childhood-to-adolescence* evolution of IgE antibodies to pollens and plant foods in the BAMSE cohort. J Allergy Clin Immunol, 2013.
- 41. Wickman, M., et al., *The BAMSE project:* presentation of a prospective longitudinal birth cohort study. Pediatr Allergy Immunol, 2002. **13** Suppl 15: p. 11-3.
- 42. Lodrup Carlsen, K.C., *The environment and childhood asthma (ECA) study in Oslo: ECA-1 and ECA-2.* Pediatr Allergy Immunol, 2002. **13 Suppl 15**: p. 29-31.
- 43. Lodrup Carlsen, K.C., et al., *Asthma in every fifth child in Oslo, Norway: a 10-year follow up of a birth cohort study.* Allergy, 2006. **61**(4): p. 454-60.
- 44. Devulapalli, C.S., et al., Severity of obstructive airways disease by age 2 years predicts asthma at 10 years of age. Thorax, 2008. **63**(1): p. 8-13.
- 45. Riiser, A., et al., *Does bronchial hyperresponsiveness in childhood predict active asthma in adolescence?* Am J Respir Crit Care Med, 2012. **186**(6): p. 493-500.
- 46. Hovland, V., et al., *The significance of early recurrent wheeze for asthma outcomes in late childhood.* Eur Respir J, 2013. **41**(4): p. 838-45.
- Hafkamp-de Groen, E., et al., Predicting asthma in preschool children with asthma-like symptoms: Validating and updating the PIAMA risk score. J Allergy Clin Immunol, 2013. 132(6): p. 1303-1310
- 48. Scheffers, F.R., et al., *The association between indoor temperature and body mass index in children: the PIAMA birth cohort study.* BMC Public Health, 2013. **13**(1): p. 1119.
- 49. Wijga, A.H., et al., Cohort profile: The Prevention and Incidence of Asthma and Mite Allergy (PIAMA) birth cohort. Int J Epidemiol, 2013.
- 50. van Rossem, L., et al., Blood pressure in 12-yearold children is associated with fatty acid composition of human milk: the prevention and incidence of asthma and mite allergy birth cohort. Hypertension, 2012. **60**(4): p. 1055-60.
- 51. Kohlboeck, G., et al., Food intake, diet quality and behavioral problems in children: results from the GINI-plus/LISA-plus studies. Ann Nutr Metab, 2012. **60**(4): p. 247-56.
- 52. von Berg, A., et al., Allergies in high-risk schoolchildren after early intervention with cow's milk protein hydrolysates: 10-year results from the German Infant Nutritional Intervention (GINI) study. J Allergy Clin Immunol, 2013. 131(6): p. 1565-73.
- Garcia Algar, O., et al., Concentrations and determinants of NO2 in homes of Ashford, UK and Barcelona and Menorca, Spain. Indoor Air, 2004. 14(4): p. 298-304.

- 54. Friguls, B., et al., [Perinatal exposure to tobacco and respiratory and allergy symptoms in first years of life]. Arch Bronconeumol, 2009. **45**(12): p. 585-90
- 55. Eller, E., et al., *Development of atopic dermatitis in the DARC birth cohort.* Pediatr Allergy Immunol, 2009
- 56. Kummeling, I., et al., *Etiology of atopy in infancy: the KOALA Birth Cohort Study*. Pediatr Allergy Immunol, 2005. **16**(8): p. 679-84.
- Notenboom, M.L., et al., Maternal fatty acid status in pregnancy and childhood atopic manifestations: KOALA Birth Cohort Study. Clin Exp Allergy, 2011. 41(3): p. 407-16.
- Guxens, M., et al., Cohort Profile: The INMA--INfancia y Medio Ambiente--(Environment and Childhood) Project. Int J Epidemiol, 2011.
- Morales, E., et al., Circulating 25-hydroxyvitamin D3 in pregnancy and infant neuropsychological development. Pediatrics, 2012. 130(4): p. e913-20.
- 60. Herr, M., et al., *Does allergic rhinitis exist in infancy? Findings from the PARIS birth cohort.* Allergy, 2011. **66**(2): p. 214-21.
- Ranciere, F., et al., Onset and persistence of respiratory/allergic symptoms in preschoolers: new insights from the PARIS birth cohort. Allergy, 2013.
- 62. Slama, R., et al., *Maternal personal exposure to airborne benzene and intrauterine growth.* Environ Health Perspect, 2009. **117**(8): p. 1313-21.
- 63. Bousquet, J., et al., GA2LEN (Global Allergy and Asthma European Network) addresses the allergy and asthma 'epidemic'. Allergy, 2009. **64**(7): p. 969-77.
- Keil, T., et al., European birth cohort studies on asthma and atopic diseases: I. Comparison of study designs -- a GALEN initiative. Allergy, 2006.
   61(2): p. 221-8.
- Rzehak, P., et al., Body mass index trajectory classes and incident asthma in childhood: results from 8 European Birth Cohorts--a Global Allergy and Asthma European Network initiative. J Allergy Clin Immunol, 2013. 131(6): p. 1528-36.
- 66. Bousquet, J., et al., Pooling Birth Cohorts in Allergy and Asthma: European Union-Funded Initiatives A MeDALL, CHICOS, ENRIECO, and GALEN Joint Paper. Int Arch Allergy Immunol, 2012. 161(1): p. 1-10.
- 67. Gehring, U., et al., Environmental exposure assessment in European birth cohorts: results from the ENRIECO project. Environ Health, 2013. 12: p. 8.
- 68. Tischer, C.G., et al., *Meta-analysis of mould and dampness exposure on asthma and allergy in eight European birth cohorts: an ENRIECO initiative*. Allergy, 2011. **66**(12): p. 1570-9.
- Vrijheid, M., et al., European birth cohorts for environmental health research. Environ Health Perspect, 2012. 120(1): p. 29-37.
- Anto, J.M., et al., Understanding the complexity of IgE-related phenotypes from childhood to young adulthood: a Mechanisms of the Development of Allergy (MeDALL) seminar. J Allergy Clin Immunol, 2012. 129(4): p. 943-54 e4.
- 71. Bousquet, J., et al., MeDALL (Mechanisms of the Development of ALLergy): an integrated approach

- from phenotypes to systems medicine. Allergy, 2011. **66**(5): p. 596-604.
- 72. Fortier, I., et al., Quality, quantity and harmony: the DataSHaPER approach to integrating data across bioclinical studies. Int J Epidemiol, 2010. **39**(5): p. 1383-93.
- 73. Fortier, I., et al., *Is rigorous retrospective* harmonization possible? Application of the DataSHaPER approach across 53 large studies. Int J Epidemiol, 2011. **40**(5): p. 1314-28.
- 74. Pinart, M., et al., Comorbidity of eczema, rhinitis and asthma in IgE-sensitized and non-sensitized children: an international population-based cohort. Lancet Respir Med, 2014. in press.
- 75. Pinart, M. and e. al, Excess comorbidity of eczema, rhinitis and asthma in IgE- and non-IgE-sensitized children: an
- international population-based study. Lancet Resp Med, 2014. in press.
- 76. Hohmann, C., et al., The MeDALL core questionnaires for a harmonized follow-up assessment of 11 European birth cohorts on asthma and allergies. Int Arch Allergy Immunol, 2014. in press.
- 77. Kotecha, S.J., et al., Effect of late preterm birth on longitudinal lung spirometry in school age children and adolescents. Thorax, 2012. **67**(1): p. 54-61.
- 78. Kerkhof, M., et al., *Transient early wheeze and lung function in early childhood associated with chronic obstructive pulmonary disease genes.* J Allergy Clin Immunol, 2013.
- Litonjua, A.A., et al., Maternal antioxidant intake in pregnancy and wheezing illnesses in children at 2 y of age. Am J Clin Nutr, 2006. 84(4): p. 903-11.
- Sordillo, J.E., et al., Multiple microbial exposures in the home may protect against asthma or allergy in childhood. Clin Exp Allergy, 2010. 40(6): p. 902-10.
- 81. Donahue, S.M., et al., *Prenatal fatty acid status and child adiposity at age 3 y: results from a US pregnancy cohort.* Am J Clin Nutr, 2011. **93**(4): p. 780-8.
- Regnault, N., et al., Sex-specific associations of gestational glucose tolerance with childhood body composition. Diabetes Care, 2013. 36(10): p. 3045-53.
- Wijga, A.H., et al., Adenotonsillectomy and the development of overweight. Pediatrics, 2009.
   123(4): p. 1095-101.
- 84. Willers, S.M., et al., BMI development of normal weight and overweight children in the PIAMA study. PLoS One, 2012. **7**(6): p. e39517.
- de Rooij, S.R. and T.J. Roseboom, The developmental origins of ageing: study protocol for the Dutch famine birth cohort study on ageing. BMJ Open, 2013. 3(6).
- 86. Veenendaal, M.V., et al., *Transgenerational effects of prenatal exposure to the 1944-45 Dutch famine.*BJOG, 2013. **120**(5): p. 548-53.
- 87. de Lauzon-Guillain, B., et al., Birth weight, body silhouette over the life course, and incident diabetes in 91,453 middle-aged women from the French Etude Epidemiologique de Femmes de la Mutuelle Generale de l'Education Nationale (E3N) Cohort. Diabetes Care, 2010. 33(2): p. 298-303.
- 88. Hercberg, S., et al., A primary prevention trial using nutritional doses of antioxidant vitamins and

- minerals in cardiovascular diseases and cancers in a general population: the SU.VI.MAX study-design, methods, and participant characteristics. SUpplementation en VItamines et Mineraux AntioXydants. Control Clin Trials, 1998. **19**(4): p. 336-51.
- 89. Kesse-Guyot, E., et al., *Mediterranean diet and cognitive function: a French study*. Am J Clin Nutr, 2013. **97**(2): p. 369-76.
- 90. Monsuez, J.J., et al., Impact of 6-year body weight change on cardiac geometry and function in ageing adults: the SUpplementation en Vitamines et Mineraux AntioXydants -2 (SU.VI.MAX-2) cardiovascular ultrasound substudy. J Hypertens, 2010. 28(11): p. 2309-15.
- 91. Clavel-Chapelon, F., et al., E3N, a French cohort study on cancer risk factors. E3N Group. Etude Epidemiologique aupres de femmes de l'Education Nationale. Eur J Cancer Prev, 1997. 6(5): p. 473-8.
- 92. Riboli, E., Nutrition and cancer: background and rationale of the European Prospective Investigation into Cancer and Nutrition (EPIC). Ann Oncol, 1992. **3**(10): p. 783-91.
- 93. Riboli, E., The role of metabolic carcinogenesis in cancer causation and prevention: evidence from the European prospective investigation into cancer and nutrition. Cancer Treat Res, 2014. **159**: p. 3-20.
- Fagherazzi, G., et al., Dietary acid load and risk of type 2 diabetes: the E3N-EPIC cohort study. Diabetologia, 2013.
- Engel, P., et al., Risk of osteoporotic fractures after discontinuation of menopausal hormone therapy: results from the E3N cohort. Am J Epidemiol, 2011. 174(1): p. 12-21.
- de Marco, R., et al., Risk Factors for Chronic Obstructive Pulmonary Disease in a European Cohort of Young Adults. Am J Respir Crit Care Med, 2010.
- 97. Norback, D., et al., Mould and dampness in dwelling places, and onset of asthma: the population-based cohort ECRHS. Occup Environ Med, 2013. **70**(5): p. 325-31.
- 98. Kauffmann, F., et al., Epidemiological study of the genetics and environment of asthma, bronchial hyperresponsiveness, and atopy: phenotype issues. Am J Respir Crit Care Med, 1997. **156**(4 Pt 2): p. \$123-9
- Bouzigon, E., et al., Effect of 17q21 variants and smoking exposure in early-onset asthma. N Engl J Med, 2008. 359(19): p. 1985-94.
- Boudier, A., et al., Ten-year follow-up of clusterbased asthma phenotypes in adults. A pooled analysis of three cohorts. Am J Respir Crit Care Med, 2013. 188(5): p. 550-60.
- Mohamed Hoesein, F.A., et al., Variants in the 15q24/25 locus associate with lung function decline in active smokers. PLoS One, 2013. 8(1): p. e53219.
- Stolk, R.P., et al., Universal risk factors for multifactorial diseases: LifeLines: a threegeneration population-based study. Eur J Epidemiol, 2008. 23(1): p. 67-74.
- 103. Ritchie, K., et al., Association of adverse childhood environment and 5-HTTLPR Genotype with late-

- life depression. J Clin Psychiatry, 2009. **70**(9): p. 1281-8
- 104. Portet, F., et al., Metabolic syndrome and localization of white matter hyperintensities in the elderly population. Alzheimers Dement, 2012. 8(5 Suppl): p. S88-95 e1.
- 105. Heijmans, B.T., et al., Persistent epigenetic differences associated with prenatal exposure to famine in humans. Proc Natl Acad Sci U S A, 2008. 105(44): p. 17046-9.
- 106. Giovanoli, S., et al., Stress in puberty unmasks latent neuropathological consequences of prenatal immune activation in mice. Science, 2013.
   339(6123): p. 1095-9.
- 107. Akbaraly, T.N., et al., Leisure activities and the risk of dementia in the elderly: results from the Three-City Study. Neurology, 2009. **73**(11): p. 854-61.
- 108. Berr, C., et al., *Olive oil and cognition: results from the three-city study*. Dement Geriatr Cogn Disord, 2009. **28**(4): p. 357-64.
- Ritchie, K., et al., Designing prevention programmes to reduce incidence of dementia: prospective cohort study of modifiable risk factors. BMJ, 2010. 341: p. c3885.
- Wheelock, C.E., et al., Application of 'omics technologies to biomarker discovery in inflammatory lung diseases. Eur Respir J, 2013.
   42(3): p. 802-25.
- 111. Poste, G., *Bring on the biomarkers*. Nature, 2011. **469**(7329): p. 156-7.
- 112. Taylor, D.R. and I.D. Pavord, Biomarkers in the assessment and management of airways diseases. Postgrad Med J, 2008. 84(998): p. 628-34; quiz 633
- Manolio, T.A., Genomewide association studies and assessment of the risk of disease. N Engl J Med, 2010. 363(2): p. 166-76.
- 114. Moffatt, M.F., et al., A large-scale, consortiumbased genomewide association study of asthma. N Engl J Med, 2010. 363(13): p. 1211-21.
- 115. Roberts, T.C., K.V. Morris, and M.S. Weinberg, Perspectives on the mechanism of transcriptional regulation by long non-coding RNAs. Epigenetics, 2013. **9**(1).
- Trynka, G., et al., Chromatin marks identify critical cell types for fine mapping complex trait variants. Nat Genet, 2013. 45(2): p. 124-30.
- 117. Jin, F., et al., A high-resolution map of the threedimensional chromatin interactome in human cells. Nature, 2013. **503**(7475): p. 290-4.
- 118. Melen, E. and G. Pershagen, *Pathophysiology of asthma: lessons from genetic research with particular focus on severe asthma.* J Intern Med, 2012. **272**(2): p. 108-20.
- 119. Ponomarenko, E., et al., *The Chromosome-Centric Human Proteome Project at FEBS Congress*. Proteomics, 2013.
- 120. Portelli, M.A., et al., Genome-wide protein QTL mapping identifies human plasma kallikrein as a post-translational regulator of serum uPAR levels. FASEB J, 2013.