

## Choking on a foreign body: a physiological study of the effectiveness of abdominal thrust manoeuvres to increase thoracic pressure

**ABSTRACT** The Heimlich manoeuvre is a well-known intervention for the management of choking due to foreign body airway occlusion, but the evidence base for guidance on this topic is limited and guidelines differ. We measured pressures during abdominal thrusts in healthy volunteers. The angle at which thrusts were performed (upthrust vs circumferential) did not affect intrathoracic pressure. Self-administered abdominal thrusts produced similar pressures to those performed by another person. Chair thrusts, where the subject pushed their upper abdomen against a chair back, produced higher pressures than other manoeuvres. Both approaches should be included in basic life support teaching.

### BACKGROUND

Foreign body airway obstruction (FBAO) is a common cause of death, particularly in older people. The National Safety Council USA reports that FBAO is the fourth leading cause of unintentional injury death, with 4864 reported deaths in 2013.<sup>1</sup> The 'Heimlich' manoeuvre is a technique for expelling an obstructing food bolus where a first-aider places their arms round the subject from behind and delivers a sharp inward and upward thrust to the abdomen below the rib cage. Heimlich described 162 cases where life was saved following successful administration of abdominal thrusts.<sup>2</sup>

European Resuscitation Council<sup>3</sup> guidance for treatment of FBAO in conscious adults is a combination of back blows and abdominal thrusts with no preference on order. The Australian and New Zealand Resuscitation Councils recommend back blows and chest thrusts for the management of FBAO in conscious adults, but advise against abdominal thrusts, citing concern about complications.<sup>4</sup>

External pressure on the abdomen should be transmitted through the diaphragm regardless of where it is applied, so there is no theoretical reason why force needs to be directed upwards. Motivated in part by three cases of near death from choking involving UK chest physicians (see online supplement), we describe experiments to address these two questions.

### METHODS

Detailed methods are available in the online supplement. Briefly, different expulsive

manoeuvres (see [box 1](#)) were performed on and by four consenting adult physiology researchers median (range): age 56.5 (46, 74) years and body mass index (BMI) 25.9

(25, 26) kg/m<sup>2</sup>. Oesophageal and gastric balloon catheters were placed to record pressures generated. Detailed statistical analysis is available in the online supplement.

### Box 1 Description of manoeuvres

#### *Circumferential 'horizontal' abdominal thrust*

The operator stands behind the participant, grasps their fists together and places thumb side of the fist over the fleshy part of the abdomen above the navel. The operator pulls sharply backwards starting with medium force and progressively increasing force, until the maximum pressure that the subject feels is acceptable is achieved.

#### *Heimlich manoeuvre*

The same procedure but with an upward direction of force.

#### *Auto 'upthrust' abdominal thrust*

The participant positions their own hands in the standard position for the abdominal manoeuvre and performs thrusts increasing to the maximal force they can tolerate.

#### *Chair thrust*

The participant positions themselves above a high backed chair, with the chair back positioned below the upper half of the abdomen, below the ribcage. Using gravity, bodyweight and arms for additional force, the participant allows the back of the chair to thrust up into their abdomen (see [figure 1](#)).


#### *Volitional maximal cough and sniff pressures*

The participant performed repeated maximal volitional cough and sniff manoeuvres. All manoeuvres were performed after exhalation to the end of a normal breath (at functional residual capacity) with mouth and glottis closed and a nose clip in situ.



**Figure 1** One of the authors (MH) performing a chair thrust on himself (see also online supplementary video).

## Research letter



**Figure 2** Oesophageal and gastric pressure responses to expulsive manoeuvres. All statistical tests Mann-Whitney test. \*Results are statistically significant ( $p < 0.05$ ). (A) Maximal oesophageal pressures ( $P_{oes}$ ) achieved by expulsive manoeuvres. Pressure was significantly higher for chair thrust ( $p = 0.008$ ) but did not differ between conventional upthrust Heimlich, circumferential abdominal thrust or self-administered autoabdominal thrust. (B) Maximal gastric pressures were significantly higher in chair thrust compared with Heimlich manoeuvres ( $p = 0.004$ ). The outlier autoabdominal thrust data point in (B) corresponds to the participant (MH) who had performed an abdominal thrust on himself described in case 2 (see online supplement).

## RESULTS

Maximum peak oesophageal ( $P_{oes}$ ) and gastric ( $P_{gas}$ ) pressures were similar for the different abdominal thrusts when performed by the experimenters or by the subjects on themselves (see figure 2 and online supplementary table E1). For the upthrust Heimlich manoeuvre,  $P_{oes}$  was  $57 \pm 17$  cm H<sub>2</sub>O and for the circumferential abdominal thrust  $53 \pm 11$  cm H<sub>2</sub>O ( $p = 0.7$ ). The chair thrust generated a significantly higher  $P_{oes}$  than both;  $115 \pm 27$  cm H<sub>2</sub>O ( $p = 0.008$  compared with Heimlich).

In one participant, three further manoeuvres were performed. The  $P_{oes}$  generated by back slaps (7 cm H<sub>2</sub>O) and chest compressions when supine (position taken as for CPR) (42 cm H<sub>2</sub>O) were lower than both the Heimlich manoeuvre (64 cm H<sub>2</sub>O) and cough (179 cm H<sub>2</sub>O) in that subject.  $P_{oes}$  from supine abdominal compressions was 86 cm H<sub>2</sub>O, comparable to abdominal thrusts when upright.

## DISCUSSION

Abdominal thrusts caused a sharp rise in abdominal and thoracic pressures, exceeding or equal to those produced by these and alternative manoeuvres in previous studies.<sup>5–8</sup>

In 12 supine cadavers, mean airway pressure was 40.8 cm H<sub>2</sub>O for chest and 26.4 cm H<sub>2</sub>O for abdominal thrusts.<sup>5</sup> In six anaesthetised and intubated healthy adult male volunteers airway pressures from abdominal thrusts, low-chest thrusts and mid-chest thrusts in the horizontal lateral and sitting positions were compared.<sup>6</sup> Low-chest thrusts in the horizontal lateral position (34.0 cm H<sub>2</sub>O) and

mid-chest thrusts in the sitting position (46.2 cm H<sub>2</sub>O) produced the highest pressures. In eight intubated and anaesthetised pigs, anterior chest thrusts and Heimlich manoeuvres, both performed in a seated position, produced airway pressures of 6.5 and 13.8 cm H<sub>2</sub>O, respectively, with lateral chest thrusts in the side-lying position producing 18.0 cm H<sub>2</sub>O.<sup>7</sup> Day *et al*<sup>8</sup> compared alveolar pressure change for back blows (17.7 cm H<sub>2</sub>O) and the Heimlich manoeuvre (36.7 cm H<sub>2</sub>O).

An upward thrust may be more likely to cause injury to the ribcage or other organs and the person performing it may be inhibited by this possibility. Given the similar pressures generated by circumferential abdominal thrusts, we recommend that the manoeuvre should be performed by inward thrust over the fleshy part of the abdomen, around the level of the navel. This is important information for first aid providers with mismatched height to victim; a smaller individual can perform the circumferential abdominal thrust and produce the same intrathoracic pressure as the Heimlich manoeuvre that requires upthrust.

Australian and New Zealand ALS guidance does not recommend abdominal thrusts.<sup>4</sup> Studies on pigs, cadavers or anaesthetised subjects are unlikely to be representative of the situation and lung volumes in an upright, conscious choking individual. In our study, manoeuvres were all performed with the subjects at functional residual capacity, which may be more representative of the situation in an emergency. Given our data, we suggest that these guidelines should be amended. Concern about possible risk needs to be

balanced against the almost certain risk of death if obstruction persists and the further reduction in risk if a circumferential approach is used.

Self-administered abdominal thrusts were as effective as operator-delivered thrusts and indeed they had been used successfully in two of our cases (online). Repeated manoeuvres can be performed quickly and effectively without relying on an external operator. People choking may be encouraged to try this before a rescuer makes an attempt. Self-administering the manoeuvre is also a clear signal to rescuers (compared with clutching one's throat, which might be misinterpreted as distress due to another cause such as a heart attack). A novel finding is that self-administered thrusts over the back of a chair generated greater pressures than operator-delivered thrusts or self-administered ones. Most food is consumed seated, so there is likely to be a chair available when choking occurs.

## Obesity

No obese subjects were included in this study, median BMI 25.9 (25–26) kg/m.<sup>2</sup> Obesity may affect abdominal thrust outcome as anatomical landmarks for hand positioning may be variable and add to the difficulty of performing the manoeuvre around an increased abdominal circumference. A higher percentage of abdominal adipose tissue may have a dissipating effect on the force applied with abdominal thrust manoeuvres and therefore lessen their effectiveness.

## CONCLUSION

The key to diagnosing complete airway obstruction is a conscious subject, in the process of eating, who is unable to breathe at all, nor to speak. Autoadministered thrusts appear as physiologically effective as first-aider-administered ones to generate expulsive intrathoracic pressures, and chair thrusts appear to be the most physiologically effective. We advise that everyone with complete airway obstruction should, in the first instance, either autoadminister abdominal thrusts or perform a chair thrust. The various manoeuvres should be more widely taught in schools, first aid courses, to staff in restaurants and publicised as widely as possible. We would like to see suitable notices in eating places.

Matthew J Pavitt, Laura L Swanton, Matthew Hind, Michael Apps, Michael I Polkey, Malcolm Green, Nicholas S Hopkinson

NIHR Respiratory Biomedical Research Unit, Royal Brompton Hospital and Harefield NHS Foundation Trust and Imperial College, London, UK

**Correspondence to** Dr NS Hopkinson, Royal Brompton Hospital, Fulham Rd, London SW3 6NP, UK; [n.hopkinson@ic.ac.uk](mailto:n.hopkinson@ic.ac.uk)

**Twitter** Follow Matthew Pavitt @DrMattPav; Nicholas Hopkinson @COPDdoc

**Acknowledgements** The authors thank the individual involved in case 1 for permission to include a description account of his choking and John Moore-Gillon for permission to include his own account of his experience (case 3 online).

**Contributors** NSH, MG, MH and MIP conceived the study. MG, MP, LS, MA and NSH performed experiments and collected data. MP and LS analysed the results and produced the first draft of the paper to which all authors contributed, and all authors have approved the final version.

**Funding** The study was supported by the NIHR Respiratory BRU at Royal Brompton and Harefield NHS Foundation Trust and Imperial College, London, who part fund MIP's salary.

**Disclaimer** The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health. NSH affirms that the manuscript is an honest, accurate and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

**Competing interests** None declared.

**Ethics approval** Imperial College Joint Research Office and Health Research Authority.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** Requests for anonymised individual patient data can be made to the corresponding author.

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2016-209540>).



## OPEN ACCESS

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**To cite** Pavitt MJ, Swanton LL, Hind M, *et al.* *Thorax* Published Online First: [please include Day Month Year] doi:10.1136/thoraxjnl-2016-209540

Received 6 October 2016

Revised 27 October 2016

Accepted 7 November 2016

*Thorax* 2016;0:1–3.

doi:10.1136/thoraxjnl-2016-209540

## REFERENCES

- 1 National Safety Council. Choking prevention and rescue tips. 2016. <http://www.nsc.org/learn/safety-knowledge/Pages/safety-at-home-choking.aspx> (accessed 6 Oct 2016).
- 2 Heimlich HJ, Hoffmann KA, Canestri FR. Food-choking and drowning deaths prevented by external subdiaphragmatic compression. *Physiological basis. Ann Thorac Surg* 1975;20:188–95.
- 3 Perkins GD, Handley AJ, Koster RW, *et al.* European Resuscitation Council Guidelines for Resuscitation 2015: Section 2. Adult basic life support and automated external defibrillation. *Resuscitation* 2015;95:81–99.
- 4 Australian and New Zealand Resuscitation Councils. ANZCOR Guideline 4—Airway. 2016. <https://resus.org.au/guidelines/> (accessed 6 Oct 2016).
- 5 Langhelle A, Sunde K, Wik L, *et al.* Airway pressure with chest compressions versus Heimlich manoeuvre in recently dead adults with complete airway obstruction. *Resuscitation* 2000;44:105–8.
- 6 Guildner CW, Williams D, Subitch T. Airway obstructed by foreign material: the Heimlich maneuver. *JACEP* 1976;5:675–7.
- 7 Lippmann J, Taylor DM, Slocombe R, *et al.* Lateral versus anterior thoracic thrusts in the generation of airway pressure in anaesthetised pigs. *Resuscitation* 2013;84:515–9.
- 8 Day RL, Crelin ES, DuBois AB. Choking: the Heimlich abdominal thrust vs back blows: an approach to measurement of inertial and aerodynamic forces. *Pediatrics* 1982;70:113–19.

THORAX

## Choking on a foreign body: a physiological study of the effectiveness of abdominal thrust manoeuvres to increase thoracic pressure

Matthew J Pavitt, Laura L Swanton, Matthew Hind, Michael Apps, Michael I Polkey, Malcolm Green and Nicholas S Hopkinson

*Thorax* published online April 12, 2017

---

Updated information and services can be found at:

<http://thorax.bmj.com/content/early/2017/04/06/thoraxjnl-2016-209540>

---

*These include:*

### References

This article cites 6 articles, 1 of which you can access for free at: <http://thorax.bmj.com/content/early/2017/04/06/thoraxjnl-2016-209540#BIBL>

### Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (250)

---

### Notes

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>