Repeatability of respiratory data in the individual subject

LINDSAY DAVIDSON

From the Department of Medicine, Queen Elizabeth Hospital, Birmingham¹

A knowledge of the repeatability of any respiratory measurement in the individual is a prerequisite when assessing the significance of the change in state produced by any experimental procedure. Such information is available in the literature only in scattered and incomplete detail.

The accompanying Table presents such data, and, as the high degree of repeatability in the individual is often not appreciated, a record of the published variation for normal groups is included for comparison.

All measurements are based on duplicate sampling compared with duplicate sampling 10 minutes later. Details of the methods are available together with a complete statistical analysis in the author's M.D. thesis (1962) of the University of Birmingham. The patients were undergoing thoracic, abdominal, and other major surgical procedures. Measurements were made pre-operatively and 24 hours post-operatively, and there was no difference between the statistical analysis

of differences pre-operatively and post-operatively.

It is hoped that the publication of these data will help investigators to define a steady state in their patients.

REFERENCES

- Bartels, H., and Rodewald, G. (1952). Der arterielle Sauerstoffdruck, die alveolär-arterielle Sauerstoffdruckdifferenz und weitere atmungsphysiologische Daten gesunder Männer. *Pflügers. Arch. ges. Physiol.*, **256**, 113.
- Greifenstein, F. E., King, R. M., Latch, S. S., and Conroe, J. H., jun. (1952). Pulmonary function studies in healthy men and women 50 years and older. *J. appl. Physiol.*, **4**, 641.
- Handbook of Respiration* (1958). National Academy of Sciences, National Research Council. Edited by D. S. Dittmer and R. M. Grebe. Saunders, Philadelphia.
- Kory, R. C., Callahan, R., Boren, H. G., and Syner, J. C. (1961). The Veterans Administration-Army Co-operative Study of Pulmonary Function I. Clinical spirometry in normal men. *Amer. J. Med.*, **30**, 243.
- Raine, J. M., and Bishop, J. M. (1963). A—a difference in O₂ tension and physiological dead space in normal man. *J. appl. Physiol.*, **18**, 284.
- Riley, R. L., Campbell, E. J. M., and Shepard, R. H. (1957). A bubble method for estimation of Pco₂ and Po₂ in whole blood. *J. appl. Physiol.*, **11**, 245.
- , Cournand, A., and Donald, K. W. (1951). Analysis of factors affecting partial pressures of oxygen and carbon dioxide in gas and blood of lungs: methods. *Ibid.*, **4**, 102.
- Shock, N. W., and Yengst, M. J. (1955). Age changes in basal respiratory measurements and metabolism in males. *J. Geront.*, **10**, 31.

TABLE

Measurement	No. of Paired Comparisons	S.D. of Difference	95% Confidence Limit of Difference	Published S.D. within a Group of Normals	Reference
Vital capacity (l.)	56	0.16	0.26	0.48	Greifenstein <i>et al.</i> (1952)
One-second forced expiratory volume (l./sec.)	56	0.11	0.25	0.67	Kory <i>et al.</i> (1961)
Maximal mid-expiratory flow rate (l./sec.)	56	0.50	1.03	1.30	Kory <i>et al.</i> (1961)
Respiratory frequency (breaths/min.)	41	1.40	2.83	4.80	Greifenstein <i>et al.</i> (1952)
Minute ventilation (l./min.)	41	0.51	1.03	1.70	Greifenstein <i>et al.</i> (1952)
Tidal volume (ml.)	41	28.00	57.00	116.00	Shock and Yengst (1955)
Oxygen consumption (ml./min.)	20	5.60	11.80	17.00	<i>Handbook of Respiration</i> (1958)
Carbon dioxide output (ml./min.)	20	6.40	13.50	11.07	<i>Handbook of Respiration</i> (1958)
Respiratory exchange ratio	20	0.04	0.084	0.140	Raine and Bishop (1963)
Bohr deadspace (ml.) ¹	20	12.60	26.60	51.50	Raine and Bishop (1963)
Deadspace tidal volume ratio (%)	20	2.00	4.20	6.20	Raine and Bishop (1963)
Alveolar oxygen tension (mm. Hg) ²	20	2.30	4.80	9.90	Raine and Bishop (1963)
Arterial oxygen tension (mm. Hg) ²	41	2.60	5.30	5.50	Bartels and Rodewald (1952)
Arterial carbon dioxide tension (mm. Hg) ³	41	1.70	3.40	1.90	<i>Handbook of Respiration</i> (1958)
Alveolo-arterial oxygen tension difference (mm. Hg)	20	2.60	5.50	7.40	Raine and Bishop (1963)
Estimated venous admixture (%)	20	2.40	5.00	—	—
Arterial carbon dioxide content (mm./l.)	35	0.17	0.35	2.10	Greifenstein <i>et al.</i> (1952)
Arterial blood pH	32	0.017	0.04	0.03	Greifenstein <i>et al.</i> (1952)

Notes: ¹ Assuming PACO₂ = PaCO₂.

² Calculated from alveolar air equation.

³ Microtonometry (Riley, Campbell, and Shepard, 1957).

⁴ Charts of Riley, Cournand, and Donald (1951).