Surgical stabilization of traumatic flail chest

F. PARÍS¹, V. TARAZONA¹, E. BLASCO¹, A. CANTÓ,
M. CASILLAS¹, J. PASTOR¹, M. PARÍS², and
R. MONTERO²

*Thoracic Surgery Service, Department of Surgery¹ and Department of Anaesthesia-Reanimation²,
Centro Hospitalario 'La Fé', Valencia 9, Spain*

París, F., Tarazona, V., Blasco, E., Cantó, A., Casillas, M., Pastor, J., París, M., and Montero, R. (1975). Thorax, 30, 521–527. Surgical stabilization of traumatic flail chest. Since 1970 we have stabilized the ribs to correct paradoxical movement of the chest wall in chest injuries, using an original technique, in order to avoid as far as possible the need for long-term chest wall stabilization by intermittent positive pressure respiration (IPPR).

The technical details of surgical stabilization are described, and the different types of stainless steel struts are shown. Type I was originally used either as an intramedullary nail or as an external brace. Types II and III were designed for external fixation of the strut to the rib.

Treatment of 29 patients with severe flail chest, classified into four groups, is shown: group I was treated by IPPR, group II by IPPR plus surgical stabilization, group III by surgical stabilization only, and group IV by surgical stabilization after exploratory thoracotomy.

The clinical results are discussed. We conclude that surgical stabilization of the paradoxical movement of the chest wall can avoid the use of the respirator or at least reduce the interval of IPPR to a short period during the initial recovery from trauma. Using type III struts, we have obtained stabilization of the flail chest in all cases even in patients with severe anterior paradoxical movement. The patients' tolerance of surgical stainless steel struts was good.

Although single uncomplicated fractures of the ribs are usually not serious, the prognosis is not the same when rib fractures are multiple or bilateral, or a fractured sternum coexists. In these circumstances part of the chest wall may move paradoxically.

The treatment of severe chest wall injury with paradoxical respiration remains a matter of controversy. Twenty years ago, fixation of ribs was considered useless because encircling sutures or screws or the insertion of thin nails did not abolish paradoxical movement. For this reason Avery, Mörch, and Benson (1956) considered stabilization of the chest wall by intermittent positive pressure respiration (IPPR) the most appropriate treatment.

Recently, however, Dor *et al.* (1972), Couraud, Bruneteau, and Durandeau (1973), Eschapase and Gaillard (1973) and others have reintroduced various procedures for rib fixation. Joudet (1973)

used struts composed of a bar with two staples at both ends for embracing the rib on both sides of the fracture. Similar struts have been described by Sanchez Lloret (1974). Since 1970 we have fixed the ribs to correct paradoxical movement of the chest wall in chest injuries using an original technique. The early results have already been reported (París, 1972).

Rib fixation has been re-evaluated in order to avoid as far as possible the need for long-term chest wall stabilization by IPPR. The disadvantages and dangers of IPPR are: the method is uncomfortable and requires meticulous care; difficulty in communication and the long period of bed rest are irksome to the patient; it can increase air leaks from the lung; it does not always correct traumatic thoracoplasty or avoid the healing of flail chest in a bad position, producing a stove-in deformity; it adds the complications of tracheostomy such as infection, tracheal stenosis,

FIG. 1. Thoracic lesions in 223 patients with thoracic trauma.

ulcerative tracheo-oesophageal fistula or damage to the innominate artery.

CLINICAL MATERIAL AND METHOD

From February 1969 until November 1974 we have admitted 233 patients with chest injuries; 210 were closed injuries and 23 open. Over half were due to road traffic accidents. Figure 1 shows the types of injury and the treatment of the 233 patients is shown in Table I.

Patients with severe paradoxical movement of the chest wall have been treated by IPPR as well as by surgical stabilization. Table II shows the treatment of 29 patients with severe flail chest classified into four groups.

TABLE I

TREATMENT OF 233 PATIENTS WITH CHEST INJURIES

Treatment	No.
Symptomatic treatment and observation	64
IPPR	27
Closed intercostal drainage	153
Surgical stabilization of flail chest	18
Thoracotomy	19

Some patients had more than one treatment procedure, for example IPPR with closed drainage.

TABLE II

TREATMENT OF 29 PATIENTS WITH SEVERE FLAIL CHEST

Group	Treatment	No.
I	IPPR	11
II	IPPR + surgical stabilization of flail chest	10
III	Surgical stabilization only	4
IV	Thoracotomy + surgical stabilization	4

The types of stainless steel struts designed for us¹, using the alloy 316 L, are presented in Figure 2. Type I was originally used either as an intramedullary nail or as an external brace anchored to the ribs by wire sutures (Figure 3).

The existence in some cases of ribs with a medullary canal too small for nailing led us to develop struts of types II and III (Fig. 2) designed for external fixation of the strut to the rib. The addition of a number of holes allows the suture (wire) not only to encircle the strut and rib, but to pass through the strut and rib, avoiding displacement of the prosthesis. The struts are appropriately curved and cut with a cold chisel at the time of operation.

¹Frammer Instruments, Valencia, Spain

FIG. 2. Types of stainless steel struts used for rib fixation.

Figure 4 shows different techniques for anchoring type II and type III struts to the ribs. As suture material we have used wire or other non-absorbable material. Figures 5 and 6 show the chest films of two patients treated by this method.

The struts were placed in position without thoracotomy except in the case of patients in whom thoracotomy was needed because of associated visceral injuries. In cases of lateral flail chest the approach for surgical stabilization was a lateral or posterolateral incision. In cases of anterior or anterolateral flail chest, a vertical midline incision for the approach to the sternum and a lateral incision for the approach to the ribs was used (Figure 7).

RESULTS

Group I (Table II), with flail chest treated only by IPPR, in our series had a high mortality of 8/11 (73%). The cause of this mortality can be explained. All patients in this group had multiple injuries: five had brain injuries and four were in severe shock on admission, and it was for these reasons that surgical stabilization was not carried out.

Group II consisted of 10 patients treated by IPPR plus surgical stabilization of the flail portion of the chest wall. At the time of admission to hospital these patients were unfit for surgical treatment, and the paradoxical movement of the chest wall was treated by IPPR. When the patient's state had improved, surgical stabilization was carried out and the patient was removed from the respirator. All patients in this group had multiple injuries. Three had cerebral injuries, one

was in shock on admission, and one was 82 years old. The mortality rate was 4/10. Two patients died in coma from associated brain damage; the paradoxical movement of the chest wall was relieved by the stabilization operation. In another patient the operation was carried out too late and death was due to ulcerative oesophagotracheal fistula and sepsis from too prolonged IPPR. If surgical stabilization had been carried out earlier in this patient, the oesophagotracheal fistula might have been avoided. The fourth patient was operated upon too soon after trauma before cardio-respiratory stability had been achieved. He had an irreversible cardiac arrest at the time of induction of anaesthesia.

Group III consisted of four patients treated by surgical stabilization only, who on admission were in good condition for surgical treatment. None of these patients had multiple injuries, cerebral damage or shock. There was no mortality in this group.

Group IV consisted of four patients treated by surgical stabilization after an exploratory thoracotomy for visceral injuries. One patient died from associated abdominal lesions. The other patient was in severe shock after gunshot injuries to the spleen, colon, diaphragm, lung, and chest wall.

In Table III the extent of rib fractures in 18 patients operated on is shown and its relation to the type of flail chest. In Table IV the site of paradoxical movement and the treatment group are given, showing that in the group of patients

TABLE III
INJURIES IN 18 PATIENTS TREATED BY SURGICAL STABILIZATION

Fractures	Site of Paradoxical Movement of Chest		
	Lateral	Anterolateral	Bilateral anterior
Multiple ribs on one side	7	1	—
Ribs on both sides	1	1	1
Ribs on one side + sternum	—	2	1
Ribs on both sides + sternum	—	2	2
Total	8	6	4

TABLE IV
SITE OF PARADOXICAL MOVEMENT AND TREATMENT GROUP

Site of Paradoxical Movement	Thoracotomy + Surgical Stabilization	Surgical Stabilization only	IPPR + Surgical Stabilization
Bilateral anterior	—	—	4
Anterolateral	—	1	5
Lateral	4	3	1

FIG. 3. Fixation with non-perforated struts used as intramedullary nails or external brace: (a) posterior intramedullary nailing and anterior bracing; (b) bilateral intramedullary nailing by two struts crossed and anchored to the anterior floating fragment; (c) similar technique for a lateral fragment in unilateral fractures.

treated by IPPR with secondary surgical stabilization, the paradoxical movement of the chest wall was anterior. In the group 'surgical stabilization only' the lesion was commonly of the lateral type. In Table V the results of surgical stabilization are given according to the site of paradoxical movement and the type of strut employed. Struts of types I and II were adequate to abolish paradoxical movement of the chest wall in lateral flail chest, but with anterior flail chest the thicker type III strut was required. The results were imper-

fect when the thinner types I and II struts were used.

DISCUSSION

Fixation of fractured ribs is, in our view, a method of great value in the treatment of severe flail chest. Surgical stabilization of the paradoxical movement of the chest wall can avoid the use of the respirator or at least reduce the interval of IPPR to a short period during initial recovery.

FIG. 4. Fixation with perforated struts: (a, b, and c), different methods of anchoring type III struts to the rib cage, in cases of anterior paradoxical movement; (d) types of suture of strut to rib.

from trauma. Its principal aim is to simplify the post-traumatic period by obviating the need for long-term IPPR and avoiding the potential dangers of this procedure. Short periods of IPPR are generally not unsafe.

Surgical stabilization of flail chest must be accompanied by adequate physiotherapy to assist in

raising the sputum. The method is not possible if the patient is in coma from associated brain injury or other cause and unable to co-operate.

Using type III struts we have obtained stabilization of flail chest in all cases, even in patients with severe anterior paradoxical movement. The patients' tolerance of surgical stainless steel struts

FIG. 5. Radiographs of a patient treated surgically to correct lateral paradoxical movement and depression of the left chest wall: (a) before fixation; (b, c) after fixation. The patient survived and is in good condition two years after injury.

FIG. 6. Postoperative films of a patient treated for severe postraumatic paradoxical movement produced by fractures of right ribs with fracture of the sternum. PA view (a) in forced inspiration, and (b) forced expiration; (c) shows lateral view. The patient survived and is in good condition 18 months after injury.

FIG. 7. In anterior or anterolateral flail chest a vertical midline incision is used for the approach to the sternum and a lateral incision for the ribs. The struts pass from one incision to another through a subpectoral plane.

was good. After consolidation of the fractures the struts are removed.

In some circumstances the patient must be treated by long-term IPPR but we must emphasize that the risk of rib and sternal surgical stabilization are increased after long periods of IPPR treatment. Then the problem is not only one of stabilization of the thoracic cage; there is the increased risk of infection of the operation wound from the neighbouring tracheostome and added

dangers from established intrapulmonary infection. Early use of surgical fixation, when indicated and possible, may forestall the onset of infection.

TABLE V
RESULTS OF SURGICAL STABILIZATION ACCORDING
TO SITE OF PARADOXICAL MOVEMENT AND
TYPE OF STRUT USED

Site of Paradoxical Movement	Strut		
	Types I and II	Type III	None
Anterior (bilateral and anterolateral)	3 imperfect ¹	6 good	—
Lateral	3 good	4 good	1 good

¹Required additional traction.

REFERENCES

- Avery, E. E., Mörch, E. T., and Benson, D. W. (1956). Critically crushed chests. A new method of treatment with continuous mechanical hyperventilation to produce alkalotic apnea and internal pneumatic stabilization. *Journal of Thoracic Surgery*, **32**, 291.
- Couraud, L., Bruneteau, A., and Durandea A. (1973). Volets thoraciques. Indications thérapeutiques en fonction de leur siège et du contexte clinique. *Annales de Chirurgie Thoracique et Cardiovasculaire*, **12**, 15.
- Dor, V., Noirclerc, M., Chauvin, G., Mermet, B., Kreitman, P., Leonardelli, M., and Amorós, J. F. (1972). Les traumatismes graves du thorax. Place de l'osteosynthèse dans leur traitement. A propos de 100 cas. *Nouvelle Presse Médicale*, **1**, 519.
- Eschapse, H. and Gaillard, J. (1973). Volets thoraciques. Principes de traitement. *Annales de Chirurgie Thoracique et Cardiovasculaire*, **12**, 1.
- Joudet, R. (1973). Personal communication.
- París, F. (1972). Repercusion sobre el mediastino de los traumatismos torácicos. *Archivos de Bronconeumología*, **9**, 270.

—, Cantó, A., Casillas, M., Pastor, J., Blasco, E., and Tarazona, V. (1971). Técnica personal en el tratamiento quirúrgico del tórax flácido traumático. *Libro de Actas de la X Reunión de la Sociedad de Medicina y Cirugía de Levante. Castellón*.

Sanchez Lloret, J. (1974). Osteosíntesis costal medi-

ante placa extraperiosteica. Técnica original. *Revista Quirúrgica Española*, 1, 69.

Requests for reprints to Dr. F. París, Servicio de Cirugía Torácica, Centro Hospitalario 'La Fé', Avenida Alferez Provisional 21, Valencia 9, Spain.