Evidence that rifampicin can be used safely for non-tuberculous diseases

G ACOCELLA, W BRUMFITT, AND J M T HAMILTON-MILLER

From the Department of Medical Microbiology, Royal Free Hospital, London

ABSTRACT The incidence of primary resistance to rifampicin in *Mycobacterium tuberculosis* has been analysed in countries where rifampicin is restricted to use for treating tuberculosis and in countries where its use is not restricted. There is no evidence that rifampicin-resistant *M tuberculosis* strains are more common where the use of the drug is unrestricted. Resistance to rifampicin is less common than is resistance to streptomycin or to isoniazid. We can thus see no danger of producing resistant strains of *M tuberculosis* if rifampicin therapy is used for short periods for non-tuberculous infections. The problem of resistant mutants arising in the non-tuberculous species being treated is overcome by combining rifampicin with trimethoprim.

Rifampicin has a uniquely wide spectrum of antimicrobial activity, which includes Gram-positive and Gram-negative organisms as well as *Mycobacterium tuberculosis*.¹ Because of this, in a relatively early phase of the clinical use of this antibiotic for non-tuberculous indications, rifampicin was employed alone in infections of the chest² and urinary tract.^{3,4} However, the observation that in these circumstances resistant mutants were rapidly selected caused the use of monotherapy with rifampicin in non-tuberculous infections to fall into disrepute.

As a logical means of overcoming this problem, a combination of rifampicin and trimethoprim was envisaged, based on a "double blockade" type of approach. The basic assumptions were confirmed in a series of microbiological studies,^{5,6} which indicated both the existence of synergism between the two compounds and the prevention of the selection of resistant mutants. The situation concerning fears for the development of resistance in *M tuberculosis*⁷ as a result of rifampicin/trimethoprim being used was summarised in a provocative editorial.⁸

Since then, the pharmacokinetic properties of the combination^{9,10} and its clinical effectiveness¹¹ have been evaluated, and confirmed the encouraging preclinical evidence. A further and most important part of the development of the combination has been, and continues to be,

devoted to the evaluation of possible risks connected with the extra-tuberculous use of rifampicin.

We agree whole-heartedly with Simmons¹² who stated: "those who advocate that rifampicin should be used much more freely should surely be certain that the risk of development of widespread rifampicin-resistant *M tuberculosis* is not only small but virtually non-existent." We, therefore, decided to carry out a comparative study on primary resistance in *M tuberculosis* to rifampicin in countries where the use of the antibiotic is restricted to tuberculosis, and in those where it is freely available for the treatment of both tuberculous and non-tuberculous infections.¹³ Our results indicated that *M tuberculosis* had remained uniformly sensitive to rifampicin after several years of use in both groups of countries. The incidence of primary resistance to rifampicin was extremely low. It was lower (although not significantly) than that observed to ethambutol and PAS (which are used only for tuberculosis), and significantly lower than that observed to streptomycin and isoniazid.

In view of the importance of this information we have decided to update and to extend the survey. The new and old data are reported in the present communication.

Method and sources

Primary resistance was defined, detected, and diagnosed as by Canetti *et al.*¹⁴ Data were

Address for reprint requests: Professor W Brumfitt, Department of Medical Microbiology, Royal Free Hospital, Pond Street, London NW3 2QG.

collected from countries and centres shown in tables 1 and 2, not only from the literature but also by personal communication with a number of acknowledged experts: Dr J Grosset (Paris), Dr FJ Guerra-Sanz and Dr Ortega-Calderon (Madrid), Dr M Tavares da Lima Filho (Sao Paulo), Dr GA Jaramillo (Bogota), Dr A Giobbi (Milan), and Dr M Lucchesi (Rome). In France, USA, and West Germany rifampicin is restricted for use in tuberculosis, while in the other countries in tables 1 and 2 it is freely available for the treatment of any infection.

Results

As is clear from table 1, the incidence of primary resistance to rifampicin is low and has not changed with time—hence, for practical purposes, it is a small clinical problem. From the data in table 2 it can be observed that much the same applies for the other major antitubercular drugs. Primary resistance has altered very little with time, and is relatively low in absolute terms.

Discussion

As in the previous survey, the level of sensitivity is similar in countries like France, where rifampicin is used only for tuberculosis, and Colombia, where the ratio of non-tubercular to tubercular use of the antibiotic is of the order of 9:1.

In agreement with the previous observations, the very few cases of acquired resistance to

rifampicin were found to be associated with the improper use of the drug in tuberculous patients, including cases of monotherapy. The latter observation is important, since it is well known that cases of bacteriological failure observed during combined antituberculosis regimens can occur with strains which are still sensitive to rifampicin.³⁰

It can, therefore, be concluded that treatment of non-tuberculous infections with rifampicin for short periods does not carry a risk of selecting rifampicin-resistant strains of *M tuberculosis*. Thus, the proposed use of the combination rifampicin-trimethoprim for periods of one week in non-tuberculous indications should in no way jeopardise the antituberculous efficacy of rifampicin.

References

- 1 Binda G, Domenichini E, Gottardi A *et al*. Rifampicin: a general review. *Arzneim Forsch* 1971; **21**:1907-77.
- 2 Citron KM, May JR. Rifamycin antibiotics in chronic purulent bronchitis. *Lancet* 1969; **2**:982-3.
- 3 Atlas E, Turck M. Laboratory and clinical evaluation of rifampicin. *Am J Med Sci* 1968; **256**:247-54.
- 4 Murdoch J McC, Speirs CF, Wright N, Wallace ET. Rifampicin. *Lancet* 1969; **1**:1094.
- 5 Kerry DW, Hamilton-Miller JMT, Brumfitt W. Trimethoprim and rifampicin: in vitro activities separately and in combination. *J Antimicrob Chemother* 1975; **1**:417-27.

Table 1 Percentage of strains of *M tuberculosis* found primarily resistant to rifampicin in the years 1969-78 in numerous different geographical locations

Country	Centre (*)	Reference (**)	Percentage primary resistance of <i>M tuberculosis</i> to rifampicin in indicated years (no of strains tested in parentheses)									
			1969	1970	1971	1972	1973	1974	1975	1976	1977	1978
Italy	MI	15 (1978)			1.05(379)	0.87(689)	0.48(617)	0.62(484)	0.00(486)	0.83(479)	0.00(443)	0.00(429)
	CE	15			←0.00(433)→							
	VR	16					←0.58(173)→					
	RO	17 (1977)			0.90(119)	1.06(283)		0.40(248)	0.90(331)	0.91(328)	0.76(261)	
	SS	17 (1974)	←0.00(70)→									
	NA	18 (1976)	←0.0(726)→						0.4(464)			
	TO	19					0.58(171)					
Argentina	NAT	20	←0.00 (3110)→									
Brazil	VAR	21							0.00(1268)			
Colombia	NAT	†						0.0(653)				
Spain	VA	22 (1978)			0.0(1216)				0.35(564)	0.2(675)	0.2(938)	
	VA	23							1.40(136)			0.00(179)
	SA	23							0.00(110)			
France	VAR	24									0.00(1907)	
	LY	25			←0.2(483)→					0.0(369)		
Tunisia	SF	26				←0.28(713)→						
West Germany	NAT	27					0.1(1036)					
USA	MAIN	28								0.3(3146)		
USA	HW	29						0.0(313)				

(*) MI = Milan; GE = Genoa; VR = Verona; RO = Rome; SS = Sassari; NA = Naples; TO = Turin; NAT = National; VAR = Various
VA = Valencia; MA = Madrid; SA = Salamanca; LY = Lyon; SF = Sfax; MAIN = Mainland; HW = Hawaii.

(**) Year quoted in parentheses indicates year of last updating completed (generally unpublished).

† Personal Communication, Jaramillo 1979.

Table 2 Comparative data on percentage of primary resistance of *M. tuberculosis* to some of the major antituberculous drugs in different countries

Percentage primary resistance of <i>M. tuberculosis</i> at the indicated years and countries																	
Antitubercular agents	Italy		Argentina		Brazil		Spain		France	Tunisia	West Germany	Mainland USA	Hawaii	Bolivia, Colombia Mexico			
	1975-77	1969	1972	1969-74	1975	1978	1970	1978									
Streptomycin	6.9	6.7	5.7	6.1	10.5	1.68	6.6	10.4	9.16	7.8	5.8	6.31	2.9	2.3	5.1	1.0	2.6
Isoniazid	3.17	3.7	3.2	3.8	9.7	5.04	8.8	7.2	6.60	5.5	2.4	7.99	3.4	1.8	4.4	0.3	4.5
p-Aminosalicylic acid	0.84	0.2	0.2	0.2	2.2	0.0	0.0	0.17	0.53	0.5	—	1.14	1.9	0.7	1.3	0.3	1.6
Ethambutol	0.84	0.0	0.0	0.0	0.0	0.0	0.0	0.17	0.21	0.5	0.05	—	0.2	—	0.7	0.0	0.6
Kanamycin	0.31	0.0	0.07	—	1.3	—	—	—	—	—	—	—	—	—	0.1	—	—
Rifampicin	0.74	0.0	0.0	0.0	0.0	0.0	0.0	0.35	0.21	0.0	0.0	0.28	0.1	—	0.3	0.0	0.0

G Acocella, W Brumfitt, and J M T Hamilton-Miller

- 6 Arioli V, Berti M, Carniti G, Rossi E, Silvestri LG. Interaction between rifampicin and trimethoprim in vitro and in experimental infections. *J Antimicrob Chemother* 1977; 3:87-94.
- 7 Morrison-Smith J. Rifampicin in clinical use. *J Antimicrob Chemother* 1976; 2:353-4.
- 8 Editorial. Rifampicin: for tuberculosis only? *Lancet* 1976; 1:290-1.
- 9 Acocella G, Scotti R. Kinetic studies on the combination rifampicin-trimethoprim in man. I. Absorption and urinary excretion after administration to healthy volunteers of single doses of the two compounds alone and in combination over a period of one week. *J Antimicrob Chemother* 1976; 2:271-7.
- 10 Hamilton-Miller JMT, Brumfitt W. Trimethoprim and rifampicin: pharmacokinetic studies in man. *J Antimicrob Chemother* 1976; 2:181-8.
- 11 Brumfitt W, Cooper J, Hamilton-Miller JMT. Rifaprim (rifampicin and trimethoprim): a comparative trial with cephadrine in patients with recurrent urinary infections. *Proceedings of the 11th International Congress of Chemotherapy* 1980; in press
- 12 Simmons NA. Synergy and rifampicin. *J Antimicrob Chemother* 1977; 3:109-11.
- 13 Acocella G, Hamilton-Miller JMT, Brumfitt W. Can rifampicin use be safely extended? Evidence for non-emergence of resistant strains of *Mycobacterium tuberculosis*. *Lancet* 1977; 1: 740-2.
- 14 Canetti G, Fox W, Khomenko A *et al.* Advances in techniques of testing mycobacterial drug sensitivity, and the use of sensitivity tests in tuberculosis control programmes. *Bull WHO* 1969; 41:21-43.
- 15 Giobbi A, Bulgheroni P, Mantellini P, Miradoli E, Pandiani C. Osservazioni sulla batteriologia della tubercolosi dopo rifampicina. *Riv Inst Vaccin Milano* 1975; 25:323-31.
- 16 Longarato-Jacometta F, Zambelli L. Osservazioni sul comportamento della resistenza batterica c.d. primaria in malati di tubercolosi polmonare di primo accertamento. *Friuli Med* 1975; 30:681-90.
- 17 Lucchesi M, Zubiani M, Bajocco M, Ginesu F. Ulteriore contributo sulla resistenza primaria del *Mycobacterium tuberculosis* ai chemioantibiotici specifici. *Lotta contro tubercolosi Malattie Polmonari Sociali* 1975; 45:1-8.
- 18 Nitti V. La resistenza primaria del microbatterio tubercolare. Aspetti batteriologici ed epidemiologici. *Arch Monaldi* 1972; 27:454-74.
- 19 Briccarello L, Bardessono F, Daglio C, Ferrano U, Giorgis GE, Maschini MT. Il fenomeno della resistenza primaria del bacillo di Koch: casistica raccolta nell'Ospedale Pneumologico San Luigi di Torino dal 1972 al 1975. *G Batteriol Virol Immunol* 1976; 69:49-59.
- 20 Amadio GE. Evolucion de los indices de resistencia bacteriana en tuberculosis en la Republica Argentina. *Arch Monaldi* 1977; 32:12-17.

- 21 Tavares da Lima Filho M. A resistencia bacilar primaria dos drogas no Brasil en especial na cidade de Sao Paulo. Proceedings of the 5th International Symposium on Chemotherapy of Tuberculosis, Madrid, 1976.
- 22 Ortega-Calderon A, March Arbos J. Evolucion de la resistencia primaria del *Mycobacterium tuberculosis* en los ultimos 5 años (1970-1976). *Arch Monaldi* 1977; **32**:1-11.
- 23 Garcia JA, Saenz MC, Prieto J, Martin F. Incidence of drug resistance on the mycobacterial strains isolated from treated and untreated patients. In: Williams JD, Geddes AM, eds. *Proceedings of the 9th International Congress of Chemotherapy*. New York: Plenum Press, 1976: 3:81-6.
- 24 Grosset J. *Bacteriological basis of the chemotherapy of tuberculosis*. PAHO Regional Seminar on Tuberculosis: Chemotherapy. Washington DC: Pan American Health Organisation, 1979.
- 25 Viallier J. La recherche de la sensibilité aux antibiotiques du bacille tuberculeux: est-elle toujours nécessaire pour la conduite correcte du traitement? *Lyon Med* 1978; **240**:425-6.
- 26 Najah S, Tchekel I. Étude de la résistance du bacille tuberculeux dans la region de Sfax, durant les années 1973 a 1975. *Rev Franc Malad Respir* 1978; **6**:179-82.
- 27 Hussels H, Matthiessen W, Kind A, Göbel D. Frequency and epidemiology of primary drug resistance in tuberculosis in the Federal Republic of Germany, including Berlin (West). Partial result of a study of the WATL. *Scand J Respir Dis* 1978; supplement **102**:65-7.
- 28 Kopanoff DE, Kilburn JO, Glassroth JL, Snider DE, Farer LS, Good RC. A continuing survey of tuberculosis primary drug resistance in the United States: March 1975 to November 1977. *Am Rev Respir Dis* 1978; **118**:835-442.
- 29 Pien FD, Michael NL, Ow CL, Vennesland K, Brodsky ML. Primary antituberculous drug resistance in Hawaii, 1957 to 1977. *Am Rev Respir Dis* 1978; **118**:701-4.
- 30 Medical Research Council. Co-operative controlled trial of a standard regimen of streptomycin, PAS and isoniazid and three alternative regimens of chemotherapy in Britain. *Tubercle* 1973; **54**:99-129.