Dose response of ipratropium bromide assessed by two methods

CHRISTOPHER J ALLEN AND ALASTAIR H CAMPBELL

From the Division of Thoracic Medicine, Repatriation General Hospital, Heidelberg, Australia

ABSTRACT The dose-response relationships of ipratropium bromide were assessed by two different techniques in two groups of 10 male patients with partially reversible airways obstruction. In a randomised double-blind fashion on four days, 10 patients were given 40 μg , 80 μg , or 120 μg of ipratropium bromide or placebo from identical containers. Baseline FEV₁ and vital capacity were measured and the measurements repeated after 40 minutes, one, two, four, and six hours, and any symptoms were elicited. In the second study, each patient received cumulative doses of 40 μg , 80 μg , and 120 μg . Baseline FEV₁ was obtained and repeated 35 minutes after each dose. The peak increase of the FEV₁ was comparable in both studies. The FEV₁ was slightly greater after 120 μg than after 40 μg . Although this reached statistical significance only in the first study, it was concluded that the cumulative dose-response technique was suitable for determining the peak response. However, this technique was unsuitable for assessing duration of effect, which could be examined only in the first study. After 120 μg of ipratropium bromide, the FEV₁ was significantly greater than after 40 μg at each time interval and it was greater than 80 μg at six hours ($p < 0.05$). No significant side effects were noted in either study. When prolonged effective bronchodilation is sought, a dose of 120 μg of ipratropium bromide may be preferable to the recommended dose of 40 μg .

The anticholinergic bronchodilator ipratropium bromide is a quarternary ammonium derivative of atropine. Its quarternary structure prevents it crossing the blood-brain barrier and causing central effects. As an aerosol it is highly specific and virtually free of side effects even in doses well in excess of those recommended.^{1 2}

It has been reported that the recommended dose, 40 μg from a metered dose inhaler, achieves a peak bronchodilator effect.³ Nevertheless, higher doses may prolong the duration of bronchodilation^{4 5}.

To assess both the peak bronchodilation and the duration of the response, we have determined the dose-response relationships of ipratropium bromide with two different techniques. In the first study we administered the test doses on different days. In this way we were able to assess the peak response, duration of action, and unwanted side effects. The second study employed a cumulative dose-response technique.⁶ It is simpler and allows several doses of a drug to be assessed over a short

period of time, and the peak effect and unwanted effects can be determined. Its limitation is that it does not allow the duration of action to be assessed.

Methods

A group of 16 male patients gave informed consent to taking part in the investigation. All had asthma characterised by episodes of wheeze and shortness of breath requiring bronchodilators for relief of symptoms, and 12 were receiving beclomethasone dipropionate by inhalation or oral corticosteroids or both. Only six of the 16 patients had positive skin tests to common allergens and all but one had chronic productive cough. Eleven were current cigarette smokers.

SINGLE DOSE STUDIES

For the first study, 10 male patients were admitted to hospital in a stable condition. Bronchodilators were withheld for at least nine hours before any test. However, steroids were continued unchanged.

In a randomised, double-blind fashion on four

Address for reprint requests: Dr AH Campbell, Repatriation General Hospital, Heidelberg 3077, Australia.

days (consecutive where possible), the 10 patients studied were given 40 μg , 80 μg , or 120 μg of aerosol ipratropium bromide or placebo from identical containers. Two unlabelled containers were used, containing either placebo or active agent (20 μg per puff), so that the patient each day received six puffs of aerosol over 90 seconds.

Each study started at about 0900 hr. Baseline measurements of forced expiratory volume in one second (FEV_1) and vital capacity (VC) were performed, and the best of three readings taken. The patient then inhaled the aerosol. At 40 minutes, one, two, four, and six hours after the aerosol measurements of FEV_1 and VC were performed. Again the best of three measurements was taken, corrected to BTPS. Symptoms were recorded before each measurement. Baseline measurements each day were within 15% of each other. Tests were postponed when the baseline values were outside this range. The significance of the results obtained was analysed using analysis of variance.

CUMULATIVE DOSE-RESPONSE STUDY

For the cumulative dose-response study, 10 male outpatients were assessed (four common to the first study). Bronchodilators were withheld as before.

The peak effect of ipratropium bromide is not reached until about 30–40 minutes.³ This peak is then maintained for two to three hours. For this reason, after baseline measurements of FEV_1 and VC, and then the inhalation of 40 μg of ipratropium bromide, 35 minutes were allowed to elapse before three measurements of FEV_1 were made. The patient then inhaled another 40 μg and the measurements were repeated beginning 35 minutes after this dose. A third dose of 40 μg was then inhaled and measurements repeated 35 minutes later.

In this way, the patient received cumulative doses of 40 μg , 80 μg , and 120 μg . The best of three FEV_1 readings were taken each time. The readings were uncorrected because of the brief duration of the investigation. Symptoms were elicited after each dose. The significance of the results were analysed using analysis of variance.

Results

SINGLE DOSE STUDIES

The baseline mean FEV_1 values before placebo and to each dose of ipratropium bromide did not differ significantly. The mean and range in litres before placebo, 40 μg , 80 μg , and 120 μg of ipratropium bromide were 1.08 (0.40–1.80), 1.06 (0.40–1.70), 1.09 (0.40–1.90).

Compared with the placebo, each dose of ipratropium bromide produced a significant increase of the mean FEV_1 ($p < 0.05$). This was found up to two hours for the 40 μg dose and up to six hours with the 80 μg and 120 μg doses. Given the baseline value, after 80 μg the FEV_1 was significantly greater than after 40 μg at four hours ($p < 0.05$) whereas for 120 μg the FEV_1 was significantly greater than 40 μg at all times except zero and was greater than 80 μg at six hours ($p < 0.05$) (fig 1). Apart from a dry cough in one patient after 120 μg there were no side effects.

CUMULATIVE DOSE-RESPONSE STUDY

The results for the cumulative dose response study are shown in fig 2. The mean baseline FEV_1 was 1.06 l (range 0.38–2.03 l). After 40 μg there was a significant increase in mean FEV_1 of 0.30

Fig 1 Mean increase $\pm$ SEM of FEV_1 after different doses of ipratropium bromide, $n=10$. $\blacktriangle$ = 120 mcg, --- = 80 mcg, $\cdots$ = 40 mcg $\blacksquare$ = placebo.

Fig 2 Mean increase $\pm$ SEM of FEV_1 after cumulative doses of ipratropium bromide, $n=10$.

(28%) compared to the mean baseline FEV₁. After a cumulative dose of 80 µg there was a further rise of 6%, and after a cumulative dose of 120 µg, a further rise of 3% in FEV₁, though neither of these latter changes was statistically significant.

No side effects were recorded.

Discussion

Using two different methods we found a small increase in the peak bronchodilatation when doses of 80 µg and 120 µg of ipratropium bromide were compared with 40 µg. The differences were not statistically significant in the cumulative dose study, but reached significance at the 5% level with the dose of 40 µg versus 120 µg administered on separate days. Ruffin and Newhouse⁵ found that 60 µg of ipratropium bromide gave a significantly greater peak maximum mid-expiratory flow rate than 20 µg and 40 µg, but no difference for peak FEV₁. Most other studies,^{4,7} have failed to show that doses of ipratropium bromide above 40 µg produce a greater response than that achieved by 40 µg. The improvement in the peak response in the present study was small. It can be concluded that a dose of 40 µg of ipratropium bromide produces close to the peak response obtainable. The investigation has shown that the cumulative dose-response technique is suitable for determining the peak response. However, the technique does not allow assessment of duration of effect. For this, the more troublesome procedure of administering doses on different days is necessary. Ruffin and Newhouse⁵ have shown previously that 60 µg of ipratropium bromide gives a significantly longer duration of action than 20 µg or 40 µg, whereas Baigelman and Chodosh⁴ found in a group of chronic bronchitics that 80 µg gave a significantly longer lasting effect than 40 µg. In the present investigation it was found that 120 µg produced significant bronchodilatation for at least six hours. It was superior to 40 µg and marginally better than 80 µg. The patients investigated may have been particularly responsive to anticholinergic drugs. They all had asthma but were investigated when relatively stable and most were receiving steroids. All but one had chronic productive cough consistent with chronic bronchitis.

Only six had positive skin reactions to common allergens. Patients with these characteristics may be more responsive to anticholinergic drugs than other asthmatics.^{8,9}

For the type of patient studied, our results show that 120 µg of ipratropium bromide offers a higher peak and longer duration of bronchodilatation than the usually recommended dose of 40 µg. The dose of 120 µg was well tolerated, and provided this is confirmed, this dose of ipratropium bromide may prove useful to obtain prolonged effective bronchodilation in clinical practice.

References

- 1 Wood MJ, Paterson JW. Evaluation of cumulative doses of Sch 1000 in asthmatics. *Postgrad Med J* 1975; **51**, Supplement 7:100.
- 2 Simonsson BG, Jonson B, Ström B. Bronchodilatory and circulatory effects of inhaling increasing doses of an anti-cholinergic drug, ipratropium bromide (Sch 1000). *Scand J Respir Dis* 1975; **56**:138-149.
- 3 Loddenkemper R. Dose and time response of Sch 1000 MDI on total (Rt) and expiratory (Re) airways resistance in patients with chronic bronchitis and emphysema. *Postgrad Med J* 1975; **51**, Supplement 7:97.
- 4 Baigelman W, Chodosh S. Bronchodilator action of the anticholinergic drug ipratropium bromide (Sch 1000) as an aerosol in chronic bronchitis and asthma. *Chest* 1977; **71**:324-328.
- 5 Ruffin RE, Newhouse MT. Ipratropium bromide (Sch 1000) monohydrate aerosol: bronchodilator effect of three dose levels in asthmatics. *Lung* 1978; **155**:141-46.
- 6 Shenfield GM, Paterson JW. Clinical assessment of bronchodilator drugs delivered by aerosol. *Thorax* 1973; **28**:124-8.
- 7 Gross HJ. Dose response study of the effect of Sch 1000 MDI as forced vital capacity (FVC), FEV_{1.0}, maximal mid-expiratory flow (MMEF) and heart rate, E.C.G. and blood pressure. *Postgrad Med J* 1975; **51**, Supplement 7:95.
- 8 Altounyan REC. Variation of drug action on airway obstruction in man. *Thorax* 1964; **19**:406-15.
- 9 Ruffin RE, Fitzgerald JD, Rebuck AS. A comparison of the bronchodilator activity of Sch 1000 and salbutamol. *J Allergy Clin Immunol* 1977; **59**:136-41.