Pulmonary alveolar lipoproteinosis complicating juvenile dermatomyositis

M P SAMUELS, J O WARNER

From Brompton Hospital, London

ABSTRACT A 9 year old boy with juvenile dermatomyositis developed pulmonary alveolar lipoproteinosis. This previously unreported association was treated successfully by bronchoalveolar lavage.

Pulmonary alveolar lipoproteinosis is a rare condition of unknown aetiology characterised by accumulation of a lipoproteinaceous material within alveolar spaces.¹ The diagnosis is confirmed by finding periodic acid-Schiff (PAS) staining material on light microscopy and characteristic lamellar bodies on electron microscopy. Although it has previously been reported to be associated with several immunological abnormalities in children and with haematological malignancies and respiratory infections in adults, an association with autoimmune disease has not been noted. We report a case of the disease associated with juvenile dermatomyositis.

Case report

Dermatomyositis was diagnosed in a 7 year old boy. He had a six month history of decreasing exercise tolerance and muscle stiffness and eventually developed a typical rash. At diagnosis his creatinine kinase was 534 IU/l and the antinuclear factor test gave a weakly positive result. He was treated with prednisolone, at a starting dose of 2 mg/kg daily. Progress was complicated by interfascial calcinosis and a fracture of his left humerus secondary to osteoporosis, but within two years he had nearly full muscle strength and the prednisolone had been reduced to less than 1 mg/kg on alternate days.

Two and a half years after the onset he developed a respiratory infection, for which he was treated with erythromycin; but he remained short of breath, had a persistent cough, and lost weight. A chest radiograph showed patchy infiltrates throughout both lung fields, but no hilar adenopathy, pleural effusion, or cardiomegaly. In view of the possibility of *Pneumocystis carinii* infection he received high dose co-trimoxazole. As his symptoms and radiographic findings did not improve he was referred for further investigation.

On admission his height and weight were on the 50th percentile. He had a heliotrope rash on his upper eyelids, generalised muscle wasting, and early clubbing. He was mildly cyanosed with a respiratory rate of 30/min, poor chest expansion, and bilateral basal inspiratory crackles on auscultation.

Lung function tests showed reduced lung volumes with a severe mixed obstructive and restrictive defect and no response to a bronchodilator; transfer coefficient (Kco) was normal at 1.92-2.35 mmol min⁻¹ kPa⁻¹ l⁻¹.

His haemoglobin was 17.5 g/l and his arterial oxygen tension (Pao₂) 5.5 kPa. Bronchial washings revealed no pathogens; they were not examined for PAS staining bodies. Open lung biopsy showed features typical of alveolar lipoproteinosis with no evidence of pulmonary fibrosis. Further investigations indicated normal white cell count, complement, immunoglobulins, opsonins, and bone marrow. Lymphocyte subsets were normal, but there was a serum dependent defect of neutrophil candida phagocytosis and killing.

Bronchoalveolar lavage was performed in view of poor exercise tolerance, hypoxaemia, and an alveolar-arterial Po₂ difference of 8 kPa. Under general anaesthesia with a double lumen endotracheal tube the right lung was lavaged with 500 ml aliquots of normal saline, warmed to body temperature, until clear fluid returned. A total of 20 l was used. The left lung was lavaged a week later. Both procedures were tolerated without complications.

There was an immediate symptomatic and physiological improvement and 12 months later exercise tolerance, lung function, and Pao₂ were normal (table). The radiographic infiltrates have shown gradual clearing (figure). Muscle strength has remained normal as prednisolone has been slowly reduced.

Discussion

Pulmonary alveolar proteinosis is associated with immunological abnormalities, including hypogammaglobinaemia, lymphopenia, and thymic hypoplasia.² Secondary infections, in particular with opportunist organisms, are often found and, with respiratory failure, contribute to the high mor-

Results of lung function tests before and after bronchoalveolar lavage

	Before lavage		After lavage		
	June 1986	Oct 1986	Nov 1986	Jan 1987	Apr 1987
Forced expiratory volume in 1 s (ml)	750	510	630	1020	1450
Forced vital capacity (ml)	870	940	980	1370	1650
Total lung capacity (ml)	1920	1590	1900	2140	2110
Airway resistance (kPa l ⁻¹ s)	0.24	0.75	0.43	0.27	0.34
Alveolar-arterial oxygen difference (kPa)		8.9		5.5	4.1

Address for reprint requests: Dr M P Samuels, Brompton Hospital, London SW3 6HP.

Accepted 28 July 1988

A

B

Radiographic appearances before (A) and after (B) bronchoalveolar lavage.

tality.² In adults it has been associated with haematological malignancy and silicosis. There has been a single report in association with autoimmunity but without clinically evident disease.³ A 3 year old girl had a low IgA concentration and a high positive smooth muscle antibody titre but no abnormal deposition of immunoglobulin in the lung was detected by fluorescence studies.

Our patient had clinically florid dermatomyositis with a serum dependent defect of neutrophil phagocytosis, but no immunoglobulin abnormalities. Lung biopsy showed no evidence of vasculitis or interstitial lung disease, although an intercostal muscle biopsy performed at the same time showed evidence of myositis. Whether the dermatomyositis has any bearing on the development of lipoproteinosis is questionable. Patients with alveolar lipoproteinosis have frequently either been taking corticosteroids at the onset or been treated with them. Patients having corticosteroids for other conditions, however, do not develop alveolar proteinosis. The initial trigger in our patient is likely to have been a respiratory infection, with either dermatomyositis, corticosteroid, or the serum defect (or a combination of these) causing an abnormal response in surfactant uptake. Abnormalities have been found both in type II alveolar cells and in alveolar macrophages, both of which are concerned in surfactant uptake and recycling.¹

Spontaneous remission may occur in patients with pulmonary alveolar proteinosis,⁴ but bronchoalveolar lavage is indicated in those with appreciable dyspnoea, PaO_2 less than 8.7 kPa, or an alveolar-arterial PO_2 difference of more than 4.3 kPa. The primary effect of lavage is mechanical with immediate clinical and physiological improvement followed over several weeks by radiographic improvement.⁵ A single lavage may cause complete remission, as in our patient, suggesting that the primary abnormality is no longer present.

Lavage has sometimes been performed in small children and infants with cardiopulmonary bypass, to overcome hazards and technical difficulties.⁶ Fortunately, our patient was able to tolerate the smallest available double lumen

endotracheal tube, though a double lumen Swan-Ganz catheter passed transbronchoscopically has been used in a smaller child.⁷ Whether this method or some form of extracorporeal circuit in infants and young children will improve outcome is unknown.

Pulmonary alveolar proteinosis occurs in neonates with clinical and radiographic features similar to those of hyaline membrane disease.⁸ It may be a cause of chronic respiratory distress in both infants and older children.⁹ In the absence of an infective agent, where immunocompetence is impaired this diagnosis should be considered.

References

- 1 Claypool WD, Rogers RM, Matuschak GM. Update on the clinical diagnosis, management and pathogenesis of pulmonary alveolar proteinosis (phospholipidosis). *Chest* 1984;85:550-8.
- 2 Colon AR, Lawrence RD, Mills SD, O'Connell EJ. Childhood pulmonary alveolar proteinosis. Report of a case and review of the literature. *Am J Dis Child* 1971;121:481-5.
- 3 Gray ES. Autoimmunity in childhood pulmonary alveolar proteinosis. *Br Med J* 1973;iii:296-7.
- 4 Kariman K, Kylstra JA, Spock A. Pulmonary alveolar proteinosis: prospective clinical experience in 23 patients for 15 years. *Lung* 1984;162:223-31.
- 5 Gale ME, Karlinsky JB, Robins AG. Bronchopulmonary lavage in pulmonary alveolar proteinosis: chest radiograph observations. *Am J Radiol* 1986;146:981-5.
- 6 Seard C, Wasserman K, Benfield JR, Cleveland RJ, Costley DO, Heimlich EM. Simultaneous bilateral lung lavage (alveolar washing) using partial cardiopulmonary bypass. Report of two cases in siblings. *Am Rev Respir Dis* 1970;101:877-84.
- 7 Moazen F, Schmidt JH, Chesrown SE, et al. Total lung lavage for pulmonary alveolar proteinosis in an infant without the use of cardiopulmonary bypass. *J Pediatr Surg* 1985;20:398-401.
- 8 Knight DP, Knight JA. Pulmonary alveolar proteinosis in the newborn. *Arch Pathol Lab Med* 1985;109:529-31.
- 9 Coleman M, Dehner L, Sibley RK, Burke BA, L'Heureux PR, Thompson JR. Pulmonary alveolar proteinosis: an uncommon cause of chronic neonatal respiratory distress. *Am Rev Respir Dis* 1980;121:583-6.