Lung function 22–35 years after treatment of idiopathic spontaneous pneumothorax with talc poudrage or simple drainage

PETER LANGE, JANN MORTENSEN, STEFFEN GROTH

From Medical Department P and Chest Clinic and the Department of Clinical Physiology and Nuclear Medicine, Bispebjerg Hospital, Copenhagen, Denmark

ABSTRACT The long term effects on lung function of treatment of idiopathic spontaneous pneumothorax by talc poudrage or simple chest drainage were assessed in 114 patients. Subjects were studied 22–35 years after their first pneumothorax with measurement of static and dynamic lung volumes. Eighty patients had been treated with talc pleurodesis and 34 with simple pleural drainage. At the follow up examination 17 had died from what appear to be unrelated causes and two had emigrated. Of the remainder, 75 participated fully in the study, 14 completed a postal questionnaire, and six failed to respond. The group treated with talc poudrage showed a mild restrictive impairment of lung function with a mean total lung capacity (TLC) of 89% predicted. In subjects treated with simple drainage mean TLC was 96% predicted. One subject treated with talc poudrage had extensive pleural calcification and a substantial reduction in lung function (TLC 58% predicted) with some evidence of lung fibrosis. None of the subjects had developed a mesothelioma. We conclude that, although talc poudrage may cause a mild restrictive impairment of lung function and pleural thickening on the chest radiograph, the long term outlook is good.

Introduction

Idiopathic spontaneous pneumothorax is a relatively rare condition with an incidence of about four or five cases/100 000 people a year.^{1 2} It occurs in previously healthy young adults and is caused by a rupture of a small, often apically located, pulmonary bleb.³ The male to female ratio is about 6:1.

There has been some interest in the effect of pneumothorax on lung function at the time of the pneumothorax^{4 5} or shortly after resolution.^{6–9} There are few reports, however, on the long term effect of spontaneous pneumothorax on pulmonary function^{2 10} and insufficient patients have been studied to assess the impact of the initial treatment.

We studied lung function in subjects who were still alive 22 years or more after a first episode of idiopathic spontaneous pneumothorax. Results were related to the initial treatment, which was either talc poudrage or simple drainage. Particular interest was directed to

possible late effects of talc poudrage, including malignancy.

Methods

PATIENTS

The study population consisted of 114 consecutive patients treated at the chest clinics at Øresundshospitalet and Bispebjerg Hospital, Copenhagen, for their first episode of idiopathic spontaneous pneumothorax during 1951–63. The diagnosis was based on the presentation of a spontaneous pneumothorax in a patient with no prior history of pulmonary disease.

Ninety nine patients were men and 15 women. At the time of the first pneumothorax 99 (87%) of the subjects were under 41 years of age. The initial treatment was talc poudrage followed by drainage via an intercostal tube¹¹ in 60 patients and simple drainage via an intercostal tube in 54 patients. In the latter group 20 patients (37%) had an ipsilateral relapse and this was treated with talc poudrage in all instances. Among the 80 patients treated with talc poudrage for their first episode or for a relapse there was relapse at the side of treatment in two patients (2.5%).

The 80 patients treated with talc poudrage either for

Address for reprint requests: Dr Peter Lange, Medical Department P/ Chest Clinic, Bispebjerg Hospital, Bispebjerg Bakke 23, DK-2400 Copenhagen NV, Denmark.

Accepted 14 April 1988

the first episode of pneumothorax (60 patients) or for the recurrence of pneumothorax (20 patients) have been designated group 1 and the 34 patients treated with simple drainage only group 2.

In autumn 1985 all subjects were traced through the population register. Seventeen had died and two had emigrated. The remaining 95 were invited to undergo an interview and physical examination. Seventy five consented. Of the remainder, 14 filled in a questionnaire but did not wish to attend the examination centre, mainly for reasons of geographical distance; six failed to respond but we were able to ascertain that they had not been in contact with a local chest clinic recently.

PULMONARY INVESTIGATION

Forced expiratory volume in one second (FEV₁) and vital capacity (VC) were measured with a Pulmonet III (Gould) spirometer, the best of at least three recordings being used. Total lung capacity (TLC) and residual volume (RV) were measured by a closed circuit helium equilibrium method. The examinations were carried out according to the recommendations of the European Steel and Coal Union.¹² The results were converted to BTPS and related to predicted values.¹³ Chest radiographs in two projections were obtained. Medical details recorded after the first episode of spontaneous pneumothorax were noted.

ANALYSIS OF RESULTS

Results were analysed by Student's *t* test for unpaired data and the χ^2 test.

Results

Seventeen of the 114 subjects had died. Three had died from lung cancer but none of the remaining deaths was

due to respiratory failure or mesothelioma.

Mean values for TLC, VC and FEV₁ were lower than predicted (table), but the impairment was slight and in most subjects values were within the normal range. Subjects in group 1 (talc poudrage) had significantly lower TLC values than group 2.

Of the 75 patients who had a chest radiograph, 51 showed pleural thickening. There was no significant difference in the incidence of pleural changes on the chest radiograph between group 1 (76%) and group 2 (55%). Subjects with pleural thickening had significantly lower values for TLC (89% predicted) and VC (80% predicted) than those without pleural thickening (TLC 98% predicted and VC 88% predicted). The pleural changes were usually small but two subjects treated with talc poudrage had more extensive pleural thickening with calcification. One of the two subjects, a woman, was the only subject with severe restrictive impairment of lung function (TLC 58% predicted, FEV₁/VC 77%). She had had three pneumothoraces (one on the left side and two on the right) and had been treated with talc pleurodesis on the left side and hypertonic glucose pleurodesis on the right, but both treatments had been complicated by persistent pleural effusions requiring several pleurocenteses. Further investigations, including computed tomography of the thorax and measurement of transfer factor for carbon monoxide (9.6 ml/min/mm Hg: 48% predicted) and pulmonary epithelial permeability to inhaled technetium-99m labelled DTPA (3.9%/min: 450% predicted) suggested that she also had a moderate degree of pulmonary fibrosis.

The mean TLC in subjects with recurrent pneumothorax was significantly lower (88% predicted) than that in subjects with no recurrence (95% predicted).

Fourteen subjects had airflow limitation (FEV₁/

Mean (SD) lung function values

	TLC				VC				FEV ₁			
	Litres	% pred	p		Litres	% pred	p		Litres	% pred	p	
Total studied (n = 75)	6.31 (1.05)	92 (15)			3.64 (0.81)	83 (15)			2.58 (0.78)	74 (19)		
Talc poudrage (n = 46)	6.31 (1.20)	89 (15)	} <0.05		3.68 (0.88)	82 (16)	} NS		2.54 (0.84)	72 (21)	} NS	
No talc poudrage (n = 29)	6.30 (0.77)	96 (12)			3.57 (0.67)	85 (11.5)			2.63 (0.66)	78 (15)		
Pleural changes (n = 51)	6.14 (0.94)	89 (13)	} <0.01		3.55 (0.80)	80 (15)	} <0.025		2.51 (0.77)	72 (18)	} NS	
No pleural changes (n = 24)	6.64 (1.18)	98 (15)			3.81 (0.80)	88 (13)			2.70 (0.80)	78 (20)		

VC $\leq$ 65% and RV/TLV $\geq$ 40%), and in five this was severe (FEV₁ $\leq$ 50% predicted). All but two were lifelong heavy smokers.

Discussion

This study shows that idiopathic spontaneous pneumothorax may be followed by a mild restrictive impairment of lung function. This restrictive pattern was associated with pleural thickening.

In a five to six year follow up of 58 subjects with spontaneous pneumothorax Hallgrímsson found that four of the eight subjects with restrictive impairment of lung function (TLC $\leq$ 80% predicted) had pleural adhesions.² A similar association between pleural thickening and a restrictive impairment of lung function has been suggested in other studies of the short term changes after spontaneous pneumothorax^{7, 8} and in studies of pleural adhesions and pleural thickening after artificial pneumothorax.^{14, 15} It has been suggested that spontaneous pneumothorax may be an early sign of pulmonary emphysema,^{7, 8} but our results do not support this hypothesis.

In our study subjects treated with talc poudrage had slightly more impairment of lung function than those treated with simple drainage. The restrictive changes did not, however, cause symptoms except in one subject. In this subject the impairment of lung function that followed talc poudrage was probably not due to pleural thickening alone, as there was some evidence of lung fibrosis in addition.

Asbestos pollution of the talc has been suggested as a risk factor for the development of mesothelioma.¹⁶ This was not seen in this study despite our observation time of 22–35 years. As the latent period between asbestos exposure and the development of mesothelioma has been longer than 40 years in some studies our work does not entirely exclude the possibility of asbestos pollution of the talc.

When the choice is made between the greater risk of relapse associated with simple drainage and the transitory discomfort and slightly greater restrictive impairment of lung function with talc poudrage, it is comforting to know that, regardless of the choice, the long term prognosis with regard to lung function is good.

This study was supported by grants from the National Union for the Fight against Tuberculosis and Lung Diseases.

References

- 1 Melton LJ, Hepper NGG, Offord KP. Incidence of spontaneous pneumothorax in Olmsted County, Minnesota: 1950 to 1974. *Am Rev Respir Dis* 1979;**120**: 1379–82.
- 2 Hallgrímsson JG. Spontaneous pneumothorax in Iceland with special reference to the idiopathic type: a clinical and epidemiological investigation. *Scand J Thorac Cardiovasc Surg* 1978;**21**(suppl):9–85.
- 3 Kjaergaard H. Spontaneous pneumothorax in the apparently healthy. *Acta Med Scand* 1932;**43**(suppl): 1–159.
- 4 Gilmartin JJ, Wright AJ, Gibson GJ. Effects of pneumothorax or pleural effusion on pulmonary function. *Thorax* 1985;**40**:60–5.
- 5 Anthonisen NR. Regional lung function in spontaneous pneumothorax. *Am Rev Respir Dis* 1977;**115**:873–6.
- 6 Heckscher TH, Andree Larsen O, Lassen NA. A clinical method for determination of regional lung function using intravenous injection of Xe¹³³. *Scand J Respir Dis* 1966;**62**(suppl):31–9.
- 7 De Troyer A, Yernault JC, Rodenstein D, Englert M, De Coster A. Pulmonary function in patients with primary spontaneous pneumothorax. *Bull Eur Physiopathol Respir* 1978;**14**:31–9.
- 8 Williams MH, Kane C. Pulmonary function in patients who have recovered from spontaneous pneumothorax. *Dis Chest* 1965;**47**:153–6.
- 9 Baronofsky ID, Warden HG, Kaufman JL, Whatley J, Hanner JM. Bilateral therapy for unilateral spontaneous pneumothorax. *J Thorac Surg* 1957;**34**:310–9.
- 10 Fleetham JA, Forkert L, Clarke H, Anthonisen NR. Regional lung function in presence of pleural symphysis. *Am Rev Respir Dis* 1980;**122**:33–8.
- 11 Bethune N. Pleural poudrage—a new technique for the deliberate production of pleural adhesion as a preliminary to lobectomy. *J Thorac Surg* 1935;**4**:251–61.
- 12 Quanjer PH, ed. Standardised lung function testing. Report of working party of the European Community for Coal and Steel. *Bull Eur Physiopathol Respir* 1983;**19**(suppl 5):1–96.
- 13 Dirksen A, Groth S. Calculation of reference values for lung function tests. *Bull Eur Physiopathol Respir* 1986;**22**:231–7.
- 14 Patiala J, Karvonen MJ. Studies on pulmonary function after pneumothorax therapy. *Acta Tuberc Scand* 1954;**29**:193–210.
- 15 Autio V. The reduction of respiratory function by parenchymal and pleural lesions. *Acta Tuberc Scand* 1959;**37**:112–25.
- 16 Research Committee of the British Thoracic Association and the Medical Research Council Pneumoconiosis Unit. The survey of the long-term effects of talc and kaolin pleurodesis. *Br J Dis Chest* 1979;**73**:285–8.