Effect of a 5-lipoxygenase inhibitor on leukotriene generation and airway responses after allergen challenge in asthmatic patients

Kok P Hui, Ian K Taylor, Graham W Taylor, Paul Rubin, James Kesterson, Neil C Barnes, Peter J Barnes

Abstract

The effect of a single oral dose (800 mg) of zileuton (A-64077), a specific 5-lipoxygenase inhibitor, on the early and late airway responses to inhaled allergen was studied in a randomised, double blind, placebo controlled, and crossover trial in nine subjects with atopic asthma. Leukotriene generation was also assessed in vivo by measuring urinary leukotriene (LT) E₄ excretion, and ex vivo by measuring calcium ionophore stimulated whole blood LTB₄ production. Zileuton almost completely inhibited ex vivo LTB₄ production but reduced urinary excretion of LTE₄ by only about half. There was a trend for the early asthmatic response to be less on the day of zileuton treatment, but this did not reach statistical significance ($p = 0.08$). The zileuton induced reduction in maximum fall in FEV₁ in the early asthmatic response was, however, significantly related to the reduction in urinary LTE₄ excretion ($r = 0.8$), but not to the reduction in LTB₄ generation ex vivo. There was no significant change in the allergen induced late asthmatic response, or in the increase in airway responsiveness to methacholine following antigen. The results provide some support for the hypothesis that the cysteinyl leukotrienes have a role in the allergen induced early asthmatic response. More complete in vivo inhibition of 5-lipoxygenase may be needed to produce a significant reduction in airway response to allergen challenge.

Department of
Thoracic Medicine,
National Heart and
Lung Institute,
London SW3 6LY
K P Hui
P J Barnes

Department of
Clinical
Pharmacology, Royal
Postgraduate Medical
School, London
W12 0NN
I K Taylor
G W Taylor

Department of
Thoracic Medicine,
London Chest
Hospital, London E2
9JX
N C Barnes

Abbott Laboratories,
Abbott Park, Illinois
60064, USA
P Rubin
J Kesterson

Reprint requests to:
Dr N C Barnes

Accepted 3 December 1990

The leukotrienes, products of the 5-lipoxygenase pathway of arachidonic acid metabolism, are thought to be important mediators in the pathogenesis of asthma because their biological activities produce changes that are similar to those seen in asthma.¹⁻³ The cysteinyl leukotrienes, LTC₄, LTD₄, and LTE₄, are potent bronchoconstrictor agents in normal and asthmatic subjects,⁴⁻⁶ and LTE₄ has been reported to increase airway hyper-responsiveness.⁷ LTB₄ is a potent chemotactic agent for leucocytes,⁸ and may be important in mediating the inflammatory process in asthmatic airways. LTD₄ can increase mucus production in human airway preparations⁹ and airway microvascular leakage in animals.¹⁰ Several clinical studies with cysteinyl leukotriene antagonists in asthmatic subjects have

shown a slight reduction in the early asthmatic response to inhaled allergen^{11,12} and in cold air induced bronchoconstriction.¹³ The compounds studied were relatively weak cysteinyl leukotriene antagonists, however, and they would in addition leave the effects of the dihydroxy acid LTB₄ unopposed. An alternative approach is to inhibit the 5-lipoxygenase enzyme to reduce synthesis of all the leukotrienes. Several different classes of 5-lipoxygenase inhibitors have been found to be effective in animal models of asthma.¹⁴⁻¹⁶ No significant reduction of allergen induced bronchoconstriction was, however, proposed by piriprost¹⁷ or nafazatrom¹⁸ in two clinical studies in man—but nafazatrom was found not to prevent ex vivo leukotriene production in man after oral dosing despite having 5-lipoxygenase inhibitory activity in vitro (the activity of piriprost in man was not studied).

After exposure to inhaled allergen there is an early and late bronchoconstrictor response in asthmatic subjects and an increase in airway responsiveness. Several mediators, including the leukotrienes, are generated during antigen challenge and may be important in mediating the airway responses. Zileuton, a hydroxamic acid 5-lipoxygenase inhibitor,¹⁹ has been shown to inhibit calcium ionophore stimulated human neutrophil production of LTB₄ in man ex vivo after oral dosing,²⁰ and is effective in inhibiting allergen induced contraction of tracheal smooth muscle in the guinea pig.²¹ We have investigated the efficacy of zileuton (800 mg) on the generation of cysteinyl leukotriene in vivo (assessed by urine LTE₄ concentration) induced by inhaled allergen and on the generation of LTB₄ ex vivo (by calcium ionophore stimulated whole blood) in asthmatic subjects. We assessed the inhaled allergen induced airway responses of the early and late asthmatic response and changes in airway responsiveness.

Methods

SUBJECTS

Eleven non-smoking men with atopic asthma (mean age 28, range 19-44 years) with an FEV₁ above 70% predicted were recruited (table). Subjects were having inhaled salbutamol only, apart from one who was also inhaling steroids and taking oral theophyllines. All subjects had been shown to have a dual asthmatic response to allergen on screening. None had had an upper respiratory tract infection or exacerbation of asthma

within six weeks of starting the study. Non-steroidal anti-inflammatory drugs were not allowed during the study period. The study was approved by the ethics committee of the National Heart and Lung Hospitals, and fully informed written consent was obtained from each subject.

Subjects had to be healthy at the physical examination and to have normal results in biochemical and haematological tests and a normal electrocardiogram before entry into the study. These were repeated during the study and at the end to assess the safety and tolerability of zileuton.

STUDY PROTOCOL

Allergen challenge

On a screening day the atopy of each subject was confirmed by obtaining a positive skin-prick test response to either *Dermatophagoides pteronyssinus* or mixed grass pollen allergen extracts (Pharmacia, Milton Keynes) dissolved in normal saline. Subjects first inhaled the diluent, aerosolised with a Wright nebuliser (volume 2 ml, flow rate 7 l/min, output 0.2 ml/min), through a face mask with open mouthed tidal breathing for two minutes, and FEV₁ was then measured for 15 minutes to exclude a bronchoconstrictor response to diluent. Cumulative doses of allergen were then inhaled in the same way to identify subjects with a dual asthmatic response. The initial allergen concentration used for inhalation was that which caused a 2–3 mm diameter weal in the skinprick test. Doubling concentrations of allergen were inhaled until an early asthmatic response (fall in FEV₁ of at least 20% of the post-diluent value) was seen. No further allergen was then given, FEV₁ was measured up to eight hours to detect any late asthmatic response, which was defined as a fall in FEV₁ of at least 15% of post-diluent value. If a dual response was seen, the final allergen dose reached was used on the two allergen challenge study days. Of the 17 subjects screened, six subjects with an isolated early asthmatic response were not enrolled into the main study.

Two to three weeks later subjects returned for the first allergen challenge study. Zileuton (800 mg) or matching placebo was ingested in the morning, and allergen challenge was performed three hours later. FEV₁ was measured

every 10 minutes for the first hour after challenge, and then hourly for the next seven hours. At least two weeks later subjects returned for the second study to continue with an identical protocol provided that the baseline FEV₁ was within 15% of that on the first study day. If not, they were asked to return for another study day.

FEV₁ was measured with a dry bellow spirometer (Vitalograph, Buckingham) with the subject sitting upright. The mean of three good expiratory efforts was taken. Before each test subjects rested in a warm room for at least 15 minutes.

Non-specific airway responsiveness was assessed according to the method of Cockcroft *et al*²² on the day before and after each allergen test day, at the same time of the day for each subject. The provocative dose of methacholine needed to reduce FEV₁ by 20% of the post-diluent value (PC₂₀ FEV₁) was determined. Methacholine was inhaled in the same way as allergen.

Assessment of 5-lipoxygenase inhibition

5-Lipoxygenase activity after allergen challenge was assessed by measuring urinary LTE₄ excretion.²³ Immediately before challenge subjects emptied their bladder and urine was then collected for four hours. Urine volume and pH were measured and aliquots of 30 ml were stored at –70°C. LTE₄ was extracted from the urine by reverse phase high performance liquid chromatography, tritiated LTE₄ being used as an internal standard,²⁴ and measured by radioimmunoassay (Amersham International, Amersham). The efficacy of zileuton was also assessed by calcium ionophore stimulated whole blood LTB₄ production *ex vivo*. Three and a half, four, five, seven, nine, and 10 hours after it was given 5 ml of whole blood was incubated with 5 µl of 50 mM calcium ionophore for 15 minutes and centrifuged, and the supernatant was stored at –70°C. LTB₄ was extracted by high performance liquid chromatography, and assayed by radioimmunoassay.

Drug level

Plasma level of drug was measured at 3.5, 4 and 5 hours post-dose on both placebo and active days. Zileuton was extracted by high performance liquid chromatography and measured by ultraviolet absorbance as developed by Abbott Laboratories.

ANALYSIS

PC₂₀ values were log transformed before analysis. FEV₁ and PC₂₀ values were compared by analysis of variance. Paired data were analysed by the Wilcoxon sign rank test and the relation between two variables by regression analysis. Results are presented as means with the standard errors of the mean in parentheses unless otherwise stated. *p* < 0.05 was accepted as significant.

Results

SUBJECTS

Of the 11 subjects who showed a late response

Characteristics of the asthmatic subjects

Subject No	Age (y)	FEV ₁ l (% pred)	PC ₂₀ methacholine* (mg/ml)	Drugs†
1	22	4.9 (98)	0.44	Salbutamol
2	33	3.0 (81)	0.19	Salbutamol
3	21	4.9 (117)	0.2	Salbutamol
4	22	4.1 (98)	0.26	Salbutamol
5	24	4.8 (117)	3.4	Salbutamol
6	27	4.3 (100)	0.15	Salbutamol
7	27	4.6 (105)	1.18	Salbutamol
8	23	2.6 (84)	0.57	Salbutamol
9	29	4.6 (112)	1.24	Salbutamol, beclomethasone, theophylline

*The provocative dose of inhaled methacholine needed to cause a 20% fall in baseline FEV₁.

†Inhaled, apart from the theophylline (oral) taken by subject 9.

Figure 1 Effect of zileuton (800 mg orally) on early and late asthmatic responses after allergen challenge. ● Zileuton; ○ placebo. Values are means with 1 SEM.

Figure 3 Time course of effect of zileuton on calcium ionophore stimulated whole blood leukotriene (LT) B_4 production. Values are means with 1 SEM. * $p < 0.005$.

to inhaled allergen, two were withdrawn. One developed an exacerbation of asthma following a chest infection after the first study day (placebo). The other subject who had an FEV_1 of just over 70% predicted was withdrawn when his FEV_1 fell to below 70% predicted before allergen challenge. The remaining nine subjects completed the study, and all results were included in the analysis. Zileuton was well tolerated and there were no important biochemical, haematological, or electrocardiographic changes after treatment.

AIRWAY RESPONSES

Baseline values before allergen challenge on placebo and treatment days were closely matched (fig 1). No period or treatment effect

was found by two way analysis according to the method of Hills and Armitage.²⁵ There was a trend for the fall in FEV_1 during the early asthmatic response to be less with zileuton, but this did not reach statistical significance at any point (maximum fall in FEV_1 : placebo 1.08 (0.25) l, zileuton 0.83 (0.21) l; $p = 0.18$). The maximum difference between placebo and zileuton was 40 minutes after allergen challenge, which is after the nadir of the fall in FEV_1 (fig 1; difference in FEV_1 -0.33 (0.16) l; $p = 0.08$). There was no difference between placebo and zileuton in the late asthmatic response (maximum fall in FEV_1 : placebo 1.16 (0.24) l, zileuton 0.92 (0.27) l; $p = 0.22$). There were no significant differences in the areas under the curve during the early or late phases.

Figure 2 Effect of zileuton on airway responsiveness to inhaled methacholine before and after allergen challenge. $PC_{20} FEV_1$ is the provocative dose (mg/ml) of inhaled methacholine needed to cause a fall in FEV_1 of 20% of the post-diluent value, and individual values for each subject before and after allergen challenge are shown. The geometric mean and 95% confidence interval for each day are also shown.

Figure 4 Effect of zileuton (▨) and placebo (■) on (a) mean ex vivo leukotriene (LT) B_4 production and (b) urine LTE_4 . Mean ex vivo LTB_4 production refers to mean whole blood calcium ionophore stimulated LTB_4 production for four hours after allergen challenge. Urine LTE_4 is the total urinary excretion of LTE_4 over four hours. Values are means with 1 SEM.

Figure 5 Scatter diagrams showing the linear regression relationships between zileuton induced change in maximum fall in FEV_1 during the early asthmatic response (EAR) and reduction in urinary leukotriene (LT) E_4 production.

Baseline airway responses to inhaled metacholine before allergen challenge were very similar for placebo and zileuton (geometric mean PC_{20} $FEV_1 = 0.41$ (95% confidence interval 0.20–0.84) and 0.44 mg/ml (0.15–0.78). The increase in airway responsiveness after allergen challenge did not differ significantly after placebo and zileuton treatment (before/after challenge PC_{20} FEV_1 metacholine: placebo 2.13, zileuton 1.89; fig 2).

EFFECTS ON IN VIVO AND EX VIVO LEUKOTRIENE GENERATION

Zileuton substantially inhibited calcium ionophore stimulated whole blood LTB_4 production with maximum inhibition (93.1% of baseline; $p < 0.005$) four hours post-dose; after the dose had been given significant inhibition was still present at 10 hours after the dose (63.5% of baseline; $p < 0.005$; fig 3). Mean urinary LTE_4 excretion after allergen challenge was reduced by about half by zileuton treatment (placebo 111.5 (23.5), zileuton 58.2 (14.8) ng/mmol creatinine; $p < 0.01$; fig 4).

The zileuton induced change in maximum fall in FEV_1 for the early asthmatic response correlated with the reduction in urinary LTE_4 production ($r = 0.8$, $p < 0.01$; fig 5) but not with the inhibition of calcium ionophore stimulated whole blood LTB_4 production ($r = 0.04$, $p > 0.5$).

PLASMA ZILEUTON CONCENTRATIONS

Mean plasma zileuton concentrations three and a half, four, and five hours after ingestion were 3.37 (0.36), 3.03 (0.39) and 2.6 (0.27) $\mu\text{g/ml}$ respectively on the days of zileuton treatment; no drug was detected on the placebo day. There was no relation between the peak drug concentrations and change in FEV_1 during the early and late asthmatic responses, reduction in urinary LTE_4 excretion, or calcium ionophore stimulated whole blood LTB_4 production.

Discussion

We found that zileuton at the dose used in the study partially reduced urinary LTE_4 excretion

and substantially reduced calcium ionophore stimulated whole blood LTB_4 production ex vivo. There was a trend towards a reduction in the fall in FEV_1 during the early asthmatic response with zileuton, but neither this nor the late asthmatic response or increase in airway responsiveness differed significantly between zileuton and placebo. The change in the early asthmatic response induced by zileuton correlated with the change in urinary LTE_4 concentrations but not with the change in ex vivo LTB_4 production.

In vivo activation of the 5-lipoxygenase pathway in asthmatic patients during an asthmatic attack has been shown by the increased urinary excretion of LTE_4 ,²³ and by increased concentrations of cysteinyl leukotrienes in bronchoalveolar lavage fluid recovered by fiberoptic bronchoscopy during the early²⁶ and late²⁷ responses to inhaled allergen challenge. In the present study the activation of 5-lipoxygenase in airways was assessed by urinary excretion of LTE_4 as this is simple and non-invasive and did not interfere with measurements of lung function. Urinary LTE_4 excretion after allergen challenge is likely to reflect airway leukotriene generation, as there was no evidence of systemic effects in any subjects. Zileuton reduced the urinary excretion of LTE_4 by about half, though not down to the range found in non-asthmatic subjects in this laboratory.²³ This suggests that 5-lipoxygenase activity was only partially inhibited, and may explain the non-significant reduction in bronchoconstriction during the early asthmatic response. A significant reduction of airway response to allergen may be seen only if more complete inhibition of leukotrienes generation can be achieved, as the leukotrienes are very potent biological agents: the cysteinyl leukotrienes are thousands of times more potent than histamine in causing bronchoconstriction,²⁸ and LTB_4 is one of the most potent chemotactic agents known.²⁹

The early bronchoconstrictor response after allergen challenge is thought to be due to release of mediators, such as histamine and leukotrienes, from inflammatory cells via IgE mediated mechanisms. Although not statis-

tically significant, the maximum reduction of the fall in FEV₁ produced by zileuton was after the nadir in the fall in FEV₁ following allergen challenge. This time course of action is consistent with the known action of leukotrienes and histamine, inhaled cysteinyl leukotrienes causing maximum bronchoconstriction at about 15 minutes,^{4,6} and histamine somewhat earlier. Leukotrienes are not stored by resting cells (unlike histamine), and are generated only after stimulation, so they would be expected to contribute more to the latter part of the early bronchoconstrictor response after allergen challenge. In a study by Britton and coworkers, in which a leukotriene antagonist, L649923, reduced the early response to allergen, the maximum effect was seen after the nadir of the early asthmatic response.¹¹ In another study an antihistamine was effective in the first 15 minutes of the early asthmatic response after allergen challenge.³⁰ The significant correlation between inhibition of in vivo LTE₄ generation and reduction in early asthmatic response also supports the role of the leukotrienes as mediators in the early asthmatic response. Although the pattern of change in the early asthmatic response is similar to that in the study of Britton *et al*,¹¹ the lack of statistical significance in our study may be due to a type II error.

Although leukotriene production has been shown during the late asthmatic response following allergen challenge,²⁷ zileuton did not reduce the late fall in FEV₁ significantly in the present study. This may be because the activity of zileuton in the single dose of 800 mg used in this study may have been insufficient by that time, particularly when it was only partially effective during the early phase as measured by urinary LTE₄ levels. In two patients where serum drug concentrations seven hours after the dose were available little zileuton was detected.

Previous studies of 5-lipoxygenase inhibitors in man have depended on ex vivo methods of assessing its activity. In the present study zileuton was more effective in inhibiting ex vivo calcium ionophore stimulated whole blood LTB₄ production than in inhibiting leukotriene production in vivo as reflected by urinary LTE₄ excretion, and there was no correlation between these two variables. There are several possible explanations. The lesser effect in vivo may be due to inadequate penetration of zileuton into airway tissues, or a higher plasma concentration may be needed to suppress airways 5-lipoxygenase adequately. Alternatively, different inflammatory cells may have different susceptibilities to the effect of zileuton, such that the mast cells, macrophages, or eosinophils in the airways may not have been affected as much as neutrophils in the peripheral blood. In addition, the cellular generation of the cysteinyl leukotrienes may be less affected than that of LTB₄. The lack of relation between the changes in leukotriene generation ex vivo and the early asthmatic response suggest that this is not a useful method for assessing the clinical efficacy of 5-lipoxygenase inhibitors in man.

In this study a single oral dose (800 mg) of

zileuton was only partially effective in inhibiting activation of 5-lipoxygenase in vivo by inhaled allergen despite near complete ex vivo inhibitory activity. There was a trend towards a reduction in the early asthmatic bronchoconstrictor response but this was not significant. Measurement of leukotriene production in vivo rather than ex vivo may be more useful in future studies of 5-lipoxygenase inhibitors in man. Our results would fit a role for the leukotrienes in the early asthmatic response to inhaled allergen, particularly if inhibition of leukotriene related airway responses requires near complete inhibition of the 5-lipoxygenase enzyme in vivo. Further studies with higher or repeated doses of zileuton or with more potent 5-lipoxygenase inhibitors in man are needed.

- 1 Samuelsson B. Leukotrienes: mediators of immediate hypersensitivity reactions and inflammation. *Science* 1983; 220:568-75.
- 2 Hannarström S. Leukotrienes. *Ann Rev Biochem* 1983; 52:355-77.
- 3 Drazen JM, Austen KF. Leukotrienes and airway responses. *Am Rev Respir Dis* 1987;136:985-8.
- 4 Barnes NC, Piper PJ, Costello JF. Comparative effects of inhaled leukotriene C₄, leukotriene D₄, and histamine in normal human subjects. *Thorax* 1984;39:500-4.
- 5 Adolph E, Morris MM, Hargreave FE, O'Byrne PM. Airway responsiveness to leukotrienes C₄ and D₄ and to methacholine in patients with asthma and normal controls. *N Engl J Med* 1986;315:480-4.
- 6 Davidson AB, Lee TH, Scanlon PD, *et al*. Bronchoconstrictor effects of leukotriene E₄ in normal and asthmatic subjects. *Am Rev Respir Dis* 1987;135:333-7.
- 7 Arm JA, Spur BW, Lee TH. The effects of inhaled leukotriene E₄ on the airway responsiveness to histamine in subjects with asthma and normal subjects. *J Allergy Clin Immunol* 1988;82:654-60.
- 8 Ford-Hutchinson AW, Bray MA, Doig MV, Shipley ME, Smith MJH. Leukotriene B₄, a potent chemokinetic and aggregating substance from polymorphonuclear leukocytes. *Nature* 1980;286:264-5.
- 9 Marom S, Shelhamer JH, Bach MK, Morton DR, Kaliner M. Slow reacting substances leukotrienes C₄ and D₄ increase the release of mucus from human airways in vitro. *Am Rev Respir Dis* 1982;126:449-51.
- 10 Evans TW, Rogers DF, Aursudkig B, Chung KF, Barnes PJ. Regional and time-dependent effects of inflammatory mediators on airway microvascular permeability in the guinea pig. *Clin Sci* 1989;76:161-6.
- 11 Britton JR, Hanley SP, Tattersfield AE. The effect of an oral leukotriene D₄ antagonist L649,923 on the response to inhaled antigen in asthma. *J Allergy Clin Immunol* 1987;79:811-6.
- 12 Fuller RW, Black PN, Dollery CT. Effects of the oral leukotriene D₄ antagonist LY 171883 on inhaled and intradermal antigen and LTD₄ in atopic subjects. *J Clin Allergy Immunol* 1989;83:939-44.
- 13 Israel I, Juniper EF, Callaghan JT, *et al*. Effect of a leukotriene antagonist, LY 171883, on cold air-induced bronchoconstriction in asthmatics. *Am Rev Respir Dis* 1989;140:1348-53.
- 14 Kruetner W, Sherwood J, Sehring S, *et al*. Antiallergic activity of Sch 37224, a new inhibitor of leukotriene formation. *J Pharmacol Exp Ther* 1988;247:997-1003.
- 15 Johnson HG, Stout BK. Activity of a novel hydroquinone inhibitor of leukotriene synthesis (U-66,858) in the rhesus monkey *Ascaris* reactor. *Int Arch Allergy Appl Immunol* 1988;87:204-7.
- 16 Ku EC, Raychaudhuri A, Ghai G, *et al*. Characterisation of CGS 8515 as a selective 5-lipoxygenase inhibitor using in vitro and in vivo models. *Biochim Biophys Acta* 1988; 959:332-42.
- 17 Mann JS, Holgate ST. Effect of piriprost (U-60257), a novel lipoxygenase inhibitor, on allergen and exercise induced asthma. *Thorax* 1986;41:746-52.
- 18 Fuller RW, Maltby N, Richmond R, *et al*. Oral nafazatrom in man: effect on inhaled antigen challenge. *Br J Clin Pharmacol* 1987;23:677-81.
- 19 Carter GW, Young PR, Albert D, *et al*. Development of A-64077, a promising agent for the treatment of leukotriene-mediated diseases [abstract]. In: *Proceedings of the 7th International Conference on Prostaglandins and Related Compounds*. 1990:192.
- 20 Summers JB, Gunn BP, Martin JG, *et al*. Structure-activity

- analysis of a class of orally active hydroxamic acid inhibitors of leukotriene biosynthesis. *J Med Chem* 1988;31:1960-4.
- 21 Malo PE, Wiley YD, Shaughnessy TK, Bell R, Summers JB, Brooks DW, Carter GW. Inhibition of antigen-induced contractions of guinea-pig tracheal strips (GPTS) by lipoxygenase (5-LO) inhibitors [abstract]. *Am Rev Respir Dis* 1988;139(suppl):A92.
 - 22 Cockcroft DW, Killian DN, Mellon JJA, Hargreave FE. Bronchial reactivity to inhaled histamine: a method and clinical survey. *Clin Allergy* 1977;7:235-43.
 - 23 Taylor GW, Taylor I, Black P, *et al.* Urinary leukotriene E₄ after antigen challenge and in acute asthma and allergic rhinitis. *Lancet* 1989;i:584-8.
 - 24 Richmond RR, Turner NC, Maltby NH, *et al.* Single step procedure for the extraction and purification of leukotrienes B₄, C₄ and D₄. *J Chromatology* 1987;417:241-51.
 - 25 Hills M, Armitage P. The two-period cross-over trial. *Br J Clin Pharmacol* 1979;5:7-20.
 - 26 Chan-Yeung M, Chan H, Tse KS, Salari H, Lam S. Histamine and leukotrienes release in bronchoalveolar fluid during plicatic acid-induced bronchoconstriction. *J Allergy Clin Immunol* 1989;84:762-8.
 - 27 Dias P, Gonzalez MC, Galleguillos FR, *et al.* Leukocytes and mediators in bronchoalveolar lavage during allergen-induced late phase reactions. *Am Rev Respir Dis* 1989;139:1383-9.
 - 28 Dahlén SE, Hedqvist P, Hammarström S, Samuelsson B. Leukotrienes are potent constrictors of human bronchi. *Nature* 1980;288:484-6.
 - 29 Bray MA. The pharmacology and pathophysiology of leukotriene B₄. *Br Med Bull* 1983;39:249-54.
 - 30 Rafferty R, Ng WH, Phillips G, *et al.* The inhibitory actions of azelastine hydrochloride on the early and late bronchoconstrictor responses to inhaled allergen in atopic asthma. *J Allergy Clin Immunol* 1989;84:649-57.