Asthma following occupational exposure to *Lycopodium clavatum* in condom manufacturers

P Cullinan, J Cannon, D Sheril, A Newman Taylor

Department of Occupational and Environmental Medicine, Royal Brompton National Heart and Lung Institute, London SW3 6LR
P Cullinan
J Cannon
A Newman Taylor
343 Ripple Road, Barking, Essex
D Sheril

Reprint requests to:
Dr P Cullinan

Received 15 June 1992
Returned to authors
29 July 1992
Revised version received
17 September 1992
Accepted 6 October 1992

Abstract

Occupational asthma in two women employed in the manufacture of condoms is reported. The spores of *Lycopodium clavatum*, used as a rubber dusting agent, were identified as the causative agent.

(Thorax 1993;48:774-775)

Case reports

CASE 1

A 46 year old woman, a non-smoker without previous respiratory problems, presented with an 18 month history of cough, wheeze, eye irritation, and pruritus. Her symptoms occurred during and following periods at work and resolved when away from work.

For four years she had worked on the production line of a condom factory where she loaded latex condoms onto templates before electronic testing. The condoms were pre-coated with a mixture of lycopodium and

maize starch dust to prevent adherence. During one shift the patient might load 20 000 condoms in a dusty atmosphere and in order to complete the shift she used repeated doses of inhaled salbutamol. Respiratory protection was not worn.

She was referred after four weeks of sick leave. At this time she had a forced expiratory volume in one second (FEV₁) of 2.15 l and a forced vital capacity (FVC) of 2.55 l, improving after inhalation of a bronchodilator to FEV₁ 2.25 and FVC 2.60 l. Her histamine PC₂₀ was greater than 16 mg/ml. She had immediate skin prick reactions to extracts of grass pollen, cat fur, house dust, *Lycopodium clavatum*, and maize starch. Specific IgE antibodies to an extract of natural latex were absent. Her chest radiograph was normal.

Single blind inhalation tests, with the patient wearing latex gloves, were conducted on separate days using a dust tipping method in a sealed chamber.¹ Thirty minutes exposure to a control dust (lactose) provoked no change in her baseline FEV₁ of 2.20 l (fig). Exposure for the same duration to 0.1% *Lycopodium clavatum* in lactose provoked a 12% fall in FEV₁ 10 minutes after the challenge, and a smaller fall (maximum 6%) at 12 hours. Exposure to a 1.0% mixture for 30 minutes provoked both an early response (maximum 31% fall in FEV₁ at 15 minutes) and a less marked late response, starting nine hours after the test inhalation with a maximum fall in FEV₁ of 14% at 12 hours. The immediate response was accompanied by nasal symptoms including epistaxis, but no pruritus. Her histamine PC₂₀ remained unchanged throughout the period of testing. Four days later she was exposed in the same way to maize starch, provoking itchy skin immediately after exposure but no asthmatic reaction.

Following these findings the patient left the factory with complete resolution of her symptoms.

CASE 2

A 53 year old woman working on the same production line presented with a similar history. She had no past respiratory history but had smoked until 1965. Seven years after starting this work she developed a running nose and itching eyes and, after a further 10 years, shortness of breath and chest tightness. Her symptoms completely resolved when she was not at work. She had immediate skin prick reactions to *Lycopodium clavatum* and maize starch, as well as to common inhalant allergens. Shortly after referral she was found to have widespread gastric carcinoma and further investigations became inappropriate.

Discussion

Lycopodium dust is the spore of the common club moss, *Lycopodium clavatum*, found commonly across central and northern Europe, Asia, and North America. The lycopodium used in the UK is gathered by hand in China

Effect of inhalation of 0.1% and 1.0% *Lycopodium clavatum* on forced expiratory volume in one second (FEV₁) in case 1.

or Russia and exported in unadulterated form. Its uniform spores are approximately 25 μm in diameter and have high concentrations of fatty acids. The resulting fine, emollient powder is useful in the handling of processed rubber products and in the past was used to dust surgical gloves. The powder has also been used by pharmacists as a coating for pills and suppositories, by homeopaths in the treatment of breathing disorders, and in cosmetics. *Lycopodium* dust is flammable and is used in small quantities by professional magicians. Recent technological changes have greatly reduced its industrial use but it continues to be used in condom manufacture and by magicians.

The first reported cases of allergy to *lycopodium* are of occupational asthma in a pharmacist and of rhinitis in a woman using a dry shampoo containing *lycopodium*.² In 1951 Salen³ reported a further 12 cases of asthma, nine in opera singers using a stage makeup based on *lycopodium*, two in women using dry shampoo, and one in a Swedish boy apparently exposed only to *lycopodium* spores in the wild. All the subjects had positive skin reactions to *lycopodium* extracts. In 1988 two of 30 workers at a condom factory in France were reported to have developed occupational asthma attributed to *lycopodium*. One had a positive skin prick test to *Lycopodium clavatum* and both showed reductions in peak flow rate immediately after specific bronchial challenge; no late reactions were reported.⁴ In 1989 a 45 year old atopic man was reported to have developed asthma following recreational exposure to condoms coated with *lycopodium*.⁵ Specific skin prick testing was positive. Although he did not undergo bronchial challenge testing his symptoms resolved when using *lycopodium*-free condoms.

Those working in the manufacture of condoms are also exposed to rubber and to corn starch, both potential sensitizers. Processed rubber is a mixture of natural latex and various chemicals used as accelerators and preservatives: both may provoke contact dermatitis⁶ and occupational asthma attributed to latex has been described in subjects wearing surgical gloves.⁷ Corn starch may also cause urticaria⁸ and anaphylactic reactions.

In the first patient reported here the combination of a dual response to *Lycopodium* spores and the absence of a response to corn starch and of specific IgE antibodies to latex make it probable that her symptoms were attributable to *lycopodium* sensitisation alone. After recognition of the cause of these two cases of asthma in early 1992 the largest producer of condoms in the UK has ceased to use *lycopodium* in their manufacture; unless new manufacturing uses are found for it *lycopodium* as a cause of occupational allergic disease should become an historical curiosity.

- 1 Newman Taylor A, Davies R. Inhalation challenge testing. In: Weill H, Turner-Warwick M, eds. *Occupational lung diseases: research approach and methods*. New York: Marcel Dekker, 1982;143–67.
- 2 Peshkin M. Bronchial asthma and other allergic manifestations in pharmacists. *JAMA* 1924;82:1854–5.
- 3 Salen E. *Lycopodium* allergy. *Acta Allergol* 1951;IV: 308–19.
- 4 Catalina P, Chamoux A, Gabrillargues D, Catalina M, Royfe M, Wahl D. Contribution à l'étude des asthmes d'origine professionnelle: l'asthme à la poudre de *lycopode*. *Arch Mal Prof* 1988;49:143–8.
- 5 Juhlin L. Asthma and rhinitis associated with *lycopodium* spores on condoms. *Lancet* 1989;i:563.
- 6 Nutter A. Contact urticaria to rubber. *Br J Dermatol* 1979;101:597–8.
- 7 Marcos C, Lazaro M, Fraj J, Quirce S, de la Hoz B, Fernandez-Rivas M, et al. Occupational asthma due to latex surgical gloves. *Ann Allergy* 1991;67:319–23.
- 8 Fisher A. Contact urticaria and anaphylactoid reaction due to corn starch surgical glove powder. *Contact Dermatitis* 1987;16:224–5.