jects with asthma or allergic rhinitis. However, provocation with nebulised allergen extract is different from natural exposure to pollen grains. The results of our study show that in subjects with pollen-induced rhinitis the cessation of exposure to pollens is associated with a decrease in methacholine responsiveness.

We would like to thank the volunteers who kindly helped us with this study, Dr Alfonso Romar for statistical analyses, and Ms Maria José Pérez and Bertha Camps for technical assistance. Data of pollen counts were provided by Dr Luis Caballero.

1 Boulet LPh, Morin D, Milot J, Turcotte H. Bronchial responsiveness increases after seasonal antigen exposure in

- non-asthmatic subjects with pollen-induced rhinitis. *Ann Allergy* 1989;63:114-9.
- 2 Moreno RH, Hogg JC, Paré PD. Mechanics of airway narrowing. *Am Rev Respir Dis* 1986;133:1171-80.
- 3 Quanjer PhH (Ed). Standardized lung function testing. *Bull Eur Physiopathol Respir* 1983;19(suppl 5):1-95.
- 4 Cockcroft DW, Killian DN, Mellon JJA, Hargreave FE. Bronchial reactivity to inhaled histamine: a method and clinical survey. *Clin Allergy* 1977;7:235-43.
- 5 Sterk PJ, Daniel EF, Zamel M, Hargreave FE. Limited bronchoconstriction to methacholine using partial flow-volume curves in nonasthmatic subjects. *Am Rev Respir Dis* 1985;132:272-7.
- 6 Aubier M, Levy J, Clerici Ch, Neukirch F, Herman D. Different effects of nasal and bronchial glucocorticosteroid administration on bronchial hyperresponsiveness in patients with allergic rhinitis. *Am Rev Respir Dis* 1992;146:122-6.
- 7 Boonsawat W, Salome ChM, Woolcock AJ. Effect of allergen inhalation on the maximal response plateau of the dose-response curve to methacholine. *Am Rev Respir Dis* 1992;146:565-9.

Adventitia

A system is said to be chaotic in a scientific sense when the initial conditions of that system cannot be defined. A system under these conditions will appear to obey physical and chemical laws during short term observations but, due to this initial uncertainty, the ultimate manifestations of manipulations of the system cannot be predicted with accuracy. The human body is such a system. Under most, if not all, clinical conditions it is impossible to describe with the necessary accuracy the initial state of health of the body prior to the onset of a disease process. Most studies, no matter how rigid the inclusion and exclusion criteria are, must therefore contain a random sampling of patients whose outward appearance may appear similar but whose initial conditions may be very different. The implications of this may be very far reaching.

Cellular and laboratory animal preparations have better potential for establishing more uniform initial biological conditions than clinical human experimentation. As a result it is more likely that these experiments would demonstrate treatment benefits since there are fewer outside influences on these more simple systems to effect outcome – that is, there is less chaos present. This in part may explain the disparate results seen in some cellular and animal models of treatment regimes when applied to human populations. However, they do represent the necessary first steps in establishing efficacy of novel therapies and, in fact, may eventually be used to help establish the minimal requirements necessary for truly matching the initial conditions of subjects undergoing human experimentation.

Chaos presents many difficult challenges for the clinical researcher. Under most circumstances the overt manifestations of a disease are preceded by a non-specific prodrome. At the time a diagnosis is established, therefore, the biological system has undergone numerous interactions which may vary immensely from subject to subject. The threshold for an individual to present for treatment varies considerably, making it difficult consistently to manipulate the system at the same time in the

course of a disease. The consequence of this is to increase the chaos of the biological system. As the system deviates from its resting state in the course of these reactions, it may reach a point where the system can no longer be manipulated by the planned intervention. If enough subjects in a clinical study reach this point the intervention may be perceived from a statistical standpoint to have no benefit – that is, a false negative result.

Earlier identification of a potential subject with the specific clinical entity to be studied, accompanied by the exclusion or statistical separation of patients found later in the course of the disease, should in part reduce the effect of chaos on the outcome of the intervention. Much work is currently being conducted in this area and new earlier markers for ARDS and sepsis are presently being evaluated. Additional work needs to be done to determine how better to match the initial conditions of patients entered into a study. The relative importance of all the potential influences of other organ system deficits on outcomes needs to be determined, such as the patient's nutritional state, recent and remote infectious disease history, and medication history. Finally, the use of empiricism may be required until the above information is available. Under these constraints investigators may be forced to apply a proposed intervention to a large initial group of patients in whom an exhaustive database has been obtained. A detailed statistical evaluation might then demonstrate a subgroup of patients which appeared to have benefited from the intervention. The final step would be to apply the therapy again to this identified subgroup of patients in a randomised, prospective study which should then ultimately answer the question of efficacy. Although this could increase the cost and potentially slow down the process of clinical research, it would result in studies less confounded by chaos. The data from these studies would be less tainted with false positive and false negative results.

ERIC H GLUCK