Short paper

Hypoxaemia and release of endothelin-1

Robert I Cargill, David G Kiely, Roland A Clark, Brian J Lipworth

Abstract

Background – Secretion of the vasoconstrictor peptide endothelin-1 from vascular endothelium is increased by various stimuli. Whether hypoxaemia affects plasma levels of endothelin-1 in humans is unknown, but this may be important in the haemodynamic response to hypoxaemia. The plasma endothelin-1 concentrations in hypoxaemic humans has therefore been measured.

Methods – Plasma levels of endothelin-1 were measured by specific radioimmunoassay in 10 control subjects at rest and following 30 minutes of acute hypoxaemia (SaO₂ 75–80%) induced by breathing a nitrogen/oxygen mixture, and in 10 patients with hypoxaemic cor pulmonale.

Results – The plasma endothelin-1 concentration in control subjects was increased from a mean (SE) of 0.90 (0.11) pmol/l at baseline to 2.34 (0.34) pmol/l during hypoxaemia. In patients with cor pulmonale the plasma endothelin-1 concentration was 2.96 (0.34) pmol/l, raised in comparison with control subjects at rest but similar to levels in controls during hypoxaemia.

Conclusions – Plasma levels of endothelin-1 were increased by hypoxaemia in humans. The raised levels observed in patients with cor pulmonale may largely be attributable to the effects of hypoxaemia, although the pathophysiological significance of these observations remains to be established.

(Thorax 1995;50:1308–1310)

Keywords: endothelin, cor pulmonale, hypoxaemia.

The endothelins are a group of structurally similar peptides synthesised by vascular endothelium. These peptides act via two specific receptors. Stimulation of the type A endothelin receptor (ET_A), the main subtype found on vascular smooth muscle cells,¹ results in profound and longlasting vasoconstriction, whilst stimulation of the type B receptor (ET_B) causes release of vasodilator metabolites and transient vasodilation.² Interestingly, however, in rat pulmonary resistance vessels in vitro ET_B activation may be important in mediating endothelin-1 induced vasoconstriction.³ In humans, endothelin-1 is the most potent vaso-

constrictor substance known through its relatively selective activation of the ET_A receptor.⁴ A role in the maintenance of vascular tone is suspected but this remains speculative at present.

Synthesis of this peptide is increased by a number of humoral and physical stimuli, including hypoxia, which may be pathophysiologically relevant.⁵ In vitro studies have shown that acute hypoxia increases endothelin-1 production by human endothelial cells⁶ whilst, in experimental animals in vivo, hypoxia increases plasma concentrations of endothelin-1.⁷ The effect of hypoxaemia on endothelin-1 in humans is unknown but may be relevant in the cardiovascular adaptations to hypoxaemia and the circulatory abnormalities seen in conditions such as cor pulmonale.

We have therefore studied the effects of hypoxaemia on plasma endothelin-1 levels in healthy subjects and also in patients with hypoxaemic cor pulmonale.

Methods**SUBJECTS***Normal controls*

Ten young male volunteers of mean (SE) 28.1 (2.2) years were studied. None was taking prescribed medication and all had a normal clinical history and examination, 12-lead electrocardiogram, echocardiogram, and haematological and biochemical screen, and forced expiratory volume in one second (FEV₁) of >90% predicted.

Cor pulmonale patients

Ten patients (six men) of mean (SE) age 73.4 (1.7) years with clinically stable cor pulmonale secondary to chronic obstructive pulmonary disease were studied. All had obstructive pattern spirometry (FEV₁/forced vital capacity (FVC) <70%), arterial hypoxaemia while breathing air (PaO₂ <8.0 kPa), and had or gave a history of having peripheral oedema despite normal left ventricular function (on echocardiogram or radionuclide ventriculography), normal renal function (serum creatinine <120 mmol/l), and normal serum albumin (>35 g/l). Patients with other significant cardiovascular disease were excluded.

In this group, FEV₁ in litres was 0.75 (0.06) (range 0.36–1.11), FEV₁ as % of predicted 33.8 (5.2) (range 19–61), PaO₂ on air 6.30

Department of
Clinical
Pharmacology,
Ninewells Hospital and
Medical School,
Dundee DD1 9SY, UK
R I Cargill
D G Kiely
B J Lipworth

Department of
Respiratory Medicine,
King's Cross Hospital,
Dundee DD3 8EA
R A Clark

Reprint requests to:
Dr R I Cargill,
Department of Medicine,
Raigmore Hospital,
Inverness IV2 3UP, UK.

Received 11 May 1995
Returned to authors
4 July 1995
Revised version received
13 July 1995
Accepted for publication
24 July 1995

(0.34) kPa (range 4.35–7.80), and PaCO_2 on air 6.17 (0.50) kPa (range 3.84–8.80).

PROTOCOL

Informed consent was obtained to the protocol previously approved by the Tayside committee for medical research ethics. Baseline venous blood samples were taken into chilled EDTA tubes after 30 minutes supine rest while breathing air. Control subjects only were then rendered hypoxaemic (steady state SaO_2 75–80% measured by pulse oximetry) for 30 minutes by breathing a variable nitrogen/oxygen mixture before taking a further blood sample.

ENDOTHELIN-1 ASSAY

Samples were centrifuged at 2000 rpm at 4°C for 15 minutes and stored at –70°C. Assay was carried out using a commercially available radioimmunoassay kit (Nicholls Institute Diagnostics, San Juan Capistrano, CA, USA) after solid phase extraction on C18 silica columns. The lower limit of detection was 0.4 pmol/l and the intra-assay coefficient of variation was 4.5%. The anti-endothelin antibody used was 100% specific for endothelin-1 and had cross-reactivity of 96% with endothelin-3 and 7% with proendothelin.

DATA ANALYSIS

After testing for normality of distribution, between group comparisons were made by analysis of variance followed by Duncan's multiple range testing with $p < 0.05$ considered significant.⁸ Results are expressed as mean (SE) with 95% confidence intervals (CI) for significant differences.

Levels of endothelin-1 in 10 controls when normoxaemic (○) and following 30 minutes hypoxaemia (●), and in 10 patients with cor pulmonale (△). Boxes indicate sample means with standard error bars and asterisks indicate values significantly ($p < 0.05$) different from normoxaemic controls.

Results

The plasma endothelin-1 concentration in normoxaemic controls was 0.90 (0.11) pmol/l. After 30 minutes hypoxaemia it increased significantly to 2.34 (0.34) pmol/l (95% CI for mean difference 0.41 to 2.48). In patients with cor pulmonale the plasma endothelin-1 concentration was 2.96 (0.34) pmol/l, significantly greater than control subjects when normoxaemic (95% CI for mean difference 1.02 to 3.09) but not when hypoxaemic.

The plasma endothelin-1 concentration therefore increased 2.6-fold in response to hypoxaemia in controls and was raised 3.3-fold in patients with cor pulmonale compared with normoxaemic controls. Results from individual subjects with sample means are depicted in the figure.

Discussion

These findings indicate that hypoxaemia increases plasma levels of endothelin-1 in humans. In patients with cor pulmonale, endothelin-1 was increased to levels comparable to those in normal subjects rendered acutely hypoxaemic. It is interesting to note that baseline levels in controls were similar to those observed in other series,⁹ and that the increase following hypoxaemia and in patients with cor pulmonale is comparable to the rise seen in normal subjects at high altitude.¹⁰

These observations in humans are largely confirmatory of in vitro and animal studies which have shown hypoxia to be a potent stimulus for endothelin-1 synthesis and gene expression.^{6,7} In normal humans Therkelson *et al* found that 15 minutes of hypoxaemia caused only a small increase in plasma endothelin-1 which was not statistically significant, perhaps due to the shorter duration of the stimulus.¹¹ Abnormally high levels of endothelin-1 have been described in a series of patients with pulmonary hypertension of varying aetiology, not all of whom were hypoxaemic,⁹ and thus other stimuli may also be implicated. In the present series, by excluding subjects with other significant cardiovascular diseases such as hypertension and congestive heart failure, we feel hypoxaemia is the most significant stimulus responsible for increased endothelin-1 levels in cor pulmonale. The in vitro evidence would suggest that the increased endothelin-1 levels observed were due to increased synthesis although, as endothelin-1 clearance mechanisms in humans have not been fully characterised,⁵ the possibility of decreased removal cannot be discounted.

The potent vasoactive properties of endothelin-1 may also be responsible for some of the circulatory abnormalities seen during hypoxaemia, although plasma levels may not accurately reflect local concentrations and hence vasoconstrictor activity. Whether endothelin-1 acts as a mediator of acute hypoxic pulmonary vasoconstriction is unknown. Support for this hypothesis might be drawn from studies using endothelin receptor blockers which can attenuate acute hypoxic pulmonary vasoconstriction in rats¹² and prevent development

of pulmonary hypertension following chronic hypoxia.¹³ These drugs now need to be tested in humans where it is likely that endothelin-1 plays a significant part in the cardiovascular response to hypoxaemia.

Manipulation of the endothelin system may therefore be a useful measure in patients with hypoxaemic lung disease, either to prevent or treat the cardiopulmonary consequences of chronic hypoxaemia.

The authors thank Gillian Pirie and Wendy Coutie for performing endothelin assays.

- 1 Davenport AP, O'Reilly G, Kuc RE. Endothelin ET_A and ET_B mRNA and receptors expressed by smooth muscle in the human vasculature: majority of the ET_A sub-type. *Br J Pharmacol* 1995;114:1110-6.
- 2 Bigaud M, Pelton JT. Discrimination between ET_A- and ET_B-receptor-mediated effects of endothelin-1 and [Ala^{1,3,11,15}]endothelin-1 by BQ-123 in the anaesthetised rat. *Br J Pharmacol* 1992;107:912-8.
- 3 MacLean MR, McCulloch KM, Baird M. Endothelin ET_A- and ET_B-receptor-mediated vasoconstriction in rat pulmonary arteries and arterioles. *J Cardiovasc Pharmacol* 1994;23:838-45.
- 4 Arai H, Hori S, Aramori I, Ohkubo H, Nakanishi S. Cloning and expression of a cDNA encoding an endothelin receptor. *Nature (London)* 1990;348:730-2.
- 5 Haynes WG, Webb DJ. The endothelin family of peptides: local hormones with diverse roles in health and disease. *Clin Sci* 1993;84:485-500.
- 6 Kourembanas S, Marsden PA, McQuillan LP, Faller DV. Hypoxia induces endothelin gene expression and secretion in cultured human endothelium. *J Clin Invest* 1991;88:1054-7.
- 7 Shirakami G, Nakao K, Saito Y, Magaibuchi T, Jougaski M, Mukoyama M, *et al.* Acute pulmonary alveolar increases lung and plasma endothelin-1 levels in conscious rats. *Life Sci* 1991;48:969-76.
- 8 Brown RA, Beck JS. Comparison of several groups: the analysis of variance. In: *Medical statistics on microcomputers*. London: BMJ Publishing, 1990:34-42.
- 9 Stewart DJ, Levy RD, Cernacek P, Langleben D. Increased plasma endothelin-1 in pulmonary hypertension: marker or mediator of disease. *Ann Intern Med* 1991;114:464-9.
- 10 Goerre S, Wenk M, Bartsch P, Luscher TF, Niroomand F, Hohenhaus E, *et al.* Endothelin-1 in pulmonary hypertension associated with high altitude exposure. *Circulation* 1995;91:359-64.
- 11 Therkelson K, Jensen KA, Freundlich M, Thorsauge H, Bunemann L, Bøgeskov Nielsen L. Endothelin-1 and cerebral blood flow: influence of hypoxia, hypercapnia and indomethacin on circulating endothelin levels in healthy volunteers. *Scand J Clin Lab Invest* 1994;54:441-51.
- 12 Oparil S, Chen SJ, Meng QC, Elton TS, Yano M, Chen YF. Endothelin receptor antagonist prevents acute hypoxia induced pulmonary hypertension in the rat. *Am J Physiol* 1995;12:L95-100.
- 13 Eddahibi S, Raffestin B, Clozel M, Levame M, Adnot S. Protection from pulmonary hypertension with an orally active endothelin receptor antagonist in hypoxic rats. *Am J Physiol* 1995;37:H828-35.