Short papers

Ca²⁺-independent nitric oxide synthase activity in human lung after cardiopulmonary bypass

René Delgado, Armando Rojas, Luis A Glaría, Mabel Torres, Felix Duarte, Rigoberto Shill, Manuel Nafeh, Enma Santin, Noel González, Miriam Palacios

Abstract

Background – Because surgery involving cardiopulmonary bypass induces a systemic inflammatory response, the effect of cardiopulmonary bypass on nitric oxide (NO) generation was investigated in human lung tissue.

Methods – Nitric oxide synthase (NOS) activity was measured by the conversion of ¹⁴C-L-arginine to ¹⁴C-L-citrulline in tissue biopsy samples obtained before and after cardiopulmonary bypass.

Results – The Ca²⁺-independent production of NO found before cardiopulmonary bypass was extremely low (1.5 (0.5) pmol citrulline/mg/min), but was increased after the bypass operation (23.6 (11) pmol/mg/min).

Conclusions – Ca²⁺-independent NOS activity was detected in human lung after cardiopulmonary bypass. This finding may provide an important insight into the pathogenesis of the tissue damage and acute phase response observed after such surgery.

(Thorax 1995;50:403-404)

Keywords: cardiopulmonary bypass, nitric oxide, nitric oxide synthase.

After open heart operations with cardiopulmonary bypass patients show symptoms resembling an acute phase response such as fever, complement activation, and synthesis of acute phase proteins.¹ Furthermore, increased concentrations of cytokines such as tumour necrosis factor (TNF), interleukin 1 (IL-1), and interleukin 6 (IL-6) have been detected, suggesting that they may be important mediators in this systemic inflammatory response. The participation of oxygen-derived free radicals is also likely.^{1,2}

Nitric oxide (NO) synthesised from L-arginine by the constitutive and Ca²⁺-dependent low output pathway (cNOS) functions as an intracellular and intercellular signal for the stimulation of soluble guanylate cyclase and therefore plays a part in the regulation of cell function and communication.³ On the other hand, NO is also synthesised by a cytokine inducible and Ca²⁺-independent high output

pathway (iNOS) and is an important cytotoxic effector molecule which accounts, at least in part, for the vasodilatation and tissue damage seen in endotoxic shock.⁴

Because several alterations observed in the lung after cardiopulmonary bypass are similar to those seen during haemorrhagic shock and endotoxaemia,⁵ we have examined whether such alterations could be related to the L-arginine:NO pathway.

Methods

All subjects gave their informed consent to the studies which had the approval of the Hermanos Ameijeiras Hospital ethics committee according to the National Medical Ethics Regulations.

Tissue samples from the middle lobe of the lung were taken before (n=3) and after (n=7) cardiopulmonary bypass from patients undergoing open heart operations. None of the patients had previous pulmonary disease. Indications for surgery included mitral (n=4) or aortic (n=3) valve dysfunction. Coronary artery disease was diagnosed in two patients and one had congenital heart disease. Before surgery blood tests showed no evidence of haematological or biochemical disorders and arterial oxygen tensions (P_O₂) were normal (10–13.2 kPa). After cardiopulmonary bypass a significant reduction in haemoglobin and altered blood clotting parameters were detected in all patients. Glucose levels were increased in three patients. The mean (SE) duration of the cardiopulmonary bypass operation in the seven patients was 122 (30) minutes.

Tissue samples were washed twice with sucrose buffered solution (sucrose 320 mM, Hepes 10 mM, ethylenediaminetetraacetate (EDTA) 100 μM, pH 7.4) and homogenised in 1 ml ice cold homogenisation buffer (DL-dithiothreitol 1 mM, Tris 50 mM, sucrose 320 mM, EDTA 1 mM, soybean trypsin inhibitor 10 μg/ml, aprotinin 5 μg/ml, leupeptin 10 μg/ml, pH 7.0). The homogenate was then centrifuged at 100 000g for 20 minutes at 4°C. NOS activity in the supernatant (cytosolic) fractions was measured by the conversion of radiolabelled ¹⁴C-L-arginine to ¹⁴C-L-citrulline in the presence or absence of 1 mM ethylene glycol-bis (β-aminoethyl ether tetraacetic acid) (EGTA)

Center of
Pharmaceutical
Chemistry,
Pharmacology and
Toxicology
Department, PO Box
16042, Havana, Cuba
R Delgado
A Rojas
L A Glaría
M Torres
E Santin
M Palacios

Hermanos Ameijeiras
Hospital,
Cardiosurgery Unit,
Havana, Cuba
F Duarte
R Shill
M Nafeh
N González

Reprint requests to:
Dr R Delgado.

Received 21 March 1994
Returned to authors
20 April 1994
Revised version received
26 August 1994
Accepted for publication
5 December 1994

Ca^{2+} -independent nitric oxide synthase activity in cytosolic fractions from tissue biopsy samples of human lung before and after cardiopulmonary bypass. Closed circles represent paired data before and after cardiopulmonary bypass. Mean values for each group are indicated by bars and are statistically different ($p < 0.05$).

or N^G -monomethyl-L-arginine (L-NMMA) as previously described.⁶

Statistical significance was assessed by the Student's t test. A p value of less than 0.05 was considered significant.

Results

Lung tissue samples from patients after cardiopulmonary bypass showed significant Ca^{2+} -independent NOS activity (23.6 (11) pmol/min/mg) compared with that found before cardiopulmonary bypass (1.5 (0.5), $p < 0.05$, figure).

Discussion

The cellular location of Ca^{2+} -independent NOS (iNOS) remains to be established, but during extracorporeal circulation there is a sequestration of polymorphonuclear cells within the alveolar vasculature⁷ and human neutro-

phils have been shown to express iNOS.⁸ As mentioned above, alveolar changes occurring after cardiopulmonary bypass are similar to those observed in endotoxaemia and haemorrhagic shock. Induction of the NO pathway has been reported in both pathological processes.^{3,4,9}

Endothelial NO production (cNOS) has been implicated in maintaining low pulmonary vascular resistance in physiological circumstances.

Because both isoforms have recently been immunohistochemically localised in human lung,¹⁰ the cNOS/iNOS balance may have important implications in vascular pulmonary homeostasis. It is therefore not clear at present whether the appearance of the Ca^{2+} -independent NOS activity after cardiopulmonary bypass will be damaging or protective. Although iNOS is implicated in tissue and cell injury, it is likely that this activity may modulate both pulmonary vascular hypertension and cardiogenic pulmonary oedema after cardiopulmonary bypass.

- Butler J, Rocker GM, Westaby S. Inflammatory response to cardiopulmonary bypass. *Ann Thorac Surg* 1993;55:552-9.
- Markewitz A, Faist B, Lang S, Endres S, Hultner L, Reichart B. Regulation of acute phase response after cardiopulmonary bypass by immunomodulation. *Ann Thorac Surg* 1993;55:389-94.
- Moncada S, Palmer RMJ, Higgs EA. Nitric oxide: physiology, pathophysiology and pharmacology. *Pharmacol Rev* 1991;43:109-42.
- Kilbourn RG, Jubran A, Gross SS, Griffith OW, Levi R, Adams J, et al. Reversal of endotoxin mediated shock by N^G -methyl-arginine as inhibitor of nitric oxide synthesis. *Biochem Biophys Res Commun* 1990;172:1132-8.
- Packer DJ, Karp RB, Kirkling JW, Bedard P. Lung water and alveolar and capillary volumes after intracardiac surgery. *Circulation* 1972;45(Suppl I):139-46.
- Salter M, Knowles RG, Moncada S. Widespread tissue distribution and changes in activity of Ca^{2+} -independent nitric oxide synthase. *FEBS Lett* 1991;291:145-9.
- Ratcliff NB, Young GW, Hackel DB, Mikat E, Wilson JW. Pulmonary injury secondary to extracorporeal circulation. *J Thorac Cardiovasc Surg* 1973;65:425-32.
- Salvemini D, De Nucci G, Gryglewski RJ, Vane JR. Human neutrophils and mononuclear cells inhibit platelet aggregation by releasing nitric oxide like factor. *Proc Natl Acad Sci USA* 1989;89:6328-32.
- Thiemermann C, Szabo C, Mitchell JA, Vane JR. Vascular hyporeactivity in hemorrhagic shock is mediated by nitric oxide. *Proc Natl Acad Sci USA* 1993;90:267-71.
- Kobzik L, Bredt DS, Lowenstein CJ, Drazen J, Gaston B, Sugarbaker D, et al. Nitric oxide synthase in human and rat lung: immunocytochemical and histochemical localization. *Am J Respir Cell Mol Biol* 1993;9:371-7.