Bronchiectasis and homozygous (P_1ZZ) α_1 -antitrypsin deficiency in a young man

William Rodriguez-Cintron, Kalpalatha Guntupalli, Armando E Fraire

Department of Pulmonary Medicine, Baylor College of Medicine and Ben Taub General Hospital W Rodriguez-Cintron

Department of Medicine, Baylor College of Medicine, and Department of Pulmonary/Critical Care Medicine, Ben Taub General Hospital, Houston, Texas 77030, USA K Guntupalli

Department of Pathology, University of Massachusetts Medical Center, Worcester, Massachusetts, USA A E Fraire

Reprint requests to: Dr K Guntupalli.

Received 20 October 1993
Returned to authors 7 January 1994
Revised version received 31 March 1994
Accepted for publication 5 April 1994

Abstract

Pulmonary complications of α_1 -antitrypsin deficiency are most commonly manifested by panlobular emphysema. This report describes histologically proven bronchiectasis in a 21 year old man with massive haemoptysis and homozygous deficiency of α_1 -antitrypsin. Neither panlobular emphysema nor cirrhosis of the liver were present. Bronchiectasis must be considered part of the spectrum of the pulmonary pathology that may be encountered in individuals with α_1 -antitrypsin deficiency.

(Thorax 1995;50:424-425)

Keywords: α_1 -antitrypsin deficiency, bronchiectasis.

The occurrence of α_1 -antitrypsin deficiency in association with cirrhosis of the liver and pan acinar pulmonary emphysema has been well documented. However, the association of α_1 -antitrypsin deficiency with bronchiectasis or repeated bronchopulmonary infections, although reported, is not well documented. Various reports have provided clinical and bronchographic documentation of α_1 -antitrypsin deficiency in association with bron-

chiectasis, but only two cases to our knowledge have provided histopathological verification of the bronchiectasis. The purpose of this report is (1) to present the clinical and radiographic features of this uncommon association in a young man with massive haemoptysis, a P_1ZZ homozygous phenotype, and histopathologically proven bronchiectasis; and (2) to review the cumulative experience reported thus far in the medical literature.

Case report

In August 1987 a 21 year old non-smoking white man presented to Ben Taub General Hospital in Houston, Texas after coughing two cups of blood. His past medical history included a laparotomy at the age of two for suspected hepatosplenomegaly. However, the results of this exploration were unknown to the patient and his family. He remained well until the age of 16 when he developed frequent episodes of cough, excessive sputum production, and occasional bouts of blood-tinged sputum. At the age of 18 he underwent a thoracotomy for removal of his left lower lobe following an episode of massive haemoptysis. Histopathological examination of the resected left lower lobe showed grossly dilated bronchi. Microscopic evaluation showed several abnormally dilated large bronchi near the pleural surface with chronic inflammation of the bronchial wall. There was also focal destruction and interruption of the bronchial wall musculature and multiple aggregated lymphoid cells, a finding which corresponded microscopically to bronchiectasis with follicular bronchitis (figure).

Physical examination revealed a young man in no acute distress. The pulse was 120/min, respiration rate 20/min, and blood pressure was 130/80 mm Hg. There was no clubbing, cyanosis, or orthostatic hypotension, breath sounds were vesicular, and the heart rhythm was normal. A spleen tip was palpable in the left upper abdominal quadrant, and the liver was non-palpable. There was no peripheral lymphadenopathy. A chest radiograph was unremarkable except for postoperative changes at the left lung base. A fiberoptic bronchoscopy performed two days later showed a moderate amount of dark blood oozing from the bronchial orifice corresponding to the left upper lobe which did not clear after saline lavage. Similar oozing of dark blood was noted in the right upper lobe but cleared readily after saline lavage. The haemoptysis ceased spontaneously. Pulmonary function testing two weeks later showed an FEV₁ of 3.71 (79% predicted), an FVC of 4.69 (80% predicted), and a TLC of 6.95 (90% predicted). The transfer factor for carbon monoxide (TLCO) and liver function tests were normal. A bronchogram with radiopaque contrast material showed saccular bronchiectasis in the lingula and upper lobe, which now occupied the left hemithorax, and mild bronchiectasis of the right upper lobe. Chest computed tomographic scanning confirmed the bronchographic findings and identified a small cirrhotic liver, gastric varices, and

Chronic inflammation (arrow) extending deep into mucous glands and focal destruction with interruption of the muscle layer in the bronchial wall (double arrow). Stain: haematoxylin and eosin, original magnification $\times 240$ reduced to 58% in origination.

Reports in the medical literature of α_1 -antitrypsin deficiency in association with bronchiectasis

Reference	No. of cases	Total P_1ZZ	HPVB
Ericksson (1965) ⁶	35	35	Yes (2)
Orell (1971) ⁷	8	6	Yes (4)
Longstreth (1975) ¹	1	1	No
Kagan (1975) ¹¹	1	1	No
Larsson (1978) ⁸	28	246	No
Scott (1977) ²	1	1	No
Jones (1985) ³	3	3	No
Current case (1994)	1	1	Yes (1)

HPVB = histopathological verification of bronchiectasis.

splenomegaly. However, a percutaneous needle biopsy of the liver showed no microscopic evidence of cirrhosis. Periodic acid Schiff positive granules were not identified within hepatocytes. Serum protein electrophoresis showed the absence of the usual spike seen in the α_1 -globulin region. The serum level of α_1 -antitrypsin was 18 mg/dl (8% of normal). The α_1 -antitrypsin phenotype of the patient's parents and one sibling were all P_1ZZ . Sweat levels of chloride and immunoelectrophoresis were both normal.

Discussion

Although no clear risk for the development of bronchiectasis in homozygous α_1 -antitrypsin deficient individuals has ever been established, a few reports have helped to confirm an association between this deficiency and bronchiectasis and pulmonary infections.¹⁻³ It has been suggested that the absence of this protease inhibitor accentuates the damage due to the unopposed protease activity which occurs in bronchitis and its accompanying neutrophilic infiltration of the bronchial wall.^{3,4} Repeated bronchopulmonary infection with severe P_1ZZ have been reported.⁵ It has been postulated that in the α_1 -antitrypsin-deficient individual repeated episodes of ordinary bronchitis, of whatever cause, may lead to the development of bronchiectasis.⁴ This suggestion seems to be supported by the study by Ericksson⁶ in which most of the 35 cases who underwent necropsy had bronchitis, although only two of the 35 had bronchiectasis.

Orell and Mazodier reviewed the pulmonary histopathology of 38 cases of α_1 -antitrypsin deficiency collected from the literature and added eight cases of their own.⁷ These authors noted that "chronic bronchiectasis was a common but not a constant finding" in the 38 patients in the literature, but they did not mention bronchiectasis specifically in their review of the 38 cases. Of their own eight cases they chose to report their findings in only six, but did not state why two cases were excluded. Interestingly, one of these six cases was contributed by Ericksson, which is acknowledged in the table.⁷ Four of the six cases had bronchiectasis, of whom three had peripheral cylindrical bronchiectasis and one had minor bronchiectasis. These cases differ considerably from the one reported here in that they had

severe panlobular emphysema, often with features of vanishing lung and bronchiectasis as a secondary feature.⁷ In contrast, our case had bronchiectasis but no histopathological evidence of panlobular emphysema.

Our review of the literature suggests that the association of α_1 -antitrypsin deficiency and bronchiectasis may not be as uncommon as is usually believed. Most reported patients with this association have been in middle to late life, usually smokers, and only six appear to have been documented with histopathological evaluation of the bronchiectasis. Thus, our patient may represent only the seventh case reported with histopathological verification of the bronchiectasis, and the first to be diagnosed during life. In agreement with most previous study reports, our patient proved to have a P_1ZZ phenotype.

Studies from Sweden indicate that the frequency of the association of α_1 -antitrypsin deficiency and bronchiectasis may be as high as 10%.⁶⁻⁸ However, the true frequency of bronchiectasis in α_1 -antitrypsin-deficient individuals remains to be determined. Conversely, the frequency of α_1 -antitrypsin deficiency in series of bronchiectasis (from all causes) is also unknown.⁴ Although α_1 -antitrypsin is the major anti-elastase of the normal lower respiratory tract, other anti-elastases also exist.^{9,10} Because of the possible protection by anti-elastases other than α_1 -antitrypsin that are not measured, the association between bronchiectasis and anti-elastase deficiency may not be well established.

The ease with which α_1 -antitrypsin can be aerosolised into the tracheobronchial tree with return of anti-elastase defences may make this a therapeutic option in patients with bronchiectasis attributed to α_1 -antitrypsin deficiency.⁵

1 Longstreth G, Weitzman SA, Browning RJ, Lieberman J. Bronchiectasis and homozygous alpha-1-antitrypsin deficiency. *Chest* 1975;67:233-5.

2 Scott J, Anderson CL, Shankar PS, Staurides A. Alpha-1-antitrypsin deficiency with diffuse bronchiectasis and cirrhosis of the liver. *Chest* 1977;71:535-8.

3 Jones DK, Gooden D, Cavanaugh P. Alpha-1-antitrypsin deficiency presenting as bronchiectasis. *Br J Dis Chest* 1985;79:301-4.

4 Stockley RA. Chronic bronchitis: the antiprotease/protease balance and the effect of infections and corticosteroids. *Clin Chest Med* 1988;9:643-56.

5 Hubbard RC, Brantly ML, Sellers SE, Mitchell ME, Crystal RG. Anti-neutrophil-elastase defenses of the lower respiratory tract in α_1 -antitrypsin deficiency directly augmented with an aerosol of α_1 -antitrypsin. *Ann Intern Med* 1989;111:206-12.

6 Ericksson S. Studies in alpha-1-antitrypsin deficiency. *Acta Med Scand* 1965;177:1-85.

7 Orell SR, Mazodier P. Pathological findings in alpha-1-antitrypsin deficiency. In: Mittmon C, ed. *Pulmonary emphysema and proteolysis*. New York and London: Academic Press, 1971:69-89.

8 Larsson C. Natural history and life expectancy in severe alpha-1-antitrypsin deficiency. *Acta Med Scand* 1978;204:345-51.

9 Gadek JE, Fells GA, Zimmerman RL, Rennard SI, Crystal RG. Antielastases of the human alveolar structures. *J Clin Invest* 1981;68:889-98.

10 Tobin MJ, Cook PJJ, Hutchison DCS. Alpha-1-antitrypsin deficiency: the clinical and physiological features of pulmonary emphysema in subjects homozygous for Pi type Z. *Br J Dis Chest* 1983;77:14-27.

11 Kagan E, Soskolne CL, Zwi S, Hurwitz S, Maier GMG, Ipp T, et al. Immunologic studies in patients with recurrent bronchopulmonary infections. *Am Rev Respir Dis* 1975;111:441-51.