Detection of anti-cytokeratin 8 antibody in the serum of patients with cryptogenic fibrosing alveolitis and pulmonary fibrosis associated with collagen vascular disorders

Naomi Dobashi, Jiro Fujita, Yuji Ohtsuki, Ichiro Yamadori, Takeo Yoshinouchi, Tadashi Kamei, Michiaki Tokuda, Satoko Hojo, Hiroki Okada, Jiro Takahara

Abstract

Background—It has been suggested that the humoral immune system plays a role in the pathogenesis of cryptogenic fibrosing alveolitis (CFA). Although circulating autoantibodies to lung protein(s) have been suggested, none of the lung proteins have been characterised. The purpose of this study was to determine the antigen to which the serum from patients with pulmonary fibrosis reacted.

Methods—The anti-A549 cell antibody was characterised in a patient with CFA using Western immunoblotting and immunohistochemical staining of A549 cells. As we identified that one of the antibodies against A549 cells was anti-cytokeratin 8, the expression of mRNA of cytokeratin 8 in A549 cells was evaluated. In addition, we attempted to establish an enzyme linked immunosorbent assay to measure the levels of anti-cytokeratin 8 antibody in the serum of patients with CFA and pulmonary fibrosis associated with collagen vascular disorders (PF-CVD).

Results—Initially two anti-A549 cell antibodies were detected in the serum of patients with pulmonary fibrosis, one of which was characterised as anti-cytokeratin 8 antibody by Western immunoblotting. We were able to establish an ELISA to measure anti-cytokeratin 8 antibody and found significantly higher levels in patients with CFA and PF-CVD than in normal volunteers, patients with sarcoidosis, pneumonia, and pulmonary emphysema.

Conclusions—One of the anti-A549 cell antibodies in the serum of patients with CFA was against cytokeratin 8. The serum levels of anti-cytokeratin 8 antibody were increased in patients with CFA and PF-CVD. These results suggest that anti-cytokeratin 8 antibody may be involved in the process of lung injury in pulmonary fibrosis.

(Thorax 1998;53:969-974)

Keywords: A549 cell; antibody; cytokeratin 8; idiopathic pulmonary fibrosis; collagen vascular disorders

Cryptogenic fibrosing alveolitis (CFA) and pulmonary fibrosis associated with a collagen vascular disorder (PF-CVD) are inflammatory

lung diseases of unknown aetiology that are characterised by the accumulation of neutrophils and mononuclear cells.^{1,2} Increased numbers of neutrophils, eosinophils, and CD4 positive lymphocytes as well as raised levels of immunoglobulin have been reported in the bronchoalveolar lavage fluid of patients with CFA.^{2,4} This suggests an immunological pathogenesis for the disease although no definite aetiological factors have been indicated.

Wallace *et al* recently reported the presence of circulating autoantibodies in patients with CFA.^{5,6} However, none of the target antigens have been characterised. We have therefore evaluated the autoantibody against A549 cells and have identified one of the antigens to be cytokeratin 8.

Methods

SUBJECTS

Twenty one patients (11 women) of median age 68 years (range 44-86) with a diagnosis of pulmonary fibrosis were studied from 1990 to 1996. Of the 21 patients, 12 had IPF and nine had PF-CVD (four rheumatoid arthritis, four dermatomyositis, and one Sjogren syndrome). There were 13 non-smokers, two ex-smokers, and six current smokers. The diagnoses were made on clinical, radiological, physiological, and histological grounds. The criteria used included history of exertional dyspnoea and cough, fine crackles on physical examination, compatible findings on the chest radiograph (diffuse basal reticulonodular shadowing), physiological abnormalities of restrictive lung defects including decreased transfer factor, and abnormal Pao₂ at rest and/or with exertion. Histological confirmation was obtained in all cases by open lung biopsy. Histological examination of all patients with PF-CVD showed a pattern of usual interstitial pneumonia (UIP). In all patients high resolution computed tomographic scanning of the lungs (HRCT) was performed. None of the patients received immunosuppressive treatment such as corticosteroids or cyclophosphamide. Lung function tests (% vital capacity (VC) and forced expiratory volume in one second (FEV₁)) were performed in all patients. We also studied 15 normal subjects (seven women) of median age 33 years, all of whom were non-smokers and, as a control for different pulmonary diseases, we studied five patients with sarcoidosis (four women) of median age 46 years, all non-smokers, 16 with pneumonia (six women) of

First Department of Internal Medicine, Kagawa Medical University, 1750-1 Miki-cho, Kita-gun, Kagawa 761-0793, Japan

N Dobashi
J Fujita
M Tokuda
S Hojo
H Okada
J Takahara

Department of Respiratory Medicine, Gamagohri City Hospital, Aichi, Japan
T Yoshinouchi

Department of Internal Medicine, Kagawa Prefectural Central Hospital, Kagawa, Japan
T Kamei

Department of Pathology, Okayama University Medical School, Okayama, Japan
I Yamadori

Department of Pathology, Kochi Medical School, Kochi, Japan
Y Ohtsuki

Correspondence to:
Dr J Fujita.

Received 27 February 1998
Returned to authors
27 April 1998
Revised manuscript received
26 May 1998
Accepted for publication
15 June 1998

Figure 1 Western immunoblot analysis using serum from a patient against lysates of several cell lines. Lane 1, HL60; lane 2, K562; lane 3, U937; lane 4, Daudi; lane 5, PC9; lane 6, A549; lane 7, Jurkat; lane 8, TK130; lane 9, TO1019; lane 10, MN321. Although only a single band (arrow **) was detected in HL60, K562, Daudi, Jurkat, TK130, TO1019, and MN321, two bands (arrow * and **) were detected in PC9 and A549.

median age 65 years, six of whom were non-smokers and 10 were smokers, and 10 patients with pulmonary emphysema (all men) of median age 73 years, all of whom were smokers.

The protocols for this study were approved by the institutional review board for human studies and informed written consent was obtained from the subjects.

BLOOD SAMPLES

Peripheral venous blood samples with and without ethylene diamine tetra-acetic acid (EDTA) were obtained before breakfast. After centrifugation at 1000*g* for 10 minutes at 4°C the serum was frozen and stored at -70°C until used. Arterial blood oxygen and carbon dioxide tensions (Pao₂ and Paco₂) were measured with a blood gas analyser. To evaluate the antibodies against several cell lines the serum of a 66 year old female patient with CFA was used.

CELL LINES

The following cell lines were used: A549, PC9 (derived from adenocarcinoma of the lung), TK130, TO1019, MN321 (small cell carcinoma of the lung), HL60 (acute promyelocytic leukaemia), K562 (chronic myelogenous leukaemia), U937 (lymphoma), and Jurkat (leukaemia).

Figure 2 Immunohistochemical staining of A549 cells with anti-cytokeratin 8 monoclonal antibody. The cytoplasm of A549 cells was clearly and strongly stained. The nuclei were completely negative (avidin-biotin peroxidase complex method, ×580).

Table 1 Immunohistochemical staining of A549 cells by several monoclonal antibodies against cytokeratins

Antibodies	Cytokeratins	Results	Molecular weight (kD)
35βH11 (MA902)	CK 8	(+)	52.5
CK18	CK18	(+)	45
CAM5.2	CK 8, CK 18	(+)	52.5, 45
CK14	CK14	(-)	50
CK16	CK16	(-)	48
CK17	CK17	(-)	46
CK19	CK19	(-)	40

SDS-PAGE ELECTROPHORESIS AND WESTERN BLOTTING

SDS-polyacrylamide gel electrophoresis was performed according to Laemmli's method⁷ with a slight modification. Lysates of cell lines were mixed with sodium dodecyl sulphate (SDS 2.0%) and heated (100°C, 5 min). The samples were then applied to a 10/20% SDS polyacrylamide gel, electrophoresed (60 mA, 120 min), fixed in 50% methanol, 10% acetic acid, and stained with Coomassie Blue. Standard molecular weight markers purchased from Daiichi-Kagaku (Tokyo, Japan) comprised egg lysozyme (MW 14 400), trypsin inhibitor (MW 20 100), carbonic anhydrase (MW 30 000), aldolase (MW 42 400), bovine albumin (MW 66 000), and phosphorylase B from rabbit muscle (MW 97 400). Commercially available purified keratin from human epidermis (Sigma Chemical Co, lot 95H11291, St Louis, Missouri, USA) and recombinant human cytokeratin 8 (Progen Biotechnik GmbH, lot 503209, Heidelberg, Germany) were also used as the positive control.

Proteins were electrophoretically transferred onto nitrocellulose membrane⁸ and were detected by immunoblotting using the serum from one patient, peroxidase conjugated goat anti-human IgG antibody (Sigma Immuno-Chemicals, lot 094H-4810, St Louis, Missouri, USA), and stained with 4CN PLUS for chromogenic detection of horse radish peroxidase (NEM Life Science Products, Boston, Massachusetts, USA). Western immunoblots for cytokeratin 8 were also performed using a mouse monoclonal anti-human cytokeratin antibody (CAM5.2, lot 61217, Becton Dickinson Immunocytometry Systems, San Jose, California, USA) which detects human cytokeratin 8 and cytokeratin 18, and gold conjugated anti-mouse IgG antibody (Sigma Chemical Co, lot 98F8882). Silver staining was then performed using Immuno-Blot gold enhancement kit (Bio-RAD Laboratories, Hercules, California, USA). We also evaluated peroxidase conjugated goat anti-human IgA (Sigma ImmunoChemicals, lot 023H-8868) and anti-human IgM (Sigma ImmunoChemicals, lot 094H-8832) as second antibodies.

IMMUNOHISTOCHEMISTRY OF A549 CELLS BY SEVERAL ANTICYTOKERATIN ANTIBODIES

To evaluate the expression of several cytokeratins in A549 cells, immunohistochemical staining by anti-human monoclonal antibodies against several cytokeratins was performed. A549 cells were immunohistochemically stained, employing the avidin-biotin peroxi-

Figure 3 Expression of cytokeratin 8 and β -actin mRNA in several lung cancer cell lines by the RT-PCR method. Lane 1, PC9 (adenocarcinoma); lane 2, A549 (adenocarcinoma); lane 3, TK130 (small cell lung cancer); lane 4, TO1019 (small cell lung cancer); lane 5, MN321 (small cell lung cancer). DNA ladder (100 bp) and X174 Hae III digest were used as DNA size markers. Although all cell lines expressed mRNA for cytokeratin 8 (approximately 244 bp), A549 and PC9 expressed increased amounts of mRNA.

Figure 4 Western immunoblot analysis using the anti-human cytokeratin antibody CAM5.2 which detects cytokeratin 8 and cytokeratin 18 and patient's serum. Lane 1, CAM5.2 antibody against A549; lane 2, patient's serum against A549; lane 3, CAM5.2 antibody against PC9; lane 4, patient's serum against PC9. The positive band with a higher molecular weight (arrow *) corresponded to cytokeratin 8. The positive band with a lower molecular weight was not stained by CAM5.2 antibody, suggesting that this band was not cytokeratin 18.

dase complex method (Dako LSAB kit-peroxidase, Dako Corp, Kyoto, Japan) using mouse monoclonal antibodies to cytokeratins. The monoclonal antibodies used were as follows: 35 β H11 (MA902) which detects cytokeratin 8 (Enzo Diagnostics Inc, New York, USA, 1:5000 dilution), anti-human cytokeratins 18 and 19 (ScyTek Lab, Logan, Utah, USA, each 1:30 dilution), CAM5.2

which detects both cytokeratin 8 and 18 (Becton Dickinson Immunocytometry Systems, San Jose, California, USA, 1:20 dilution), anti-human cytokeratins 14, 16 and 17 (Novo Casstra Lab Ltd, Newcastle upon Tyne, UK, 1:20, 1:30 and 1:30 dilution, respectively). In order to retrieve and increase the immunoreactivities, cytokeratins 14, 17, 18, CAM5.2 and MA902 were preincubated with 0.1% pronase at 37°C for 20 minutes. Cytokeratins 16 and 19 were autoclaved at 132°C for 12 minutes.

EXPRESSION OF MRNA FOR CYTOKERATIN 8 IN A549 CELLS BY RT-PCR

To confirm the expression of cytokeratin 8 mRNA from A549 cells, reverse transcriptase-polymerase chain reaction (RT-PCR) was performed. Total RNA from each cell line was isolated by a single step acid guanidium-phenol chloroform extraction. Total RNA (1 μ g) was reverse transcribed into cDNA using random primers (Takara Shuzo Co, Kyoto, Japan) and avian myeloblastosis virus reverse transcriptase (Life Science Inc, Petersburg, Florida, USA) in a final volume of 20 μ l. The cDNA was amplified by PCR. The primer sequences for cytokeratin 8 were as follows: sense primer AACAAACCTTAGGCGGCAGCT, antisense primer GCCTGAGGAAGTTGATCTCG. PCR beads (Ready-To-Go, Amersham Pharmacia Biotech Ltd, Uppsala, Sweden) containing 50 pmol of each primer, 1 μ l of cDNA, and 22 μ l of water were used. Thirty cycles were performed consisting of a 60 second denaturation at 95°C and annealing and polymerase extension for two minutes at 72°C. All PCRs were terminated with a 10 minute extension at 72°C. Ten μ l aliquots of the PCR products were electrophoresed on 10/20% SDS polyacrylamide gel and analysed by direct visualisation after ethidium bromide staining. The expected size of the amplified fragment was 244 nucleotides.

ENZYME-LINKED IMMUNOSORBENT ASSAY FOR ANTI-CYTOKERATIN 8 AUTOANTIBODY IN HUMAN SERUM

An enzyme-linked immunosorbent assay (ELISA) was established to measure anti-cytokeratin 8 antibody levels in human serum samples. Serum was added to wells coated with recombinant human cytokeratin 8 (0.5, 1, 2, and 4 μ g/ml). After incubation and washing, the solid phase bound anti-human cytokeratin 8 autoantibody was further incubated with peroxidase conjugated goat anti-human IgG antibody (Sigma ImmunoChemicals, lot 094H-4810, diluted 1:1000). After further washes TMB Peroxidase EIA Substrate Kit (Bio-RAD Laboratories, Hercules, California, USA) was used to measure the amount of solid phase bound antibodies. The assay was calibrated using a standard solution of the patient's serum in which the anti-cytokeratin 8 antibody had been determined by Western immunoblotting. Data were expressed as mean values from duplicate determinations.

Figure 5 Western immunoblot analysis using the patient's serum against A549 (lane 1), PC9 (lane 2), and keratin from human epidermis (lane 3). The patient's serum strongly reacted with the keratin from human epidermis (arrow *). The positive band with a lower molecular weight (arrow **) did not correspond to the keratin from human epidermis.

Figure 6 Western immunoblot analysis using the patient's serum against A549 (lane 1), PC9 (lane 2), and recombinant human cytokeratin 8 (lane 3). The patient's serum reacted strongly only with recombinant human cytokeratin 8.

Figure 7 Standard curves of ELISA revealing anti-cytokeratin 8 antibody in patient's serum. Numbers represent concentrations of recombinant cytokeratin 8 ($\mu\text{g/ml}$) to coat wells.

STATISTICAL METHODS

The results were expressed as mean (SE) values. Comparisons of values between groups were analysed with the Mann-Whitney U test, a p value of less than 0.05 being considered as significant.

Results

Figure 1 shows the Western immunoblot analysis of one patient's serum against lysates of several cell lines. Although only one band (MW 45 000) was detected in HL60 cells, K562 cells, Daudi cells, Jurkat cells, TK130, and TO1019, two bands (MW 52 000 and 45 000) were found in PC9 and A549 cells. There were no positive reactions when anti-human IgA and IgM antibodies were used (data not shown).

Immunohistochemical staining of A549 cells with anti-cytokeratin 8 monoclonal antibody was restricted to the cytoplasm, and the nuclei were completely negative (fig 2). Table 1 summarises the results of immunohistochemical staining of A549 cells. The cytoplasm of A549 cells was also stained with anti-cytokeratin 18 monoclonal antibody. However, A549 cells were negative with anti-cytokeratin 14, 16, and 17 monoclonal antibodies. There were a few positive cells in the cytoplasm with anti-cytokeratin 19 antibody.

Figure 3 shows the expression of cytokeratin 8 mRNA in several lung cancer cell lines by RT-PCR. Although all cell lines expressed mRNA for cytokeratin 8, A549 and PC9 cells expressed increased amounts of mRNA.

Western immunoblot analysis using a patient's serum and anti-human cytokeratin (CAM5.2) which detects cytokeratin 8 and cytokeratin 18 against A549 cells and PC9 cells showed that the patient's serum and anti-human cytokeratin reacted with A549 cells and PC9 cells (fig 4). The positive band which had a higher molecular weight (approximately 52 000) corresponded to cytokeratin 8. Western immunoblot analysis using a patient's serum against A549 cells, PC9 cells, and human keratin showed that the patient's serum

Figure 8 Levels of anti-human cytokeratin 8 antibodies in serum samples from patients and from normal volunteers. Mean (SE) levels of anti-human cytokeratin antibodies in the serum of patients with pulmonary fibrosis (0.48 (0.10)) were significantly higher than those of normal volunteers (0.1 (0.004), $p < 0.001$, Mann-Whitney U test) and of patients with sarcoidosis (0.08 (0.03), $p < 0.01$), pneumonia (0.09 (0.02), $p < 0.001$), and pulmonary emphysema (0.11 (0.02), $p < 0.001$).

reacted strongly with the human keratin (fig 5). Western immunoblot analysis using a patient's serum against A549 cells, PC9 cells, and recombinant human cytokeratin 8 showed that the patient's serum reacted strongly with recombinant human cytokeratin 8 (fig 6).

Figure 7 shows standard curves of ELISA which detect anti-cytokeratin antibody in the patient's serum. Using this ELISA it was possible to measure the anti-human cytokeratin antibody in the patients' serum samples. Mean (SE) levels of anti-human cytokeratin antibodies in serum from patients with pulmonary fibrosis (0.48 (0.10) compared with the control patient's serum which was determined to 1.0) were significantly higher than those of normal volunteers (0.1 (0.004), $p < 0.001$; fig 8). In addition, levels of anti-human cytokeratin antibodies in serum samples from patients with pulmonary fibrosis were significantly higher than those in patients with sarcoidosis (0.08 (0.03), $p < 0.01$), pneumonia (0.09 (0.02), $p < 0.001$), and pulmonary emphysema (0.11 (0.02), $p < 0.001$). In patients with pulmonary fibrosis there was no significant difference between levels of anti-human cytokeratin antibodies in the serum of eight smokers (six current smokers and two ex-smokers, 0.404 (0.102)) and 13 non-smokers (0.524 (0.164)).

Discussion

In this study we have demonstrated that one of the antibodies against A549 cells in patients with CFA and PF-CVD is anti-cytokeratin 8 antibody. Measurement of this antibody by ELISA showed that the levels in serum from patients with CFA and PF-CVD were significantly higher than levels in serum from normal volunteers and patients with other pulmonary diseases.

The current consensus is that, whatever the aetiological agent(s), CFA represents a persistent immunological reaction in the lung which

results in scar formation.⁹ Data from bronchoalveolar lavage have shown increased numbers of neutrophils, eosinophils, and CD4 positive lymphocytes³ as well as raised levels of immunoglobulin⁴ in the bronchoalveolar lavage (BAL) fluid of patients with CFA. In addition, there is evidence of a more general immune dysregulation with hypergammaglobulinaemia,³ circulating immune complexes,^{10, 11} and the production of recognised non-organ specific autoantibodies such as antinuclear factor and rheumatoid factor.¹² Clinical studies have shown that collagen vascular disorders, which have a known immunological pathogenesis, are associated with an identical pattern of alveolitis,¹³ and it has also been reported that there may be large numbers of B lymphocytes and plasma cells in the lungs of patients with CFA.¹⁴ Furthermore, it has been reported that the B cell growth factor activity¹⁵ and immunoglobulin⁴ in BAL fluid increases in patients with CFA compared with normal controls.

The presence of autoantibodies in patients with CFA has been implied and subsequently investigated.^{5, 6, 12, 16} The nature and location of the antigen(s) are believed to be associated with pulmonary epithelial lining cells and, perhaps, with capillary endothelial cells on the basis of ultrastructural studies¹⁷ and a previous demonstration of immune complexes at these sites.¹⁸

It has been reported that the serum from patients with CFA is positive on cryostat sections of lung tissues by an indirect immunofluorescence technique.¹⁶ Other investigators using ELISA techniques have found antibodies to the nuclear antigen topoisomerase II (170 kD) in 38% of patients¹⁹ and hepatitis C virus in 28% of patients.²⁰ Wallace *et al* have reported that circulating IgG autoantibodies to 70–90 kD protein(s) which are associated with alveolar epithelial lining cells are detected in serum from patients with CFA.^{5, 6} These findings suggest that there are several pulmonary autoantigens to which autoantibodies in the serum react.

We have shown that anti-cytokeratin 8 antibodies are present in serum samples from patients with CFA and PF-CVD. Cytokeratin is antigenically related to keratin and α -helical coiled-coil protein.²¹ Some studies have shown that antibodies can penetrate cells *in vivo*.^{22–25} Antibodies that bind to cytokeratin 8 may therefore be pathologically significant. Defects in cytoskeletal organisation are known to be capable of disrupting cell function²⁶ and may be characteristic of some pathological processes.²⁷ It has been reported that antibodies to cytoskeletal proteins are associated with some organ-specific autoimmune diseases.²¹ Cytokeratin-specific antibodies appear to increase in diseases such as alcoholic liver disease and rheumatic disorders,^{28, 29} and it has been suggested that anticytokeratin antibodies are formed in response to epithelial cell injury or death.^{30–33} However, none of these reports has described the autoantibody as specific to cytokeratin 8.

Few studies have evaluated the clinical significance of anti-cytokeratin 8 autoantibody in serum.³⁰ There appears to be a high incidence of antibodies against cytoskeletal proteins in normal serum which suggests that these antibodies

may be a part of the natural/physiological autoantibody repertoire.³⁰ However, since physiological autoantibodies are usually IgM,^{34,35} the existence of the IgG type anti-cytokeratin 8 antibody in the serum may have been pathogenetically relevant to pulmonary fibrosis.

The significance of anti-cytokeratin 8 autoantibody in the pathogenesis of CFA and PF-CVD should be discussed. In the lung cytokeratin 8 is expressed in bronchoepithelial cells.³⁶ Cytokeratin 8 may have been released from epithelial cells following cell injury. The resulting antibody-antigen interaction with immune complex formation could have a significant role in the perpetuation of the disease process, either by direct injury of epithelial cells or via local macrophage activation as they are cleared by phagocytosis.

In addition, it has been shown that wild type p53 is overexpressed in regenerated bronchoepithelial cells in patients with CFA.^{37,38} It has also been reported that the expression of the human cytokeratin 8 gene is regulated by the p53 gene.³⁹ It is therefore possible that regenerated bronchoepithelial cells that had overexpressed p53 protein might have been selectively destroyed by anti-cytokeratin 8 antibody. However, we cannot rule out the possibility that the existence of cytokeratin 8 autoantibody is a non-specific consequence of lung injury. The significance of anti-cytokeratin 8 antibody in terms of disease severity, stage, prognosis, and possible response to treatment should therefore be evaluated in future studies.

In this study the serum of patients with CFA reacted strongly with two antigens of molecular weights 52 000 and 45 000. Although we could characterise one protein with a molecular weight of 52 000 as cytokeratin 8, the other antigen still needs to be identified.

In conclusion, our results have shown that one of the anti-A549 cell antibodies in the serum of patients with CFA was anti-cytokeratin 8. In addition, levels of anti-cytokeratin 8 antibodies in the serum increased in patients with pulmonary fibrosis including PF-CVD. These results suggest that anti-cytokeratin 8 antibodies may play a significant role in the process of lung injury in pulmonary fibrosis.

- 1 Crystal RG, Bitterman PB, Rennard SI, et al. Interstitial lung diseases of unknown cause. Disorders characterized by chronic inflammation of the lower respiratory tract. *N Engl J Med* 1984;310:154-66.
- 2 Peterson MW, Monick M, Hunninghake GW. Prognostic role of eosinophils in pulmonary fibrosis. *Chest* 1987;92:51-6.
- 3 Cherniack RM. Bronchoalveolar lavage constituents in healthy individuals, interstitial pulmonary fibrosis and selected comparison groups. *Am Rev Respir Dis* 1990;141(Suppl):S169-93.
- 4 Reynolds HY, Fulmer JD, Kazmierowski JA, et al. Analysis of cellular and protein content of bronchoalveolar lavage fluid from patients with idiopathic pulmonary fibrosis and chronic hypersensitivity pneumonitis. *J Clin Invest* 1977;59:165-75.
- 5 Wallace WAH, Roberts SN, Caldwell H, et al. Circulating antibodies to lung protein(s) in patients with cryptogenic fibrosing alveolitis. *Thorax* 1994;49:218-24.
- 6 Wallace WAH, Schofield JA, Lamb D, et al. Localization of a pulmonary autoantigen in cryptogenic fibrosing alveolitis. *Thorax* 1994;49:1139-45.
- 7 Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680.
- 8 Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of polypeptides from polyacrylamide gels to nitrocellulose sheets. *Proc Nat Acad Sci USA* 1979;76:4350-4.
- 9 Kradin RL, Divertie MB, Colvin RB, et al. Usual interstitial pneumonitis is a T cell alveolitis. *Clin Immunol Immunopathol* 1986;40:224-35.
- 10 Haslam PL, Thomson B, Mohammed I, et al. Circulating immune complexes in patients with cryptogenic fibrosing alveolitis. *Clin Exp Immunol* 1979;37:381-90.
- 11 Dreisen RB, Schwarz MI, Theofilopoulos AN, et al. Circulating immune complexes in the idiopathic interstitial pneumonias. *N Engl J Med* 1978;298:353-7.
- 12 Turner- Warwick M, Doniach D. Auto-antibodies in idiopathic pulmonary fibrosis. *BMJ* 1965;1:886-91.
- 13 Wallaert B. Subclinical alveolitis in immunologic systemic disorders. *Lung* 1990;168(Suppl):974-83.
- 14 Haslam PL. Evaluation of alveolitis by studies of lung biopsies. *Lung* 1990;168(Suppl):984-92.
- 15 Emura M, Nagi S, Takeuchi M, et al. In vitro production of B cell growth factors and B cell differentiation factors by peripheral blood mononuclear cells and bronchoalveolar lavage T lymphocytes from patients with idiopathic pulmonary fibrosis. *Clin Exp Immunol* 1990;82:133-9.
- 16 Ehringer R. Similar frequency of auto-antibodies against type II pneumocytes and Clara cells in patients with interstitial lung disease and healthy persons. *Klin Wochenschr* 1991;69:297-302.
- 17 Corrin B, Dewar A, Rodriguez-Roisin R, et al. Fine structural changes in cryptogenic fibrosing alveolitis and asbestosis. *J Pathol* 1985;147:107-19.
- 18 Schwartz MI, Dreisen RB, Pratt DS, et al. Immunofluorescent pattern in idiopathic interstitial pneumonia. *J Lab Clin Med* 1978;91:929-35.
- 19 Meliconi R, Bestagno M, Sturani C, et al. Autoantibodies to DNA topoisomerase II in cryptogenic fibrosing alveolitis and connective tissue diseases. *Clin Exp Immunol* 1989;76:184-9.
- 20 Ohta K, Watanabe J, Hirai K, et al. High prevalence serum antibodies to hepatitis C virus in idiopathic pulmonary fibrosis. *Am Rev Respir Dis* 1992;145:A218.
- 21 Kurki P, Virtanen I. The detection of human antibodies against cytoskeletal components. *J Immunol Methods* 1984;67:209-23.
- 22 Alarcon-Segovia D, Ruiz-Arguelles A, Fishbein E. Antibody to nuclear ribonucleoprotein penetrates live human mononuclear cells through Fc receptors. *Nature* 1978;271:67-9.
- 23 Alarcon-Segovia D, Ruiz-Arguelles A, Fishbein E. Antibody penetration into living cells. I. Intracellular immunoglobulin in peripheral blood mononuclear cells in mixed connective tissue disease and systemic lupus erythematosus. *Clin Exp Immunol* 1979;35:364-75.
- 24 Alarcon-Segovia D, Ruiz-Arguelles A, Fishbein E. Antibody penetration into living cells. II. antiribonucleoprotein IgG penetrates into Tc lymphocytes causing their deletion and the abrogation of suppressor function. *J Immunol* 1979;122:1855-62.
- 25 Alarcon-Segovia D. Penetration of antinuclear antibodies into immunoregulatory T cells: pathogenetic role in the connective tissue diseases. *Clin Immunol Allergy* 1981;1:117-26.
- 26 Sturgess JM, Chao J, Turner JAP. Transposition of ciliary microtubules: another cause of impaired ciliary motility. *N Engl J Med* 1980;303:318-22.
- 27 Runger-Brandt E, Gabbiani G. The role of cytoskeletal and cytocontractile element in pathologic processes. *Am J Pathol* 1983;110:361-92.
- 28 Kurki P, Miettinen A, Salaspuro M, et al. Cytoskeleton antibodies in chronic active hepatitis, primary biliary cirrhosis, and alcoholic liver disease. *Hepatology* 1983;3:297-302.
- 29 Kurki P, Helve T, Virtanen I. Antibodies to cytoplasmic intermediate filaments in rheumatic diseases. *J Rheumatol* 1983;10:558-62.
- 30 Kurki P, Virtanen I. The detection of human antibodies against cytoskeletal components. *J Immunol Methods* 1984;67:209-23.
- 31 Zauli D, Crespi C, Dall'Amore P, et al. Antibodies to the cytoskeleton components and other autoantibodies in inflammatory bowel disease. *Digestion* 1985;32:140-4.
- 32 Grubauer G, Romani N, Kofler H, et al. Apoptotic keratin bodies as autoantigen causing the production of IgM-anti-keratin intermediate filament autoantibodies. *J Invest Dermatol* 1986;87:466-71.
- 33 Hintner H, Romani N, Stanzl U, et al. Phagocytosis of keratin filament aggregates following opsonization with IgG-anti-keratin filament autoantibodies. *J Invest Dermatol* 1987;88:176-82.
- 34 Tron F, Le Guern C, Cazenave PA, Bach JF. Intrastain recurrent idiotypes among anti-DNA antibodies of (NZB x NZW) F1 hybrid mice. *Eur J Immunol* 1982;12:761-6.
- 35 Tron F, Jacob L, Bach JF. Murine monoclonal anti-DNA antibodies with an absolute specificity for DNA have a large amount of idiotype diversity. *Proc Nat Acad Sci USA* 1983;80:6024-7.
- 36 Iyonaga K, Miyafuji M, Suga M, et al. Alterations in cytokeratin expression by the alveolar lining epithelial cells in lung tissue from patients with idiopathic pulmonary fibrosis. *J Pathol* 1997;182:217-24.
- 37 Kuwano K, Kunitake R, Kawasaki M, et al. p21Waf1/Cip1/Sd1 and p53 expression in association with DNA strand breaks in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 1996;154:477-83.
- 38 Hojo S, Fujita J, Yamadori I, et al. Overexpression of p53 protein in interstitial lung diseases. *Respir Med* 1998;92:184-90.
- 39 Mukhopadhyay T, Roth JA. p53 involvement in activation of the cytokeratin 8 gene in tumor cell lines. *Anticancer Res* 1996;16:105-12.