

Airway hyperresponsiveness in asthma: not just a matter of airway inflammation

Vito Brusasco, Emanuele Crimi, Riccardo Pellegrino

According to the most recent definition, bronchial asthma is a chronic inflammatory disorder of the airways associated with reversible airway obstruction and increased airway responsiveness to a variety of stimuli.¹ An intuitive inference from this definition is that a causal relationship may exist between airway inflammation and airway hyperresponsiveness. Along this line of reasoning, most of the research in the last two decades in this field was aimed at identifying inflammatory cell products possibly responsible for the pathogenesis of bronchial asthma and airway hyperresponsiveness.² However, the common observation that the asthmatic airways are equally hyperresponsive to a variety of different stimuli does challenge the idea that a single inflammatory cell or mediator may be central to the pathogenesis of asthma and airway hyperresponsiveness and focuses on the importance of an altered mechanical response of the target organ. This view has been recently corroborated by the finding that airway responsiveness of normal individuals may become similar to that of asthmatics by simply changing the pattern of breathing during the bronchial challenge.³ It is therefore legitimate to wonder how much of airway hyperresponsiveness is due to inflammation or to inherent predisposing factors.

The aims of the present review are to show that airway narrowing in asthma is the ultimate result of an interaction between complex and multiple mechanisms not necessarily and uniquely related to airway inflammation, and to revisit the evidence on which the theorem "airway inflammation equal to airway hyperresponsiveness" has been constructed.

Mechanisms of airway narrowing in asthma

Airway calibre *in vivo* is the result of a delicate balance between the force generated by the airway smooth muscle (ASM) and a number of opposing factors.⁴ The latter are mainly represented by autonomic mechanisms that tend to limit the ASM tone and by mechanical forces that oppose ASM shortening.

AIRWAY SMOOTH MUSCLE (ASM)

Although it is widely accepted that ASM contraction is a major cause for airway obstruction in asthma, there is uncertainty as to whether abnormalities of ASM play any part in the pathogenesis of airway hyperresponsiveness. In terms of force-length characteristics, the asthmatic ASM does not behave differently from the normal ASM.⁵⁻⁶ These data lend support to a purely immunological view of bronchial asthma in which the ASM cell is regarded as a normal effector of response to abnormal inflammatory stimuli. Nevertheless, the smooth muscle of

asthmatic airways may be able to generate greater force than normal because of hypertrophy and hyperplasia,⁷ thus causing excessive airway narrowing.⁸⁻⁹ Although the thickness of the ASM in the airway wall of asthmatic patients might have been overestimated¹⁰ and there is no proof that an increased mass of smooth muscle is associated with an increased force generation capacity,¹¹ the possibility that an increased contractile force developed by ASM might contribute to airway hyperresponsiveness in asthma cannot be ruled out.

Recent work has suggested that normal and asthmatic ASM may differ in terms of length shortening characteristics. The capacity and the velocity of shortening are increased after passive sensitisation of ASM.¹² The increase of shortening velocity is due to an increase in the amount and the activity of the myosin light chain kinase¹³ and increased phosphorylation of the 20 kD regulatory unit with increased cross-bridge cycling rate.¹⁴⁻¹⁷ Albeit small, the difference in shortening velocity between asthmatic and normal ASM may be relevant to the generation and maintenance of force. Under normal conditions stretching causes a transient reduction in ASM tone.¹⁸⁻¹⁹ This is considered an important mechanism for preventing excessive airway narrowing (see below) but it may be less efficient if the velocity of shortening is increased.²⁰

Moreover, sensitised ASM may develop a myogenic contractile response to stretching.²¹ This response, which is believed to be a consequence of conversion from multi-unit to single-unit ASM,²² has been invoked to explain the sustained bronchoconstriction that occurs in some asthmatics after taking one or more deep breaths.²³⁻²⁴

MECHANICAL AND GEOMETRIC FACTORS

The mechanical properties of ASM are such that, if unimpeded, it may shorten to about 20-30% of its initial length when an appropriate stimulus is applied.²⁵ *In vivo*, such a shortening would result in complete airway closure.⁸ In normal humans the maximal response to bronchoconstrictor stimuli is limited,²⁶ which suggests that some mechanisms opposing ASM shortening are operative *in vivo*.

In vitro the shortening of ASM is considerably less when elastic loads are applied.²⁷ Furthermore, human ASM shortens less than ASM from other species and this difference seems to be related to a greater amount of connective tissue present in human bronchi, which represents a parallel elastic load.²⁸ It has been suggested that a decrease in the elastic load internal to ASM may contribute to greater ASM shortening in asthma.²⁹ The cartilaginous rings and

Cattedra di
Fisiopatologia
Respiratoria,
Dipartimento di
Scienze Motorie e
Riabilitative,
Università di Genova,
Genova, Italy
V Brusasco
E Crimi

Servizio di
Fisiopatologia
Respiratoria, Ospedale
S. Croce e Carle,
Cuneo, Italy
R Pellegrino

Correspondence to:
Professor V Brusasco,
Dipartimento di Scienze
Motorie e Riabilitative
Facoltà di Medicina e
Chirurgia, Università di
Genova, Largo R Benzi 10,
16132 Genova, Italy.

plates may also limit ASM shortening, thus preventing excessive airway narrowing^{8, 28}; in rabbits, softening the tracheal cartilage by papain consistently increases airway resistance.³⁰

The most efficient force opposing ASM shortening in vivo is provided by the elastic recoil of the lung.^{8, 28} When this is reduced, as in emphysema, the airways can become more narrow more easily.³¹ Although there is no consistent proof that lung elastic recoil is reduced in asthma,^{32–35} even small decrements of transpulmonary pressure near functional residual capacity may have a deleterious effect when the ASM contracts. In normal individuals the response to methacholine is greatly enhanced by breathing just 500 ml below FRC—that is, at a lung volume at which transpulmonary pressure is only a few cm H₂O less.³⁶

The load imposed on the airways by the surrounding lung parenchyma changes during breathing. Due to their visco-plasto-elastic properties, both lung parenchyma and airways dissipate energy during breathing.^{18, 19, 37, 38} For example, the force necessary to elongate the ASM in vitro is slightly but consistently greater than that necessary to bring it back to its original length.^{18, 19, 38} Such an energy dissipation becomes as much as 15 times greater when the ASM is contracted, depending on the frequency and amplitude of the length oscillations.³⁸ Basically, energy dissipation implies less ASM force and, by inference, less bronchoconstriction. During inspiration the airways are stretched by the force of interdependence through which they are coupled with lung parenchyma. If this force is effectively applied to the external airway wall, then ASM shortening may be opposed. Increasing tidal volume therefore reduces the response to bronchoconstrictor stimuli.^{39, 40} When a stretch of similar amplitude of a deep inspiration to total lung capacity is applied, ASM is fully elongated and most of the energy generated during active shortening is dissipated.^{38, 39} In normal individuals a deep inspiration is able to reverse a bronchoconstriction fully, enough to cause as much as a 50% decrease in forced expiratory flow.^{32, 33, 41–44} In asthmatic subjects deep inspiration has less bronchodilator effect on constricted airways^{32, 33, 42, 44–46} or may even cause bronchoconstriction.^{46, 47} The importance of the ability of deep inspiration to dilate the airways has recently been explored by Skloot *et al*³ and by Pellegrino *et al*.³² In normal individuals prevention of deep inspirations during a bronchial challenge causes dyspnoea and the airway responsiveness becomes similar to that of asthmatics.^{3, 32} An inference from these data may be that airway hyperresponsiveness is more a problem of inability to dilate constricted airways rather than of increased constrictor stimuli to ASM.

In a simple model where airway narrowing is uniquely due to ASM shortening, the contractile elements may return to their relaxed status after being stretched because the actin-myosin cross bridges are detached. Then, the airways remain dilated after a deep inspiration until cross bridges reform and the tone prior to the

deep inspiration is re-established.³⁹ This is what seems to occur in normal individuals during induced bronchoconstriction.^{41–44} To understand why the bronchodilator response to deep inspiration in asthmatic subjects is blunted or even reversed, a more complex model must be invoked where ASM does not dissipate energy during cycling and/or it is prevented from doing so by external factors. For the first condition to be true, primary defects of the ASM that prevent detachment of cross bridges must be present. To date there are no data to support such a mechanism, which suggests that the behaviour of the ASM is more likely to be regulated by external forces. In asthma the airways may be less sensitive to the action of external forces because the force of interdependence is reduced^{3, 4, 33, 36, 41, 46, 47} or the non-contractile elements in the airway wall are more stiff.¹⁷ According to Fredberg *et al*¹⁷ the cross bridge cycling rate decreases if the ASM stretching is impeded, and the conversion to slow cycling latch bridges occurs which promotes maintenance of steady state tone and increased stiffness of the ASM. Thus, even a minimal transient unloading of the ASM due to the effects of deep inspiration on the non-contractile tissues external to the ASM—for example, the lung or parenchyma—could result in passive shortening of the ASM. This could explain, at least in part, the blunted bronchodilator effect of deep inspirations associated with the increase in parenchymal hysteresis in asthma.⁴⁶ Furthermore, if the velocity of shortening of the ASM in asthma is increased,^{12–17} the active force may be quickly re-established after inspiration so that the ASM is rapidly unloaded upon expiration.²⁰ Recently, Gunst *et al*⁴⁸ advanced the hypothesis that the force developed by the ASM in response to constrictor stimuli depends on the length at which the stimulus is applied. If the ASM contracts at longer length and then is shortened, the tensile force is less than if the ASM is contracted at shorter length. It has been suggested that this behaviour reflects a different arrangement of the contractile elements inside the ASM cell.⁴⁰

The lack of protection from excessive airway narrowing observed in asthma can be, at least in part, reasonably attributed to airway inflammation or inflammatory remodelling of the airway walls.^{2, 49, 50} It has been suggested that peribronchial oedema may attenuate the pulling effect of parenchymal attachments on airway walls, thus unloading the ASM and allowing more shortening for the same intensity of stimulus.⁴ The airway wall thickening that occurs in bronchial asthma^{49, 51} due to oedema, cellular infiltration, and vascular engorgement may amplify the bronchoconstrictor response as it causes airway calibre to decrease more for a given ASM shortening.⁵² However, the effect of mucosal thickening on airway narrowing is complex. Airway mucosa folds when ASM shortens,^{9, 53} a process that requires energy dissipation and hence may represent a load for the ASM. In addition, the subepithelial fibrosis characteristic of asthma makes the airway stiff, thus opposing ASM shortening.⁵⁴ On the other hand, the degree of airway narrowing is

inversely proportional to the number of mucosal folds,⁵³ which decreases with mucosal and basement membrane thickening.⁵⁵ Paradoxically, the same inflammatory abnormalities could either favour or limit airway narrowing, depending on interactions with other concomitant changes.

In conclusion, owing to the complexity and multiple interactions of the mechanism regulating airway narrowing, it would be simplistic to regard airway inflammation as the only cause of airway hyperresponsiveness.

Airway inflammation and airway hyperresponsiveness

The concept of asthma as an inflammatory disorder derives from necroscopic studies on asthmatic subjects who died because of an asthma attack or other causes^{51 56 57} and in vivo studies using bronchoalveolar lavage, induced sputum, and bronchial biopsy specimens in subjects with asthma of different severity, either at baseline⁵⁸⁻⁶⁰ or after exposure to allergens or occupational sensitisers.⁶¹⁻⁶³

The most prominent feature in necropsies of patients with fatal asthma is a marked thickening of airway walls. This is commonly associated with epithelial damage, thickening of basement membrane, marked increase of bronchial capillary bed, fluid exudation with oedema, goblet cell hyperplasia, ASM hyperplasia and hypertrophy, and intraluminal mucus and cellular debris causing complete or partial airway occlusion. The eosinophil is generally the dominating inflammatory cell in both fatal and non-fatal asthma^{64 65} but neutrophils may prevail in cases of sudden onset fatal asthma.⁶⁴ The airway walls of subjects with non-fatal asthma are also thicker than normal but to a lesser extent than those of subjects with fatal asthma.^{66 67}

Common findings in the biopsy specimens of asthmatic airways are epithelial shedding, collagen deposition below the basement membrane,^{68 69} increased numbers of eosinophils and mast cells in the mucosa. Eosinophils, mast cells and their products are also increased in the bronchoalveolar lavage fluid.⁷⁰ In this scenario the recruitment and activation of Th2 cells seem to play a central role.⁷¹

AIRWAY INFLAMMATION AS A CAUSE OF AIRWAY HYPERRESPONSIVENESS: PROS

In asthmatic subjects inhalation of allergen or occupational sensitisers causes an inflammatory (mainly eosinophilic) response in the airways which is associated with an increase in airway hyperresponsiveness.^{61 72} In both healthy humans and animals exposure to ozone causes an inflammatory (mainly neutrophilic) response and airway hyperresponsiveness.^{73 74} A temporal relationship exists between the inflammatory events occurring in the airways and the increase in airway responsiveness after exposure to allergens. Both the influx of eosinophils and the increase in airway responsiveness precede the development of the late phase response.^{75 76}

Airways hyperresponsiveness often first appears or worsens following a viral infection of

the upper airways or vaccination with live influenza virus. The increase in airways responsiveness after rhinovirus infection is associated with an increase in the number of eosinophils in the sputum.⁷⁷ Airways hyperresponsiveness in these models may be the result of an increased reflex bronchoconstriction via cholinergic and non-cholinergic excitatory neural pathways. Epithelial necrosis, reduced production of epithelial relaxing factors such as PGE₂ and neutral-peptidases, or loosening of tight junctions with overexposure of sensory nerve endings caused by viral replication are the putative underlying mechanisms.⁷⁸ In animal models infection with parainfluenza 3 virus blocks the ability of β_2 adrenoceptor agonists to inhibit antigen induced contraction of isolated ASM,⁷⁹ reduces the activity of muscarinic M₂ autoreceptors,⁸⁰ and increases the number and the releasability of bronchial mast cells.⁸¹ The last two effects have been found to be strictly associated with the development of sustained airways hyperresponsiveness. A virus induced overproduction of interferon gamma (IFN γ) from macrophages and lymphocytes has been indicated as the responsible mechanism.⁸²

AIRWAY INFLAMMATION AS A CAUSE OF AIRWAY HYPERRESPONSIVENESS: CONS

Airway inflammation can be observed both in asthmatic and in atopic non-asthmatic subjects.^{83 84} In subjects with allergic rhinitis who have never experienced asthma symptoms both local⁸⁵ and inhalational⁸⁶ allergen challenge cause an eosinophilic inflammatory response which cannot be easily distinguished from that seen in asthmatic subjects, but the functional responses are different.⁸⁶

There is no evidence to suggest that neutrophils play a role in airway hyperresponsiveness in humans.⁷⁰ In animal models airway neutrophilia induced by sulphur dioxide⁸⁷ or endotoxin⁸⁸ is associated with hyporesponsiveness rather than hyperresponsiveness.

The presence of high IgE levels in infancy seems to be a more important predisposing factor for the development of airway hyperresponsiveness than the occurrence of common viral infections in early life.⁸⁹

Epithelial damage after vaccination with live influenza virus occurs both in healthy and asthmatic subjects but airway hyperresponsiveness develops only in the latter,⁹⁰ suggesting that individual related factors make an important difference.

The results of studies on the relationships between airway hyperresponsiveness and airway inflammation at baseline are largely inconsistent (table 1). In some instances weak correlations were found between one or more of the inflammatory markers (mainly eosinophils, T lymphocytes and epithelial shedding) and the degree of airway hyperresponsiveness. However, spurious correlations may have resulted from the inclusion of non-asthmatic controls in the regression analysis. Moreover, in an almost equal number of studies no significant correlations were found between airway inflammation and airway hyperresponsiveness. In particular, it was found that airway hyperresponsiveness

Table 1 Studies in human subjects showing (+) and not showing (-) significant correlations between airway inflammation and airway hyperresponsiveness

Authors	Year	Assessment			
		BAL	BB	Sputum	Marker(s)
Kirby <i>et al</i> ¹	1987	+			Eos
Wardlaw <i>et al</i> ²	1988	+			Eos, MBP
Kelly <i>et al</i> ³	1988	-			Eos
Chan Yeung <i>et al</i> ⁴	1988	-			Eos
Pliss <i>et al</i> ⁵	1989	+			Eos
Jeffery <i>et al</i> ⁶	1989		-		Eos
Ferguson <i>et al</i> ⁷	1989	+			Eos
Gibson <i>et al</i> ⁸	1989			-	Eos
Djukanovic <i>et al</i> ⁹	1990		-		EG2
Brusasco <i>et al</i> ¹²	1990	-			Eos
Foresi <i>et al</i> ¹⁰	1990	-	-		Eos
Adelroth <i>et al</i> ^{101*}	1990	-			Eos, ECP
Bradley <i>et al</i> ⁶⁰	1991		+		EG2
Walker <i>et al</i> ¹⁰²	1991	+			Eos
Bentley <i>et al</i> ¹⁰³	1992		+		EG2
Ollerenshaw <i>et al</i> ¹⁰⁴	1992		-		Eos
Ferguson <i>et al</i> ¹⁰⁵	1992	±			Eos, ECP
Pin <i>et al</i> ¹⁰⁶	1993			+	Eos
Duddridge <i>et al</i> ^{107*}	1993	-			Eos
Iredale <i>et al</i> ¹⁰⁸	1994			-	Eos
Woolley <i>et al</i> ¹⁰⁹	1996	+	+		EG2, Eos
Chetta <i>et al</i> ¹¹⁰	1996		+		Eos
Kidney <i>et al</i> ¹¹¹	1996			-	Eos
Pizzichini <i>et al</i> ¹¹²	1996		+		Eos, MBP
Foresi <i>et al</i> ¹¹³	1997			+	EG2
Crimi <i>et al</i> ¹¹⁴	1998	-	-	-	Eos, ECP

BAL = bronchoalveolar lavage; BB = bronchial biopsy; Eos = number or percentage of eosinophils; ECP = eosinophil cationic protein; MBP = major basic protein; EG2 = cells stained with anti-ECP antibodies.

*With anti-inflammatory treatment.

may be present even in the absence of demonstrable inflammatory cells in the airway lumen or mucosa,¹¹⁴ which suggests that the presence of inflammatory cells in the airways is not necessary to sustain airway hyperresponsiveness.

In lung transplant recipients airway hyperresponsiveness was reported without airway inflammation,¹¹⁵ suggesting that an altered neural control of ASM tone may play a major role in these subjects.

OPEN QUESTIONS

Owing to the lack of evidence that airway hyperresponsiveness and airway inflammation are closely related, there are several questions that must be answered before a causal relationship between airway inflammation and airway hyperresponsiveness in asthma can be established or rejected.




Figure 1 Hypothetical mechanisms and pathways of airway hyper-responsiveness.

Firstly, can airway hyperresponsiveness develop independent of airway inflammation—for example, as a consequence of inherited abnormalities of ASM contractility or autonomic regulation? This hypothesis cannot be ruled out even if the gaussian distribution of airway hyperresponsiveness in the general population¹¹⁶ and its partial reversibility after pharmacological treatments¹¹⁷ or allergen avoidance¹¹⁸ seem to suggest a major role for acquired rather than for inherited factors.

Secondly, is a specific type of inflammation responsible for airway hyperresponsiveness? At present the most appealing hypothesis is that airway hyperresponsiveness is the consequence of repeated episodes of airway inflammation in susceptible subjects. In this connection, allergic inflammation seems to play a major part as atopy with high serum IgE levels is associated with an increased risk of airway hyperresponsiveness in humans.¹¹⁹ In bronchial biopsy specimens of asthmatic subjects the Th2 phenotype has been found to be associated with airway hyperresponsiveness.¹²⁰ Moreover, in some animal models exposure to the sensitising agent causes an inflammatory response characterised by Th2 lymphocyte phenotype, IL-5 and eosinophil activation,¹²¹ which is followed by the development of airway hyperresponsiveness.¹²² In a murine model, however, airway hyperresponsiveness was obtained even without eosinophil infiltration and was prevented by treatment with anti-IFN γ antibodies, suggesting that eosinophil recruitment and airway responsiveness are differently regulated.¹²³

Thirdly, in which way may airway inflammation lead to the development of airway hyperresponsiveness? One possibility is that the release of mediators (histamine, leukotrienes, PAF, proteases) changes ASM contractility or autonomic regulation. Another possibility is that airway inflammation leads to airway remodelling through the release of chemokines and cytokines. Eotaxin, RANTES, MCP-3, MCP-4, IL-5 and IL-8 promote recruitment, activation and survival of inflammatory cells. IL-4, TGF- β , GM-CSF, TNF and IL-1 β modify other aspects of airway biophysiology as antigen processing (Th1–Th2 switch), nitric oxide synthesis, β_2 adrenoceptor function and the homeostasis of epithelial cells, fibroblasts, myofibroblasts, and also ASM cells. The former mechanism would be responsible for transient hyperresponsiveness upon exposure to triggers, the latter for sustained baseline hyperresponsiveness.¹²⁴

Fourthly, are there inherited factors that modulate the consequences of inflammation on airway structure? This possibility is suggested by epidemiological data¹²⁵ indicating that a child with an allergic parent has an increased risk of developing the same allergic disease as affects the parent, whereas the risk of developing a different allergic disease is not significantly different from that for the general population.

Conclusions

Airway inflammation and airway hyperresponsiveness, two major characteristics of bronchial asthma, are loosely related to each other. It seems that the presence of inflammatory cells in the airways is neither sufficient nor necessary for the development of airway hyperresponsiveness. This would imply that an altered response of the target organ is a prerequisite for airway hyperresponsiveness to develop. In this scenario, chronic airway inflammation is likely to play a key role as a stimulus for structural changes (airway wall remodelling, changes in airway to lung interdependence, changes in ASM contractility) affecting the organ response to acute stimuli (fig 1). A practical conclusion is that no inferences about airway hyperresponsiveness can be made from measurements of airway inflammation and vice versa.

- 1 Guidelines for the diagnosis and management of asthma. National Heart, Lung, and Blood Institute. National Asthma Education Program Expert Panel Report. *J Allergy Clin Immunol* 1991;88:425–34.
- 2 Holgate ST. Mediators and cytokine mechanisms in asthma. *Thorax* 1993;48:103–9.
- 3 Skloot G, Permutt S, Togiag AG. Airway hyperresponsiveness in asthma: a problem of limited smooth muscle relaxation with inspiration. *J Clin Invest* 1995;96:2393–403.
- 4 Macklem PT. A theoretical analysis of the effect of airway smooth muscle load on airway narrowing. *Am Rev Respir Dis* 1996;153:83–9.
- 5 Black JL. Pharmacology of airway smooth muscle in chronic obstructive pulmonary disease and in asthma. *Am Rev Respir Dis* 1991;143:1177–81.
- 6 Vincent KS, Black JL, Yan K, et al. Comparison of in vivo and in vitro responses to histamine in human airways. *Am Rev Respir Dis* 1983;128:875–9.
- 7 Ebina M, Yaegashi H, Takahashi T, et al. Distribution of smooth muscles along the bronchial tree. A morphometric study of ordinary autopsy lungs. *Am Rev Respir Dis* 1990;141:1322–6.
- 8 Moreno RH, Hogg JC, Paré PD. Mechanics of airway narrowing. *Am Rev Respir Dis* 1985;133:1171–80.
- 9 Lambert RK, Wiggs BR, Kuwano K, et al. Functional significance of increased airway smooth muscle in asthma and COPD. *J Appl Physiol* 1993;74:2771–81.
- 10 Opazo-Sae A, Okazawa M, Kay T, et al. Airway smooth muscle (ASM) orientation and airway stiffness favour airway narrowing rather than airway shortening. *Am J Respir Crit Care Med* 1994;149:A583.
- 11 Coflesky JT, Jones RC, Reid LM, et al. Mechanical properties and structure of isolated pulmonary arteries remodeled by chronic hyperoxia. *Am Rev Respir Dis* 1987;136:388–94.
- 12 Mitchell RW, Ruhlmann H, Magnussen H, et al. Passive sensitization of human bronchi augments smooth muscle shortening velocity and capacity. *Am J Physiol (Lung Cell Mol Physiol)* 1994;267:218–22.
- 13 Jiang H, Rao K, Halayko AJ, et al. Bronchial smooth muscle mechanics of a canine model of allergic airway hyperresponsiveness. *J Appl Physiol* 1992;72:39–45.
- 14 Kamm KF, Stull JT. Myosin phosphorylation, force, and maximal shortening velocity in neurally stimulated tracheal smooth muscle. *Am J Physiol* 1985;249:C238–47.
- 15 Murphy RA. What is special about smooth muscle? The significance of covalent crossbridge regulation. *FASEB J* 1994;8:311–8.
- 16 Jiang H, Rao K, Halayko AJ, et al. Ragweed sensitization-induced increase of myosin light chain kinase content in canine airway smooth muscle. *Am J Respir Cell Mol Biol* 1992;7:567–73.
- 17 Fredberg JJ, Jones KA, Nathan M, et al. Friction in airway smooth muscle: mechanism, latch and implications in asthma. *J Appl Physiol* 1996;81:2703–12.
- 18 Ludwig MS, Robatto FM, Simard S, et al. Lung tissue resistance during contractile stimulation: structural damping decomposition. *J Appl Physiol* 1992;72:1332–7.
- 19 Gunst SJ, Mitzner W. Mechanical properties of contracted canine bronchial segments in vitro. *J Appl Physiol* 1981;50:1236–47.
- 20 Solway J, Fredberg JJ. Perhaps airway smooth muscle dysfunction contributes to asthmatic bronchial hyperresponsiveness after all. *Am J Respir Cell Mol Biol* 1997;17:144–6.
- 21 Antonissen LA, Mitchell RW, Kroeger EA, et al. Mechanical alterations of airway smooth muscle in a canine asthmatic model. *J Appl Physiol* 1979;46:681–7.
- 22 Stephens NL, Kroeger EA, Kromer U. Induction of a myogenic response in tonic airway smooth muscle by tetraethylammonium. *Am J Physiol* 1975;228:628–32.
- 23 Marthan R, Woolcock AJ. Is a myogenic response involved in deep inspiration-induced bronchoconstriction in asthmatics? *Am Rev Respir Dis* 1989;140:1354–8.
- 24 Pellegrino R, Violante B, Crimi E, et al. Time course and calcium dependence of sustained bronchoconstriction induced by deep inhalation in asthma. *Am Rev Respir Dis* 1991;144:1262–6.
- 25 Stephens NL, Kroeger EA. Ultrastructure, biophysics, and biochemistry of airway smooth muscle. In: Nadel JA, ed. *Physiology and pharmacology of the airways*. New York: Marcel Dekker, 1980: 31–121.
- 26 Woolcock AJ, Salome CM, Yan K. The shape of the dose-response curve to histamine in asthmatic and normal subjects. *Am Rev Respir Dis* 1984;130:71–5.
- 27 Okazawa M, Bai TR, Wiggs BR, et al. Airway smooth muscle shortening in excised canine lung lobes. *J Appl Physiol* 1993;74:1613–21.
- 28 Ishida K, Paré PD, Blogg T, et al. Effects of elastic loading on porcine trachealis muscle mechanics. *J Appl Physiol* 1990;69:1033–9.
- 29 Bramley AM, Thomson RJ, Roberts CR, et al. Hypothesis: excessive bronchoconstriction in asthma is due to decreased airway elastance. *Eur Respir J* 1994;7:337–41.
- 30 Moreno RH, Paré PD. Intravenous papain-induced cartilage softening decreases preload of tracheal smooth muscle. *J Appl Physiol* 1989;66:1694–8.
- 31 Cheung D, Schot R, Zwinderman AH, et al. Relationship between loss in parenchymal elastic recoil pressure and maximal airway narrowing in subjects with alpha-1-antitrypsin deficiency. *Am J Respir Crit Care Med* 1997;155:135–40.
- 32 Pellegrino R, Wilson O, Jenouri G, et al. Lung mechanics during bronchoconstriction. *J Appl Physiol* 1996;81:964–75.
- 33 Brusasco V, Pellegrino R, Violante B, et al. Relationship between quasi-static pulmonary hysteresis and maximal airway narrowing in humans. *J Appl Physiol* 1992;72:2075–80.
- 34 Mansell A, Dubrawsky C, Levison H, et al. Lung mechanics in antigen-induced asthma. *J Appl Physiol* 1974;37:297–301.
- 35 Freedman S, Tattersfield AE, Pride NB. Changes in lung mechanics during asthma induced by exercise. *J Appl Physiol* 1975;38:974–82.
- 36 Ding DJ, Martin JG, Macklem PT. Effects of lung volume on maximal methacholine-induced bronchoconstriction in normal humans. *J Appl Physiol* 1987;62:1324–30.
- 37 Von Neergaard K. Neue Auffassungen über einen Grundbegriff der Atemmechanik. Die Retraktionskraft der Lunge, abhängig von der Oberflächenspannung in den Alveolen. *Z Gesamte Exp Med* 1929;66:373–94.
- 38 Sasaki H, Hoppin FG Jr. Hysteresis of contracted airway smooth muscle. *J Appl Physiol* 1979;47:1251–62.
- 39 Gunst SJ. Contractile force of canine airway smooth muscle during cyclical length changes. *J Appl Physiol* 1983;55:759–69.
- 40 Shen X, Wu MF, Tepper RS, et al. Mechanisms for the mechanical response of airway smooth muscle to length oscillations. *J Appl Physiol* 1997;83:731–8.
- 41 Pellegrino R, Violante B, Crimi E, et al. Effects of aerosol methacholine and histamine on airways and lung parenchyma in healthy humans. *J Appl Physiol* 1993;74:2681–6.
- 42 Parham JW, Shepard RH, Norman PS, et al. Analysis of time course and magnitude of lung inflation effects on airway tone: relation to airway reactivity. *Am Rev Respir Dis* 1983;128:240–5.
- 43 Kariya ST, Thompson LM, Ingenito EP, et al. Effects of lung volume, volume history, and methacholine on lung tissue viscoelasticity. *J Appl Physiol* 1989;66:977–82.
- 44 Fish JE, Rosenthal RR, Batra G, et al. Airway responses to methacholine in allergic and nonallergic subjects. *Am Rev Respir Dis* 1976;113:579–86.
- 45 Wheatley JR, Paré PD, Engel LA. Reversibility of induced bronchoconstriction by deep inspiration in asthmatics and normal subjects. *Eur Respir J* 1989;2:331–9.
- 46 Burns CB, Taylor WR, Ingram RH Jr. Effects of deep inhalation in asthma: relative airway and parenchymal hysteresis. *J Appl Physiol* 1985;59:1590–6.
- 47 Lim TK, Pride NB, Ingram RH. Effects of volume history during spontaneous and acutely induced air-flow obstruction in asthma. *Am Rev Respir Dis* 1987;135:591–6.
- 48 Gunst SJ, Meiss RA, Wu MF, et al. Mechanisms for mechanical plasticity of tracheal smooth muscle. *Am J Physiol (Cell Physiol)* 1995;268:C1267–76.
- 49 Jeffery PK. Comparative morphology of the airways in asthma and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1994;150:S6–S13.
- 50 O'Byrne PM. Airway inflammation and airway hyperresponsiveness. *Chest* 1986;90:575–7.
- 51 Dunill MS. The pathology of asthma with special reference to changes in the bronchial mucosa. *J Clin Pathol* 1960;13:27–33.
- 52 James AL, Paré PD, Hogg JC. The mechanics of airway narrowing in asthma. *Am Rev Respir Dis* 1989;139:242–6.
- 53 Lambert RK. Role of bronchial basement membrane in airway collapse. *J Appl Physiol* 1991;71:666–73.
- 54 Okazawa M, Vedal S, Verburgt L, et al. Determinants of airway smooth muscle shortening in excised canine lobes. *J Appl Physiol* 1995;78:608–14.
- 55 Wiggs BR, Hrousis CA, Drazen JM, et al. On the mechanism of mucosal folding in normal and asthmatic airways. *J Appl Physiol* 1997;83:1814–21.
- 56 Messer JW, Peters GA, Bennet WA. Causes of death and pathologic findings in 304 cases of bronchial asthma. *Dis Chest* 1960;38:612–24.
- 57 Ebina M, Takahashi T, Chiba T, et al. Cellular hypertrophy and hyperplasia of airway smooth muscle underlying bron-

- chial asthma. A 3-D morphometric study. *Am Rev Respir Dis* 1993;148:720-6.
- 58 Kirby JG, Hargreave FE, Gleich GJ, et al. Bronchoalveolar cell profiles of asthmatic and non-asthmatic subjects. *Am Rev Respir Dis* 1987;136:379-83.
- 59 Gibson PG, Mattoli S, Sears MS, et al. Cellular characteristics of sputum from patients with asthma and chronic bronchitis. *Thorax* 1989;44:693-9.
- 60 Bradley BL, Azzawi M, Jacobson M, et al. Eosinophils, T-lymphocytes, mast cells, neutrophils, and macrophages in bronchial biopsy specimens from atopic subjects with asthma: comparison with biopsy specimens from atopic subjects without asthma and normal control subjects and relationship to bronchial hyperresponsiveness. *J Allergy Clin Immunol* 1991;88:661-74.
- 61 De Monchy JGR, Kauffman HF, Venge P, et al. Bronchoalveolar eosinophilia during allergen-induced late asthmatic reaction. *Am Rev Respir Dis* 1985;131:373-6.
- 62 Rossi GA, Crimi E, Lantero S, et al. The late-phase asthmatic reaction to inhaled allergen is associated with an early recruitment of eosinophils in the airways. *Am Rev Respir Dis* 1991;143:379-83.
- 63 Lam S, LeRiche J, Phillips D, et al. Cellular and protein changes in bronchoalveolar lavage fluid after late asthmatic reactions in patients with red cedar asthma. *J Allergy Clin Immunol* 1987;80:44-50.
- 64 Sanjiv S, Crotty TB, Kephart GM, et al. Sudden-onset fatal asthma. *Am Rev Respir Dis* 1993;148:713-9.
- 65 Bousquet J, Chanez P, Lacoste JY, et al. Eosinophilic inflammation in asthma. *N Engl J Med* 1990;323:1033-9.
- 66 Carrol N, Elliot J, Morton A, et al. The structure of large and small airways in nonfatal and fatal asthma. *Am Rev Respir Dis* 1993;147:405-10.
- 67 Kuwano K, Bosken CH, Wiggs BR, et al. Small airways dimensions in asthma and in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1993;148:1220-5.
- 68 Jeffrey PK, Wardlaw AJ, Nelson FC, et al. Bronchial biopsies in asthma. An ultrastructural, quantitative study and correlation with hyperreactivity. *Am Rev Respir Dis* 1989;140:1745-53.
- 69 Roche WR, Beasley R, Williams JH, et al. Subepithelial fibrosis in the bronchi of asthmatics. *Lancet* 1989;522-9.
- 70 Djukanovic R, Roche WR, Wilson JW, et al. Mucosal inflammation in asthma. *Am Rev Respir Dis* 1990;142:434-57.
- 71 Robinson DS, Hamid Q, Ying S, et al. Predominant Th2-lyke bronchoalveolar T-lymphocyte population in atopic asthma. *N Engl J Med* 1992;326:298-304.
- 72 Brusasco V, Crimi E, Gianiorio P, et al. Allergen-induced increase of airway responsiveness and inflammation in mild asthma. *J Appl Physiol* 1990;69:2209-14.
- 73 Selzer J, Bigby BG, Stulberg M, et al. O₃-induced change in bronchial reactivity to methacholine and airway inflammation in humans. *J Appl Physiol* 1986;60:1321-6.
- 74 Fabbri LM, Aizawa H, Alpert SE. Airway hyperresponsiveness and changes in bronchoalveolar lavage after ozone exposure in dogs. *Am Rev Respir Dis* 1984;129:288-91.
- 75 Durham SR, Craddock CF, Cookson WO, et al. Increases of airway responsiveness to histamine precede allergen-induced late asthmatic responses. *J Allergy Clin Immunol* 1988;82:764-70.
- 76 Rossi GA, Crimi E, Lantero S, et al. The late-phase asthmatic reaction to inhaled allergen is associated with an early recruitment of eosinophils in the airways. *Am Rev Respir Dis* 1991;143:379-83.
- 77 Grunberg K, Smits HH, Timmers MC, et al. Experimental rhinovirus 16 infection. Effects on cell differentials and soluble markers in sputum in asthmatic subjects. *Am J Respir Crit Care Med* 1997;158:609-16.
- 78 Kava T. Acute respiratory infections, influenza vaccination and airway reactivity in asthma. *Eur J Respir Dis* 1987;70(Suppl 150):7-38.
- 79 Buckner CK, Clayton DE, Ain-Shoka AA, et al. Parainfluenza 3 infection blocks the ability of a beta adrenergic receptor agonist to inhibit antigen-induced contraction of guinea pig isolated airway smooth muscle. *J Clin Invest* 1981;67:376-84.
- 80 Fryer AD, Jacoby DB. Parainfluenza virus infection damages inhibitory M₂ muscarinic receptors on pulmonary parasympathetic nerves in the guinea pig. *Br J Pharmacol* 1991;102:267-71.
- 81 Castleman WLR, Sorkness RL, Lemanske RF, et al. Viral bronchiolitis during early life induces increased numbers of bronchiolar mast cells and airway hyperresponsiveness. *Am J Pathol* 1990;137:821-31.
- 82 Busse WW. Viral infections in human. *Am J Respir Crit Care Med* 1995;151:1675-7.
- 83 Djukanovic R, Lai CKW, Wilson JW, et al. Bronchial mucosal manifestations of atopy: a comparison of markers of inflammation between atopic asthmatics, atopic non asthmatics and healthy controls. *Eur Respir J* 1992;5:538-44.
- 84 McFadden ER Jr, Gilbert IA. Asthma. *N Engl J Med* 1992;327:1928-37.
- 85 Sedgwick JB, Calhoun WJ, Gleich GJ, et al. Immediate and late airway response of allergic rhinitis patients to segmental antigen challenge. *Am Rev Respir Dis* 1991;144:1274-81.
- 86 Brusasco V, Oddera S, Mereu C, et al. Airway inflammation and responsiveness to allergen in rhinitis and asthma. *Am J Respir Crit Care Med* 1997;155:A978.
- 87 Shore SA, Karriya ST, Anderson K, et al. Sulfur-dioxide-induced bronchitis in dogs: effects on airway responsiveness to inhaled and intravenously administered methacholine. *Am Rev Respir Dis* 1987;135:840-7.
- 88 Folkerts G, Henricks PAJ, Slootweg PJ, et al. Endotoxin-induced inflammation and injury of the guinea pig respiratory airways cause bronchial hyporeactivity. *Am Rev Respir Dis* 1988;137:1441-8.
- 89 Martinez DF. Viral infections and the development of asthma. *Am J Respir Crit Care Med* 1995;151:1644-8.
- 90 Kerrebijn KF, Howell JBL, Jongejan RC, et al. Airway hyperresponsiveness and inflammation. In: Holgate ST, ed. *The role of inflammatory processes in airway hyperresponsiveness*. Oxford: Blackwells, 1989: 10-13.
- 91 Kirby JG, Hargreave FE, Gleich GJ, et al. Bronchoalveolar cell profiles of asthmatic and non-asthmatic subjects. *Am Rev Respir Dis* 1987;136:379-83.
- 92 Wardlaw AJ, Dunette S, Gleich GJ, et al. Eosinophils and mast cells in bronchoalveolar lavage in subjects with mild asthma. Relationship to bronchial hyperreactivity. *Am Rev Respir Dis* 1988;137:62-9.
- 93 Kelly C, Ward C, Stenton CS, et al. Number and activity of inflammatory cells in bronchoalveolar lavage fluid in asthma and their relation to airway responsiveness. *Thorax* 1988;43:684-92.
- 94 Chan-Yeung M, Leriche J, et al. Comparison of cellular and protein changes in bronchial lavage fluid of symptomatic and asymptomatic patients with red cedar asthma on follow-up examination. *Clin Allergy* 1988;18:359-65.
- 95 Pliss LB, Ingenito EP, Ingram RH. Responsiveness, inflammation, and effects of deep breaths on obstruction in mild asthma. *J Appl Physiol* 1989;66:2298-304.
- 96 Jeffery PK, Wardlaw AJ, Nelson FC, et al. Bronchial biopsy in asthma. *Am Rev Respir Dis* 1989;140:1745-53.
- 97 Ferguson AC, Wong FWM. Bronchial hyperresponsiveness in asthmatic children. Correlation with macrophages and eosinophils in bronchoalveolar fluid. *Chest* 1989;96:988-91.
- 98 Gibson PG, Mattoli S, Sears MS, et al. Cellular characteristics of sputum from patients with asthma and chronic bronchitis. *Thorax* 1989;44:693-9.
- 99 Djukanovic R, Wilson JW, Britten KM, et al. Quantitation of mast cells and eosinophils in the bronchial mucosa of symptomatic atopic asthmatics and healthy control subjects using immunohistochemistry. *Am Rev Respir Dis* 1990;142:863-71.
- 100 Foresi A, Bertorelli G, Pesci A, et al. Inflammatory markers in bronchoalveolar lavage and in bronchial biopsy in asthma during remission. *Chest* 1990;98:528-35.
- 101 Adelroth E, Rosenhall L, Johansson S, et al. Inflammatory cells and eosinophilic activity in asthmatics investigated by bronchoalveolar lavage. *Am Rev Respir Dis* 1990;142:91-9.
- 102 Walker C, Kaegi MK, Braun P, et al. Activated T cells and eosinophilia in bronchoalveolar lavages from subjects with asthma correlated with disease severity. *J Allergy Clin Immunol* 1991;88:935-42.
- 103 Bentley AM, Menz G, Storz CHR, et al. Identification of T lymphocytes, macrophages, and activated eosinophils in the bronchial mucosa in intrinsic asthma. *Am Rev Respir Dis* 1992;142:500-6.
- 104 Ollerenshaw SL, Woolcock AJ. Characteristics of the inflammation in biopsies from large airways of subjects with asthma and subjects with chronic airflow limitation. *Am Rev Respir Dis* 1992;145:922-7.
- 105 Ferguson AC, Whitelaw M, Brown H. Correlation of bronchial eosinophils and mast cell activation with bronchial hyperresponsiveness in children with asthma. *J Allergy Clin Immunol* 1992;90:609-13.
- 106 Pin I, Radford S, Kolendowicz R, et al. Airway inflammation in symptomatic and asymptomatic children with methacholine hyperresponsiveness. *Eur Respir J* 1993;6:1249-56.
- 107 Duddridge M, Ward C, Hendrick DJ, et al. Changes in bronchoalveolar lavage inflammatory cells in asthmatic patients treated with high dose inhaled beclomethasone dipropionate. *Eur Respir J* 1993;6:489-97.
- 108 Iredale MJ, Wanklin SAR, Phillips IP, et al. Non invasive assessment of bronchial inflammation in asthma: no correlation between eosinophilia of induced sputum and bronchial responsiveness to inhaled hypertonic saline. *Clin Exp Allergy* 1994;24:940-5.
- 109 Woolley KL, Adelroth E, Woolley MJ, et al. Granulocyte-macrophage colony-stimulating factor, eosinophils and eosinophils cationic protein in subjects with and without mild, stable, atopic asthma. *Eur Respir J* 1994;7:1576-84.
- 110 Chetta A, Foresi A, Del Donno M, et al. Bronchial responsiveness to distilled water and methacholine and its relationship to inflammation and remodeling of the airways in asthma. *Am J Respir Crit Care Med* 1996;153:910-7.
- 111 Kidney JC, Wong AC, Efthimiadis A, et al. Elevated B cells in sputum of asthmatics. Close correlation with eosinophils. *Am J Respir Crit Care Med* 1996;153:540-4.
- 112 Pizzichini E, Pizzichini MMM, Efthimiadis A, et al. Indices of airway inflammation in induced sputum: reproducibility and validity of cell and fluid-phase measurements. *Am J Respir Crit Care Med* 1996;154:308-17.
- 113 Foresi A, Leone C, Pelucchi A, et al. Eosinophils, mast cells, and basophils in induced sputum from patients with seasonal allergic rhinitis and perennial asthma: relationship to methacholine responsiveness. *J Allergy Clin Immunol* 1997;100:58-64.
- 114 Crimi E, Spanevello A, Neri M, et al. Dissociation between airway inflammation and airway hyperresponsiveness in allergic asthma. *Am J Respir Crit Care Med* 1998;157:4-9.
- 115 Liakakos P, Snell GI, Ward C, et al. Bronchial hyperresponsiveness in lung transplant recipients: lack of correlation with airway inflammation. *Thorax* 1997;52:551-6.
- 116 Cockcroft DW, Berscheid BA, Murdock KY. Unimodal distribution of bronchial responsiveness to inhaled histamine in a random human population. *Chest* 1983;83:751-4.

- 117 Hoag JE, McFadden ER. Long-term effect of cromolyn sodium on nonspecific bronchial hyperresponsiveness: a review. *Ann Allergy* 1991;**66**:53-63.
- 118 Platt-Mills TAE, Tovey ER, Mitchell EB, *et al.* Reduction of bronchial hyperreactivity during prolonged allergen avoidance. *Lancet* 1983;ii:675-8.
- 119 Cockcroft DW, Murdock KY, Berscheid BA. Relationship between atopy and bronchial responsiveness to histamine in a random population. *Ann Allergy* 1984;**53**:26-9.
- 120 Robinson DS, Ying S, Bentley AM, *et al.* Relationship among number of bronchoalveolar lavage cells expressing messenger ribonucleic acid for cytokines, asthma symptoms, and airway methacholine responsiveness in atopic asthma. *J Allergy Clin Immunol* 1993;**92**:397-403.
- 121 Foster PS, Hogan SP, Ramsay AJ, *et al.* Interleukin 5 deficiency abolishes eosinophilia, airway hyperreactivity, and lung damage in a mouse asthma model. *J Exp Med* 1996;**183**:195-201.
- 122 Hogan SP, Koskinen A, Matthaei KI, *et al.* Interleukin-5-producing CD4⁺ T cells play a pivotal role in aeroallergen-induced eosinophilia, bronchial hyperreactivity, and lung damage in mice. *Am J Respir Crit Care Med* 1988;**157**:210-8.
- 123 Hessel EM, Van Oosterhout AJM, Van Ark I, *et al.* Development of airway hyperresponsiveness is dependent on Interferon γ and independent of eosinophils infiltration. *Am J Respir Cell Mol Biol* 1997;**16**:325-34.
- 124 Haley KJ, Drazen JM. Inflammation and airway function in asthma: what you see is not necessarily what you get. *Am J Respir Crit Care Med* 1998;**157**:1-3.
- 125 Dold S, Wjst M, Mutius E, *et al.* Genetic risk for asthma, allergic rhinitis, and atopic dermatitis. *Arch Dis Child* 1992;**67**:1018-22.