Pulmonary neuronal M₂ muscarinic receptor function in asthma and animal models of hyperreactivity

R W Costello, D B Jacoby, A D Fryer

The dominant innervation of the airway smooth muscle is mediated by parasympathetic fibres which are carried in the vagus nerves. Activation of these cholinergic nerves releases acetylcholine which binds to M₃ muscarinic receptors on the smooth muscle causing contraction.¹ Acetylcholine also feeds back onto neuronal M₂ muscarinic receptors located on the postganglionic cholinergic nerves. Stimulation of these receptors further limits acetylcholine release, so these M₂ muscarinic receptors act as autoreceptors.²⁻³ Loss of function of these M₂ receptors, as occurs in some patients with asthma and in animal models of hyperreactivity, leads to an increase in vagally mediated hyperreactivity. In this review we shall discuss the mechanisms that may account for the loss of function of these neuronal M₂ muscarinic receptors.

Innervation of the airways by parasympathetic nerves

The vagus nerves carry preganglionic nerve fibres from the vagal nuclei in the medulla to ganglia in the airways.⁴ These parasympathetic ganglia are interspersed irregularly along the posterior aspect of the wall of the trachea and major bronchi.⁵ From these ganglia short postganglionic nerves pass forward to innervate the airway smooth muscle, the bronchial circulation, and the glandular acini.⁶⁻⁹ Histological studies have not shown postganglionic efferent fibres beyond the level of the terminal bronchi,⁴ and functional studies have not found an effect of vagal stimulation on the respiratory bronchioles and the alveoli.¹⁰⁻¹¹ The site of the most dense cholinergic innervation—the major bronchi—is also the site of bronchoconstriction in patients with asthma, suggesting an underlying pathogenic relationship.

Cholinergic receptors in the airways

Acetylcholine acts on both muscarinic and nicotinic receptors. Five different muscarinic receptors (M₁–M₅) have been genetically sequenced. These M₁–M₅ muscarinic receptors can also be identified based on differing binding affinities between different antagonists. M₁ muscarinic receptors are selectively blocked by pirenzepine, muscarinic M₂ receptors are blocked by AF-DX116 and gallamine, M₃ receptors are blocked by 4-DAMP, while M₄ receptors are antagonised by homobocine; it has been difficult to identify a selective agonist at M₅ receptors. Autoradiographic studies have demonstrated muscarinic M₁ and M₂ receptors along nerve bundles and within the cholinergic ganglia.¹²⁻¹³ Primary cultures of postganglionic cholinergic neurons from the trachea have

shown these nerves to possess messenger RNA for only the M₂ receptors.¹⁴ Airway smooth muscle cells express M₂ and M₃ muscarinic receptors; the latter mediate smooth muscle contraction.¹⁵⁻¹⁶

Physiological function of cholinergic nerves in the airways

Stimulation of the parasympathetic nerves releases acetylcholine which causes the airway smooth muscle to contract,¹⁵⁻¹⁹ the glandular tissue to secrete mucus,²⁰⁻²¹ and the bronchial circulation to dilate.²²⁻²³ Studies both in man and in animals have shown that, in addition to causing contraction, the vagus nerves also maintain a baseline tonic contraction of the airway smooth muscle.¹⁷⁻²⁴⁻²⁶ This baseline tonic contraction of the airways has been demonstrated both in human and in animal studies. For example, in normal non-asthmatic humans 80 µg of inhaled ipratropium bromide caused a 40% reduction in airway resistance, demonstrating that the vagus nerves are important in maintaining airway tone.²⁷ Furthermore, directly inhibiting the vagus nerves—for example, by cutting them—causes bronchodilation.²⁸ In vivo recording of the neural impulses in the vagi of cats and dogs has shown that even at rest there is neural activity in the parasympathetic ganglia.²²⁻²⁹⁻³⁰ The resting airway tone is higher in asthmatic subjects than in normal controls. This increased tone was completely blocked by ipratropium bromide, indicating that it was vagally mediated.²⁶ This shows that in patients with asthma there is increased vagal nerve activity at rest.

Such a powerful bronchoconstricting mechanism needs to be tightly controlled and this is best done close to the site of release of acetylcholine. Indeed, the most important local control over acetylcholine release from postganglionic cholinergic nerves is exerted by acetylcholine itself. Acetylcholine acting on inhibitory muscarinic M₂ autoreceptors located prejunctionally on postganglionic nerves limits the further release of acetylcholine (fig 1). Thus, these receptors act as autoreceptors.²⁻³⁻³¹

The function of the neuronal M₂ autoreceptor can be demonstrated in vivo with M₂ receptor antagonists such as gallamine which cause a dose dependent potentiation of vagally mediated bronchoconstriction. For example, gallamine 10 mg/kg increases vagally induced bronchoconstriction by as much as five fold in pathogen-free guinea pigs (fig 2A).³² Conversely, the muscarinic agonist pilocarpine stimulates neuronal M₂ muscarinic receptors and so decreases vagally induced bronchoconstriction. Pilocarpine 100 µg/kg reduces vagally

Department of
Medicine, University
of Liverpool, Liverpool
L69 3GA, UK
R W Costello

Department of
Environmental Health
Sciences, School of
Hygiene and Public
Health
A D Fryer

Division of Pulmonary
and Critical Care
Medicine
D B Jacoby

Johns Hopkins Asthma
and Allergy Center,
Johns Hopkins
University, Baltimore,
Maryland, USA

Correspondence to:
Dr R W Costello.

Figure 1 The pulmonary neuronal M_2 muscarinic receptor. Acetylcholine released from pulmonary vagal nerves stimulates muscarinic M_3 receptors on airway smooth muscle causing smooth muscle contraction; at the same time acetylcholine stimulates M_2 muscarinic receptors located on the postganglionic nerves. Stimulation of these neuronal M_2 muscarinic receptors limits further acetylcholine release.

mediated bronchoconstriction by about 75% (fig 2B). The presence of an M_2 autoreceptor has also been confirmed by measuring changes in induced acetylcholine release using high performance liquid chromatography in the presence of selective M_2 receptor antagonists.³³ Although first described in the airways of guinea pigs, M_2 receptors have been described in the airways of all species studied, including humans.³³⁻³⁵

Loss of function of neuronal muscarinic M_2 receptors in animal models of hyperreactivity

In antigen sensitised animals exposure to antigen causes an immediate bronchoconstriction followed by a period of increased reactivity to a variety of stimuli. This increased reactivity can be blocked with anticholinergic agents or, alternatively, by cutting the vagus nerves, suggesting that it is vagally mediated.³⁶⁻³⁷

Increased vagally mediated bronchoconstriction may arise because of an increase in the reactivity of the airway smooth muscle or because the function of the neuronal M_2 muscarinic receptor is impaired. These alternative explanations were tested in an animal model of hyperreactivity where the function of the neuronal M_2 muscarinic receptor and the smooth muscle response to acetylcholine were compared between control and antigen sensitised guinea pigs.³⁸ The bronchoconstriction induced by acetylcholine was the same in vagotomised control and antigen challenged animals, indicating that the muscarinic M_3 receptor on airway smooth muscle was functioning normally.³⁸ In contrast, in antigen challenged animals gallamine no longer potentiated and pilocarpine no longer attenuated the mag-

Figure 2 The neuronal M_2 muscarinic receptor does not function in antigen sensitised guinea pigs after challenge. Results are expressed as a ratio of the response to vagal stimulation after the antagonist gallamine (A) or the selective agonist pilocarpine (B) to the response before the compound. In control animals gallamine potentiates vagally induced bronchoconstriction and pilocarpine inhibits vagally induced bronchoconstriction. In contrast, in antigen sensitised animals after challenge gallamine does not potentiate and pilocarpine does not inhibit vagally induced bronchoconstriction, indicating loss of M_2 receptor function. Adapted from Fryer and Wills-Karp.³⁸

nitude of vagally induced bronchoconstriction (fig 2A and B). Thus, the function of M_2 muscarinic receptors is impaired in antigen sensitised animals after challenge. These findings have subsequently been confirmed in other experiments using different models of antigen challenge.³⁹⁻⁴² Increased concentrations of acetylcholine have been reported in the airways of other antigen challenged animals including mice,⁴³ dogs,⁴⁴ and guinea pigs,⁴⁵ providing indirect supportive evidence that there is loss of function of neuronal M_2 muscarinic receptors in the airways of animal models of hyperreactivity.

Antigen induced hyperreactivity is associated with an influx of inflammatory cells, particularly eosinophils and lymphocytes, into the airway walls. A number of studies have either inhibited the recruitment of these cells to the airways or neutralised specific products of these cells to establish their role in hyperreactivity.⁴⁵⁻⁴⁶ The results of these studies have specifically implicated the eosinophil in the development of antigen induced hyperreactivity. Since antigen induced vagally mediated hyperreactivity is due to loss of function of the neuronal M_2 receptor, the role of eosinophils in the loss of function of M_2 receptors has been investigated.

The selective localisation of leucocytes to sites of inflammation is mediated through the interactions of specific adhesion molecules.

Very late activation antigen 4 (VLA-4) is the major β integrin expressed by eosinophils and it recognises the counter ligand vascular adhesion molecule 1 (VCAM-1) expressed on vascular tissues, allowing migration into the airways.^{47, 48} Pretreating antigen sensitised animals with an antibody to the adhesion molecule VLA-4 before challenge prevented antigen induced eosinophil accumulation in the airways and prevented loss of function of M_2 muscarinic receptors and the development of airway hyperreactivity.³⁹ Depleting eosinophils by using a monoclonal antibody to neutralise the eosinophil chemoattractant interleukin 5 prevented antigen induced airway eosinophilia and loss of function of neuronal M_2 muscarinic receptors.⁴⁹ Thus, eosinophils are responsible for loss of pulmonary neuronal M_2 muscarinic receptor function in the airways of antigen challenged guinea pigs.

A potential mechanism for this eosinophil dependent loss of M_2 receptor function in antigen challenged animals was suggested by the finding that some eosinophil products are antagonists at M_2 muscarinic receptors. Eosinophils contain electron dense granules containing eosinophil cationic protein (ECP), eosinophil derived neurotoxin (EDN), eosinophil peroxidase (EPO), and eosinophil major basic protein (MBP).^{50, 51} These heavy granules (molecular weight 14–77 kD) comprise approximately 90% of eosinophil granular proteins. These four proteins have the common characteristic of being cytotoxic to mammalian cells and also of possessing high isoelectric points (pH range 10–11.5). In common with these eosinophil proteins, many antagonists at M_2 muscarinic receptors such as gallamine and protamine⁵² are positively charged. The cationic nature of these antagonists is important in binding to M_2 muscarinic receptors, possibly because the receptor is heavily sialated giving it a net negative charge. In receptor binding studies on M_2 and M_3 receptors MBP and, to a lesser extent, EPO displaced the agonist [3 H]N-methylscopolamine ([3 H]NMS) from guinea pig and human M_2 but not M_3 muscarinic receptors.^{53, 54} Furthermore, in the presence of the anionic compound heparin, MBP was displaced from these receptors. This suggests that the antagonism of these M_2 receptors was reversible and due to the positively charged nature of MBP. In saturation binding studies it was shown that the antagonism of MBP at M_2 receptors was allosteric rather than competitive. Eosinophil MBP may have a physiologically relevant role in the loss of M_2 receptor function since the dissociation constant for MBP at M_2 receptors is 1.4×10^{-5} M which is only minimally higher than the level of MBP found in the sputum of patients with acute asthma.⁵⁵ In contrast, eosinophil peroxidase has a low affinity for M_2 receptors and so does not appear to be physiologically relevant as an antagonist at M_2 receptors.

As binding studies have shown that heparin displaces MBP from M_2 muscarinic receptors, and because eosinophil MBP may be responsible for the loss of function of the neuronal M_2

muscarinic receptor, the effect of heparin on M_2 receptor function was tested in antigen challenged animals. In these studies it was shown that the administration of heparin acutely restored M_2 receptor function in antigen challenged guinea pigs and rats⁴⁰ (Belmonte *et al.*, unpublished). These findings suggest that M_2 receptors become dysfunctional after antigen challenge and that positively charged proteins such as MBP are acting as endogenous antagonists at M_2 receptors.

In order to establish a role for eosinophil MBP in the loss of M_2 receptor function, in vivo studies were performed with a specific neutralising antibody to eosinophil MBP.⁵⁶ In these studies antigen sensitised guinea pigs were pretreated with an antibody to MBP before challenge and the function of the M_2 receptor as well as vagally mediated hyperreactivity were tested 24 hours after challenge. In sensitised animals studied after antigen challenge there was loss of function of M_2 receptors and an increase in vagally mediated bronchoconstriction. In contrast, in antigen sensitised animals pretreated with the antibody to MBP the function of M_2 receptors was preserved and hyperreactivity inhibited (fig 3). Thus, it is likely that eosinophil products, in particular MBP, are responsible for the loss of M_2 receptor function in antigen challenged animals.

Since MBP is highly cationic and probably does not diffuse far after it is released from eosinophils, there must be a close anatomical association between eosinophils and airway nerves. In histological studies it was shown that there is a close association of both eosinophils and extracellular MBP with airway nerves in patients with asthma and in animal models of hyperreactivity (fig 4). Furthermore, the number of eosinophils per nerve was correlated with the in vivo function of neuronal M_2 muscarinic receptors.⁴² Thus, there is good evidence to indicate that eosinophil MBP is involved in the loss of function of neuronal M_2 muscarinic receptors in antigen challenged animals.

Figure 3 An antibody to eosinophil MBP protects neuronal M_2 muscarinic receptor function in antigen challenged guinea pigs. Results are expressed as a ratio of the response to vagal stimulation after pilocarpine to the response before pilocarpine. In control animals (open circles) pilocarpine inhibited vagally induced bronchoconstriction but in antigen challenged guinea pigs (closed circles) it did not. Pretreatment with an antibody to MBP before antigen challenge protected the response to pilocarpine. Adapted from Evans *et al.*⁵⁶

Loss of function of neuronal M₂ muscarinic receptors after exposure to a respiratory viral infection

Infection with a respiratory virus causes an increase in vagally mediated hyperreactivity.^{57–58} Studies in guinea pigs and in rats infected with parainfluenza virus indicate that there is loss of function of the neuronal M₂ muscarinic receptor.^{59–60} The central binding site of the M₂ muscarinic receptor is composed of negatively charged sialated glycoproteins.⁶¹ Respiratory viruses, in particular parainfluenza virus, contain the enzyme neuraminidase which cleaves sialic acid residues.^{62–63} In receptor binding studies it has been shown that the affinity of M₂ receptors for the agonist [3H] quinuclidinyl benzilate ([3H] QNB) is impaired when they are incubated with neuraminidase or when lung tissue of virally infected animals is studied.⁶⁴ These data suggest that the viral enzyme neuraminidase may be involved in the loss of function of neuronal M₂ muscarinic receptors. Inflammatory cells also appear to play a part in the loss of function of the neuronal M₂ receptor.⁶⁵ Since heparin does not restore the function of the M₂ receptor in non-sensitised virally infected animals, the inflammatory cell causing this impaired neuronal M₂ receptor function does not appear to be the eosinophil. Prior antigen sensitisation alters the immune response to a viral infection away from the normal lymphocyte response to an eosinophil rich response.⁶⁶ In antigen sensitised guinea pigs viral infections cause pulmonary eosinophilia and heparin restores the function of M₂ receptors, which suggests that the M₂ receptor dysfunction may also be eosinophil mediated (Fryer *et al*, unpublished).

Loss of function of neuronal M₂ muscarinic receptors after exposure to ozone

Exposure of animals to ozone (2 ppm for four hours) causes an increase in vagally mediated hyperreactivity in guinea pigs. In vivo studies in

guinea pigs indicate that there is an immediate loss of function of the neuronal M₂ muscarinic receptor after exposure to ozone that is long lasting.^{67–68} The mechanisms responsible for the loss of function of the neuronal M₂ receptor after exposure to ozone are not well established but appear to be dependent on inflammatory cells rather than to be the direct result of exposure to ozone.⁶⁹ Loss of function of these receptors may be eosinophil mediated, since inhibiting eosinophil recruitment with an antibody to VLA-4 or neutralising MBP prevents this loss of M₂ receptor function (Fryer *et al*, unpublished observation).

Neuronal M₂ muscarinic receptor function in patients with asthma

An increase in both vagally mediated baseline tone and vagally mediated hyperreactivity has been reported in patients with asthma. The increase in vagal hyperreactivity has been demonstrated in studies in which the vagus nerves have been inhibited pharmacologically with drugs such as atropine or ipratropium bromide. The exact contribution of the vagus nerves to the hyperreactivity varies between studies and the inconclusive nature of these studies has led some to suggest that the vagus nerves are not important in the pathogenesis of asthma. This conclusion is unfortunate as many studies either show a clear benefit in some but not all patients,^{70–71} or the investigators have not adequately inhibited vagal nerve function.^{72–74} One particular problem with the currently available anticholinergic agents is their non-selective nature; they antagonise both M₃ and M₂ receptors. Antagonism of the M₂ receptor by these agents will potentiate vagally induced bronchoconstriction and thus counteract the effect of inhibiting the M₃ receptor on the smooth muscle.⁷⁵

The presence of neuronal M₂ muscarinic receptors has also been described in humans. In these in vivo studies vagally induced bronchoconstriction is induced indirectly via a

Figure 4 Eosinophils are found in association with nerve fibres in the airways of antigen challenged guinea pigs. The photomicrograph is from a paraffin embedded section of antigen challenged guinea pig bronchus. Eosinophils (detected with chromotrope 2R) are seen in close proximity to airway nerve fibres in the airway smooth muscle (detected with an antibody to PGP 9.5, in black). The nerve fibre in the smooth muscle shown with the thin arrow in (A) is shown under higher magnification in (B). Eosinophils were also seen around the airway nerve bundles near the smooth muscle in the submucosa (thick arrows). From Evans *et al*.⁵⁶

vagal reflex with an agent such as sulphur dioxide or histamine.³⁵⁻⁷⁶ In the presence of a normal M₂ muscarinic receptor pilocarpine stimulates these receptors and thus limits the degree of vagally mediated bronchoconstriction. In addition to these indirect *in vivo* studies, *in vitro* studies on surgically resected tissue from non-asthmatic individuals have also shown the presence of the neuronal M₂ receptor.³³

Most studies on the function of the neuronal M₂ muscarinic receptor function after exposure to antigen have been carried out in animal models of hyperreactivity. This is because it is not feasible to stimulate the vagus nerve directly in humans and because lung tissue which could be used for *in vitro* studies is rarely resected from patients with asthma. However, there is evidence that there is loss of function of neuronal M₂ receptors in some patients with asthma.³⁵⁻⁷⁶⁻⁷⁷ In these *in vivo* studies the subjects had stable allergic asthma and in all but one of these studies inhaled pilocarpine had no effect on vagally induced bronchoconstriction, indicating impaired function of these receptors. The reasons for the differences in the results of these studies may reflect the techniques used to induce the vagal reflex bronchoconstriction or, alternatively, the severity of asthma. The function of the neuronal M₂ muscarinic receptor has been recently tested in people with very mild asthma with a history of wheeze during a viral infection. In this preliminary study there was normal function of the M₂ receptor at baseline and a transient loss of function of the M₂ receptor during a viral respiratory infection (Costello *et al*, unpublished observation).

The underlying mechanism of loss of function of M₂ receptors has been investigated. One recent study has shown that inhaled heparin prevented late phase hyperreactivity to allergen in sensitive asthmatic subjects. The late phase response is characterised by an inflammatory cell influx, in particular of eosinophils. It is tempting to speculate that this effect of heparin was due to an effect on M₂ receptor function analogous to that seen in antigen challenged guinea pigs, although further studies will be required to test this hypothesis.⁷⁸

Summary

In the lungs neuronal M₂ muscarinic receptors limit acetylcholine release from postganglionic cholinergic nerves. These inhibitory M₂ receptors are dysfunctional in antigen challenged guinea pigs and in humans with asthma which leads to an increase in vagally mediated hyperreactivity. *In vitro*, eosinophil products act as allosteric antagonists at neuronal M₂ muscarinic receptors. *In vivo*, displacing or neutralising MBP preserves neuronal M₂ muscarinic receptor function and prevents hyperreactivity. Thus, there is good evidence from animal studies that after antigen challenge pulmonary M₂ muscarinic receptors become dysfunctional because MBP inhibits their function. Loss of function of pulmonary neuronal M₂ muscarinic receptors has also been reported in patients

with asthma, although the clinical significance of this dysfunction and the mechanisms underlying it are not yet established.

This work has been funded by grants from The British Lung Foundation (RWC) and grants HL-54659 (DBJ) and HL 55543 (ADF) from the National Institutes of Health, USA.

- 1 Roffel AF, Elzinga CRS, Zaagsma J. Muscarinic M₂ receptors mediate contraction of human central and peripheral airway smooth muscle. *Pulm Pharm* 1990;3:47-51.
- 2 Fryer AD, MacLagan J. Muscarinic inhibitory receptors in pulmonary parasympathetic nerves in the guinea-pig. *Br J Pharmacol* 1984;83:973-8.
- 3 Blaber LC, Fryer AD, MacLagan J. Neuronal muscarinic receptors attenuate vagally-induced contraction of feline bronchial smooth muscle. *Br J Pharmacol* 1985;86:723-8.
- 4 Richardson J. Nerve supply to the lungs. *Am Rev Respir Dis* 1979;119:785-802.
- 5 Honjin R. On the nerve supply of the lung of the mouse with special reference to the structure of the peripheral vegetative and nervous system. *J Comp Neurol* 1956;105:587.
- 6 El-Bermani AW, Grant M. Acetylcholinesterase-positive nerves of the rhesus monkey bronchial tree. *Thorax* 1975;30:162-70.
- 7 El-Bermani AW. Pulmonary noradrenergic innervation of rat and monkey: a comparative study. *Thorax* 1978;33:167-74.
- 8 Spencer H, Leof D. The innervation of the human lung. *J Anat* 1964;98:599.
- 9 Baker DG, McDonald DM, Basbaum CB, *et al*. The architecture of nerves and ganglia of the ferret trachea as revealed by acetylcholinesterase histochemistry. *J Comp Neurol* 1986;246:513-26.
- 10 Nadel J, Cabezas G, Austin J. *In vivo* roentgenographic examination of parasympathetic innervation of small airways. Use of powdered tantalum and a fine focal spot X-ray tube. *Invest. Radiol.* 1971;6:9-17.
- 11 ten Berge RE, Zaagsma J, Roffel AF. Muscarinic inhibitory autoreceptors in different generations of human airways. *Am J Respir Crit Care Med* 1996;154:43-9.
- 12 van Koppen CJ, Blankesteijn WM, Klassen ABM, *et al*. Autoradiographic visualization of muscarinic receptors in pulmonary nerves and ganglia. *Neurosci Lett* 1987;83:237-40.
- 13 van Koppen C, Blankesteijn W, Klassen A, *et al*. Autoradiographic visualization of muscarinic receptors in human bronchi. *J Pharmacol Exp Ther* 1988;244:760-4.
- 14 Fryer AD, Elbon CL, Kim AL, *et al*. Cultures of airway parasympathetic nerves express functional M₂ muscarinic receptors. *Am J Respir Cell Mol Biol* 1996;15:716-25.
- 15 Madison JM, Jones CA, Tom-Moy M, *et al*. Affinities of pirenzepine for muscarinic cholinergic receptors in membranes isolated from bovine tracheal mucosa and muscle. *Am Rev Respir Dis* 1987;135:719-24.
- 16 Maeda A, Kubo T, Mishina M, *et al*. Tissue distribution of mRNAs encoding for acetylcholine receptor subtypes. *FEBS Lett* 1988;239:339-42.
- 17 Colebatch H, Halmagyi D. Effect of vagotomy and vagal stimulation on lung mechanics and circulation. *J Appl Physiol* 1963;18:881-7.
- 18 Olsen C, Colebatch H, Mebel P, *et al*. Motor control of pulmonary airways studied by nerve stimulation. *J Appl Physiol* 1965;20:202-8.
- 19 Green M, Widdicombe JG. The effects of ventilation of dogs with different gas mixtures on airway calibre and lung mechanics. *J Physiol* 1966;186:363-81.
- 20 Brody JS, Klempfner G, Staum MM, *et al*. Mucociliary clearance after lung denervation and bronchial transection. *J Appl Physiol* 1972;32:160-4.
- 21 Gallagher JT, Kent PW, Passatore M, *et al*. The composition of tracheal mucus and the nervous control of its secretion in the cat. *Proc R Soc Lond* 1976;192:49-76.
- 22 Widdicombe JG. Regulation of tracheobronchial smooth muscle. *J Physiol* 1966;43:1-37.
- 23 Phipps R, Richardson P. The effects of irritation at various levels of the airway upon tracheal mucous secretion in the cat. *J Physiol* 1976;261:563-81.
- 24 Severinghaus JW, Stupfel M. Respiratory dead space increases following atropine in man and atropine, vagal or ganglionic blockade and hypothermia in dogs. *J Appl Physiol* 1955;8:81-6.
- 25 Cabezas GA, Kessler GF, Yu DYD. Sympathetic nerve versus parasympathetic nervous regulation of airways of dogs. *J Appl Physiol* 1971;31:651-5.
- 26 Molino NA, Slutsky AS, Julia-Serda G, *et al*. Assessment of airway tone in asthma. Comparison between double lung transplant patients and healthy subjects. *Am Rev Respir Dis* 1993;148:1238-43.
- 27 Douglas NJ, Sudlow MF, Flenley DC. Effect of an inhaled atropinelike agent on normal airway function. *J Appl Physiol* 1979;46:256-62.
- 28 Karczewski W, Widdicombe JG. The effect of vagotomy, vagal cooling and efferent vagal stimulation on breathing and lung mechanics in rabbits. *J Physiol* 1969;200:25-49.
- 29 Widdicombe JG. Action potentials in vagal efferent nerve fibres to the lung of the cat. *Naunyn-Schmiedeberg's Arch Exp Pathol Pharmacol* 1961;241:451-9.
- 30 Widdicombe J. Action potentials in parasympathetic and sympathetic efferent fibres to the trachea and lungs of dogs and cats. *J Physiol* 1966;186:156-88.

- 31 Faulkner D, Fryer AD, MacLagan J. Post-ganglionic muscarinic receptors in pulmonary parasympathetic nerves in the guinea pig. *Br J Pharmacol* 1986;88:181-8.
- 32 Fryer AD, MacLagan J. Pancuronium and gallamine are antagonists for pre- and post-junctional muscarinic receptors in the guinea pig lung. *Naunyn-Schmiedeberg's Arch Pharmacol* 1987;335:367-71.
- 33 Patel HJ, Barnes PJ, Takahashi T, et al. Evidence for prejunctional muscarinic autoreceptors in human and guinea pig trachea. *Am J Respir Crit Care Med* 1995;152:872-9.
- 34 Minette P, Barnes PJ. Prejunctional inhibitory muscarinic receptors on cholinergic nerves in human and guinea-pig airways. *J Appl Physiol* 1988;64:2532-7.
- 35 Ayala LE, Ahmed T. Is there loss of a protective muscarinic receptor in asthma? *Chest* 1989;96:1285-91.
- 36 Costello RW, Evans CM, Yost BL, et al. Antigen induced hyperactivity is vagally-mediated in guinea pigs. *Am J Respir Crit Care Med* 1997;155:A572.
- 37 Santing RE, Pasman Y, Olymulder CG, et al. Contribution of a cholinergic reflex mechanism to allergen-induced bronchial hyperactivity in permanently instrumented, unrestrained guinea-pigs. *Br J Pharmacol* 1995;114:414-8.
- 38 Fryer AD, Wills-Karp M. Dysfunction of M₂ muscarinic receptors in pulmonary parasympathetic nerves after antigen challenge in guinea-pigs. *J Appl Physiol* 1991;71:2255-61.
- 39 Fryer AD, Costello RW, Yost BL, et al. Antibody to VLA-4, but not to L-selectin, protects neuronal M₂ muscarinic receptors in antigen-challenged guinea pig airways. *J Clin Invest* 1997;99:2036-44.
- 40 Fryer AD, Jacoby DB. Function of pulmonary M₂ muscarinic receptors in antigen challenged guinea-pigs is restored by heparin and poly-L-glutamate. *J Clin Invest* 1992;90:2292-8.
- 41 ten Berge RE, Santing RE, Hamstra JJ, et al. Dysfunction of muscarinic M₂ receptors after the early allergic reaction: possible contribution to bronchial hyperresponsiveness in allergic guinea-pigs. *Br J Pharmacol* 1995;114:881-7.
- 42 Costello RW, Schofield BH, Kephart GM, et al. Localization of eosinophils to airway nerves and effect on neuronal M₂ muscarinic receptor function. *Am J Physiol* 1997;273:L93-103.
- 43 Larsen GL, Fame TM, Renz H, et al. Increased acetylcholine release in tracheas from allergen-exposed IgE-immune mice. *Am J Physiol* 1994;266:L263-70.
- 44 Walters EH, O'Byrne PM, Graf PD, et al. The responsiveness of airway smooth muscle in vitro from dogs with airway hyperresponsiveness in vivo. *Clin Sci* 1986;71:605-11.
- 45 Pretolani M, Ruffie C, Lapa e Silva JR, et al. Antibody to very late activation antigen-4 prevents antigen-induced bronchial hyperactivity and cellular infiltration in the guinea pig airways. *J Exp Med* 1994;180:795-805.
- 46 Lefort J, Nahori MA, Ruffie C, et al. In vivo neutralization of eosinophil-derived major basic protein inhibits antigen induced bronchial hyperactivity in sensitized guinea pigs. *J Clin Invest* 1996;97:1117-21.
- 47 Walsh GM, Mermoud J, Hartnell A, et al. Human eosinophil but not neutrophil adherence to IL1 stimulated human umbilical vascular endothelial cells is very late antigen-4 dependent. *J Immunol* 1991;146:3419-23.
- 48 Walsh GM, Hartnell A, Wardlaw AJ, et al. IL-5 enhances the in vitro adhesion of human eosinophils, but not neutrophils, in a leukocyte integrin (CD11/18)-dependent manner. *Immunology* 1990;71:258-65.
- 49 Elbon CL, Jacoby DB, Fryer AD. Pretreatment with an antibody to IL-5 preserves the function of pulmonary M₂ muscarinic receptors in antigen challenged guinea-pigs. *Am J Respir Cell Mol Biol* 1995;12:320-8.
- 50 Gleich GJ, Loegering DA. Selective stimulation and purification of eosinophils and neutrophils from guinea pig peritoneal fluids. *J Lab Clin Med* 1973;82:522-8.
- 51 Gleich GJ, Loegering DA, Mann KG, et al. Comparative properties of the Charcot-Leyden crystal protein and the major basic protein from human eosinophils. *J Clin Invest* 1976;57:633-40.
- 52 Hu J, Wang S, Forray C, et al. Complex allosteric modulation of cardiac muscarinic receptors by protamine: a potential model for putative endogenous ligands. *Mol Pharmacol* 1992;42:311-24.
- 53 Jacoby DB, Gleich GJ, Fryer AD. Interaction of human eosinophil major basic protein with cloned human muscarinic receptors. *Am J Respir Crit Care Med* 1995;151:A821.
- 54 Jacoby DB, Gleich GJ, Fryer AD. Human eosinophil major basic protein is an endogenous allosteric antagonist at the inhibitory muscarinic M₂ receptor. *J Clin Invest* 1993;91:1314-8.
- 55 Frigas E, Loegering DA, Solley GO, et al. Elevated levels of the eosinophil granule MBP in the sputum of patients with bronchial asthma. *Mayo Clin Proc* 1981;56:345-53.
- 56 Evans CM, Fryer AD, Jacoby DB, et al. Pretreatment with antibody to eosinophil major basic protein prevents hyperresponsiveness by protecting neuronal M₂ muscarinic receptors in antigen-challenged guinea pigs. *J Clin Invest* 1997;100:2254-62.
- 57 Empey DW, Laitinen LA, Jacobs L, et al. Mechanisms of bronchial hyperactivity in normal subjects following upper respiratory tract infection. *Am Rev Respir Dis* 1976;113:523-7.
- 58 Aquilina AT, Hall WJ, Douglas RG, et al. Airway reactivity in subjects with viral upper respiratory tract infections: the effects of exercise and cold air. *Am Rev Respir Dis* 1980;122:3-10.
- 59 Fryer AD, Jacoby DB. Parainfluenza virus infection damages inhibitory M₂ muscarinic receptors on pulmonary parasympathetic nerves in the guinea-pig. *Br J Pharmacol* 1991;102:267-71.
- 60 Sorkness R, Clough JJ, Castelman WL, et al. Virus induced airway obstruction and parasympathetic hyperresponsiveness in adult rats. *Am Rev Respir Dis* 1994;150:28-34.
- 61 Peterson GL, Rosenbaum LC, Broderick DJ, et al. Physical properties of the purified cardiac muscarinic acetylcholine receptor. *Biochemistry* 1986;25:3189-202.
- 62 Scheid A, Caliguri LA, Compans RW, et al. Isolation of paramyxovirus glycoproteins. Association of both hemagglutinating and neuraminidase activities with the larger SV5 glycoprotein. *Virology* 1972;50:640-52.
- 63 Laver WG. The structure of influenza viruses. 3. Disruption of the virus particle and separation of neuraminidase activity. *Virology* 1963;20:251-62.
- 64 Fryer AD, El-Fakahany EE, Jacoby DB. Parainfluenza virus type 1 reduces the affinity of agonists for muscarinic receptors in guinea-pig heart and lung. *Eur J Pharmacol* 1990;181:51-8.
- 65 Fryer AD, Jacoby DB. Effect of inflammatory cell mediators on M₂ muscarinic receptors in the lungs. *Life Sci* 1993;52:529-36.
- 66 Coyle AJ, Erard F, Bertrand C, et al. Virus-specific CD8+ cells can switch to interleukin 5 production and induce airway eosinophilia. *J Exp Med* 1995;181:1229-33.
- 67 Schultheis AH, Bassett DJP, Fryer AD. Ozone-induced airway hyperresponsiveness and loss of neuronal M₂ muscarinic receptor function. *J Appl Physiol* 1994;76:1088-97.
- 68 Schultheis AJH. The effect of ozone on inflammatory cell infiltration and airway hyperresponsiveness in the guinea pig lung. PhD thesis, Johns Hopkins University, 1992.
- 69 Gambone LM, Elbon CL, Fryer AD. Ozone-induced loss of neuronal M₂ muscarinic receptor function is prevented by cyclophosphamide. *J Appl Physiol* 1994;77:1492-9.
- 70 Cockcroft DW, Ruffin RE, Hargreave FE. Effect of Sch100 in allergen-induced asthma. *Clin Allergy* 1978;8:361-72.
- 71 Ruffin R, Cockcroft D, Hargreave F. A comparison of the protective effect of fenoterol and Sch100 on allergen induced asthma. *J Allergy Clin Immunol* 1978;61:42-7.
- 72 Rosenthal RR, Norman PS, Summer WR, et al. Role of the parasympathetic system in antigen induced bronchospasm. *J Appl Physiol* 1977;42:600-6.
- 73 O'Byrne P, Thomson NC, Morris M, et al. The protective effect of inhaled chlorpheniramine and atropine on bronchoconstriction stimulated by airway cooling. *Am Rev Respir Dis* 1983;128:611-7.
- 74 Fish J, Rosenthal R, Summer W, et al. The effect of atropine on acute antigen mediated airway constriction in subjects with allergic asthma. *Am Rev Respir Dis* 1977;115:371-9.
- 75 Fryer AD, MacLagan J. Ipratropium bromide potentiates bronchoconstriction induced by vagal nerve stimulation in the guinea-pig. *Eur J Pharmacol* 1987;139:187-91.
- 76 Minette PJ, Lammers JWJ, Dixon CMS, et al. A muscarinic agonist inhibits reflex bronchoconstriction in normal but not asthmatic subjects. *J Appl Physiol* 1989;67:2461-5.
- 77 Okayama M, Shen T, Midorikawa J, et al. Effect of pilocarpine on propranolol induced bronchoconstriction in asthma. *Am J Respir Crit Care Med* 1994;149:76-80.
- 78 Diamant Z, Timmers MC, van der Veen H, et al. Effect of inhaled heparin on allergen-induced early and late asthmatic responses in patients with atopic asthma. *Am J Respir Crit Care Med* 1996;153:1790-5.