Socioeconomic status and chronic obstructive pulmonary disease

Eva Prescott, Jørgen Vestbo

Socioeconomic status, whether measured by education, income, or other indices of social class, has long been known to be associated with mortality from different diseases.¹⁻⁷ This knowledge for the major part stems from studies conducted in the UK where there is a tradition of interest in socioeconomic differences. A milestone within recent years was the "Black Report" prepared by a committee appointed by the UK Minister of Health in 1980.⁷ This report showed that there were still large socioeconomic differences in morbidity and mortality in men and women of all ages in England. Since then a number of epidemiological studies have been published, both in England and other European and North American countries, which confirm this relationship.⁸⁻¹²

Socioeconomic differences have been shown to exist for a number of diseases including ischaemic heart disease, many types of cancer, respiratory diseases and, in particular, mortality related to alcohol and violence. In addition, despite a general fall in mortality during recent decades, the relative disadvantage of low socioeconomic status continues to increase.³⁻⁷ The number of papers published on socioeconomic factors and morbidity and mortality is increasing sharply.¹³ The social gradient in ischaemic heart disease has been extensively studied but, although there are indicators that socioeconomic differences in respiratory diseases are greater still,² very few studies have focused on this disease entity.

Review of current literature

A social class gradient in the prevalence of respiratory symptoms in adults, in particular cough and sputum, was first observed in early studies in the UK.¹ This has later been confirmed in other studies, most of which date from the 1970s.^{2 14-26} The relationship between socioeconomic factors and symptoms, lung function parameters, and respiratory morbidity in studies in adults is summarised in table 1. Most of these studies have adjusted for smoking habits since smoking is related to both the disease and socioeconomic status.

Only a few studies have attempted to quantify the effect of socioeconomic status on indices of lung function. In a study comparing mortality in manual and non-manual workers in two British towns, respiratory symptoms and impairment of lung function was found to differ across social class.¹⁹ The difference in forced expiratory volume in one second (FEV₁) was 7-15% of FEV₁% predicted and was not affected by stratification by smoking history. In a study of 410 male non-smokers, Stebbings *et al*¹⁴ found that the difference in FEV₁ between the highest and lowest social class was

400 ml.¹⁴ A similar result was found in the Copenhagen City Heart Study in which the difference between the highest and lowest socioeconomic group was 400 ml for men and 259 ml for women after adjustment for smoking.²⁵ This difference was reflected in a threefold higher risk of admission to hospital for chronic obstructive pulmonary disease (COPD) in subjects of the lowest compared with the highest socioeconomic group, and was not affected by adjustment for smoking. Interestingly, similar risk ratios for death from respiratory disease were seen in the Whitehall study.²

Most of these studies have been conducted on middle aged adults and it is worthwhile to remember that lung function in this age group may be the result of several mechanisms.²⁷ A reduced lung function in middle age may result from a low maximally attained lung function due to either low lung function at birth or decreased growth of lung function during childhood and adolescence, a shortened plateau phase during early adult life, or an increased decline in lung function in later adult life. The important unanswered question remains: which risk factors linked to socioeconomic status are causally related to the occurrence of COPD? The link is likely to be multifactorial and includes prenatal exposures, more frequent lower respiratory tract illness in childhood, housing conditions, air pollution, environmental tobacco smoke, diet, and other life-style factors including smoking, in addition to possible genetic factors. As these factors may or may not be directly associated with socioeconomic status, they will be briefly discussed in more detail. A list of the factors of interest in this review is given in table 2.

As for prenatal exposure, the possibility that intrauterine growth of the fetus might be related to health in childhood and adult life has attracted considerable attention.²⁸⁻³⁵ Most studies concern ischaemic heart disease but a link between prenatal exposure and respiratory disease has also been suggested.^{33 36 37} Low birth weight has been associated with reduced lung function, respiratory infections, and asthma in children and adults³⁸⁻⁴⁰ regardless of length of gestation⁴¹ and whether the child had respiratory complications at birth.^{42 43} This gap may continue to widen into adult life as suggested by Barker and co-workers³⁴ who reported that, among men born 60-70 years ago in Hertfordshire, UK, those who had a lower birth weight had a lower FEV₁ adjusted for age and height. The standardised mortality rate for COPD was also associated with birth weight in these men. This was corroborated by a study on Indian men and women in which

Institute of Preventive Medicine,
Kommunehospitalet,
DK-1399 Copenhagen
K, Denmark
E Prescott

Department of Respiratory Medicine
129, H:S Hvidovre
Hospital, DK-2650
Hvidovre, Denmark
J Vestbo

Correspondence to:
Dr E Prescott.

Table 1 Studies of socioeconomic status (SES) and indices of COPD

Author/place	Year	Study population	Index of SES
Studies of adults that include measures of lung function			
Stebbins ¹⁴ (Hagerstown, Maryland)	1971	Non-smokers (410 men)*	Income, education
Cohen ¹⁶ (Baltimore, Maryland)	1977	Genetic-epidemiological (1232 men and women)*	Place of residence
Higgins ¹⁷ (Tecumseh, Michigan)	1977	Population sample (4699 men and women)*	Education, income
Lebowitz ¹⁵ (Tucson, Arizona)	1977	Population sample (3800 men and women)*	Education, income, occupation
Rasmussen ²⁶ (Copenhagen, Denmark)	1978	Non-occupationally exposed (218 men)*	Housing conditions
Krzyzanowski ¹⁸ (Cracow, Poland)	1986	Population sample (1864 men and women)**	Education
Burr ¹⁹ (Caerphilly/Bath, Scotland)	1987	Population sample (513 men)*	Manual, non-manual
Krzyzanowski ²⁰ (7 cities, France)	1988	Population sample (16 000 men and women)*	Social class, education
Bakke ²² (Bergen, Norway)	1995	Population sample (714 men and women)*	Education (3 levels)
Hole ²³ (Renfrew/Paisley, Scotland)	1996	Population sample (15 411 men and women)*	Social class (IV–V/I–III)
Prescott ²⁵ (Copenhagen, Denmark)	1998	Population sample (14 223 men and women)*	Income, education
Other studies of socioeconomic status and indices of COPD			
Marmot ² (Whitehall study, UK)	1984	Civil servants (17 530 men)**	Grade 1–4
Lebowitz ²¹ (Tucson, Arizona)	1990	583 children aged over 6*	Family income
Demissie ²⁴ (Montreal, Canada)	1996	989 children aged 6–11*	Parental SES score 1–5

*Cross sectional study; **longitudinal study.

mean FEV₁ fell by 0.09 litres with each pound decrease in birth weight in men and by 0.06 litres in women, independent of smoking.²⁸ These results are supported by animal experiments which show that lung structure may be permanently altered by calorie or protein deprivation in utero.^{44–46} Changes include reduction of elastin or collagen in the lung, enlargement of air spaces, and a reduction in elastic recoil compared with that seen in human emphysema.^{47–50}

The theory of “intrauterine programming” has been criticised.⁵¹ In addition, whereas intrauterine growth retardation because of malnourishment during pregnancy associated with low socioeconomic status may have been common more than 50 years ago, this is not likely to be a major problem in western countries today. However, today relative growth retardation may be seen in children of smoking mothers. In the Tucson studies it has been shown that children of mothers who smoke during pregnancy have impaired lung function in infancy and are also at increased risk of developing asthma,⁵² and Hanrahan *et al* have found several measures of infant lung function to be negatively affected by in utero exposure to tobacco smoke.⁵³ Whether differences continue and affect lung function into adult life remains to be seen.

Another factor related to socioeconomic status is respiratory infections during infancy. Childhood experience of respiratory disease was associated geographically with mortality from chronic bronchitis.³⁷ Early respiratory tract infections have been shown to be associated with reduced lung volumes independent of the effects of low birth weight.^{29 37 54 55} Studies of the association have consistently shown deficits of lung function of the order of 5–10%.⁵⁶ However, because of

methodological problems it is still not clear whether the relationship is causal or is merely an epiphenomenon in the mesh of associations between lung function at birth, respiratory tract infections with or without wheeze, atopy, asthma, and growth of lung function as recently discussed by Britton and Martinez.⁵⁶ Not only childhood respiratory infections but also infections in adulthood have been linked to impairment of lung function.^{57 58}

Poor housing conditions²⁶ and home dampness^{59 60} with increased house dust mites⁶¹ and gas stove usage^{62 63} are all associated with respiratory symptoms, reduced lung function, and lower socioeconomic status. Household crowding has been hypothesised to cause increased instances of respiratory infections and thus increased rates of respiratory disorders,^{32 64} although this was not confirmed in the Tucson study.²¹ Home dampness and respiratory symptoms are strongly associated, whereas the relationship between home dampness and pulmonary function is weaker in both children and adults.^{59 60 65} Children living in homes using gas for cooking are more likely to have had respiratory illnesses than those from homes using electricity in some^{66–68} but not all studies.^{69 70} In adults the use of gas for cooking was related to an increased frequency of respiratory symptoms and impaired ventilatory function among men but not among women in two studies.^{62 63} Exposure to community air pollution is an additional risk factor to the airways more often experienced by subjects of lower than of higher social class.

Smoking remains the most prominent cause of lung disease. Children from homes with smokers are more likely to have respiratory illnesses than those not exposed to tobacco smoke at home.^{71–74} Smoking in childhood and adolescence affects growth of lung function.⁷⁵ Gold *et al* clearly demonstrated this in the US Six Cities Study where a relatively modest amount of tobacco led to marked reductions in growth of lung function, particularly in teenage girls who in most western countries seem to constitute a group at high risk of persistent smoking. There is no doubt that there is a socioeconomic gradient in smoking behaviour and that this gradient is increasing. A higher

Table 2 Factors potentially associated with socioeconomic status and lung function

Intrauterine growth of lung function
Environmental exposures in childhood
Childhood respiratory infections
Smoking in childhood and adolescence
Smoking in adulthood
Occupation
Housing
Imbalance in protective factors

Outcome	Adjusted for	Result
FEV ₁ , symptoms	Smoking, sex	Association between SES and FEV ₁ but not symptoms
FEV ₁		22.3/7.2% in low/high quintile SES had poor FEV ₁ (p<0.001)
FEV ₁ , symptoms		Weak associations with both symptoms and FEV ₁
FEV ₁ , symptoms		All indices of SES were associated with both symptoms and FEV ₁
FEV ₁	Smoking	FEV ₁ reduced 10 ml per year without central heating
FEV ₁ , symptoms	Smoking, occupation	Inconsistent associations
FEV ₁ , FVC, symptoms	Smoking	200–300 ml difference in FEV ₁ , symptoms associated with SES
FEV ₁ , FVC, symptoms	Smoking, occupation	Significant association with FEV ₁ and some symptoms
FEV ₁ , FVC, symptoms	Smoking, sex, occupational exposure	COPD OR 4.2
FEV ₁ , quintile	Not adjusted	Spirometric airflow limitation OR 6.3
FEV ₁ , hospital admission	Smoking	15/26% of low SES in low/high quintile of FEV ₁
		300–400 ml difference in FEV ₁ , hospital admission RR 3
Cause-specific mortality	Unadjusted	Chronic bronchitis RR 7.3; other respiratory RR 3.1
Chronic bronchitis		Income associated with chronic bronchitis
FEV ₁ , FVC	Multiple	Boys FEV ₁ and FVC 8% lower, girls no difference

success rate in quitting is also associated with higher social class.^{76 77}

Social class, level of education, and occupation are closely correlated, and some of the social gradient in respiratory diseases may be caused by occupational exposure. The extent of confounding by occupation in epidemiological studies will depend strongly on the spectre of occupational exposure in the population under study. However, several findings indicate that confounding by occupational exposure does not fully explain the association. Women generally do not hold jobs with major exposure to dusts and fumes that potentially cause respiratory disease, yet socioeconomic differences in lung function are found in both sexes,²⁵ although most studies indicate that they are smaller in women.^{22 78 79} Furthermore, socioeconomic differences seem to exist at all ages—that is, they are present before accumulated occupational exposure has had time to cause an impairment of lung function.^{22 25 80} Finally, it was elegantly shown in a Norwegian study that education remained a significant risk factor for symptoms of COPD after adjustment for occupational exposure.²² In this study occupational airborne exposure status was assessed by self-reported past and present exposure to dust or gas as well as by classification of jobs held since leaving school. After adjustment for occupational and smoking exposure, the odds ratio of obstructive lung disease in primary educated versus university educated subjects was 2.9 (1.3–6.5) and the corresponding odds ratio for spirometric airflow limitation (dichotomised) was 5.2 (2.0–13.4).

Damage of alveolar tissue and small airways seems to be mediated by oxidants. Smoking profoundly affects the oxidant/antioxidant balance and increases oxidative stress, but antioxidants in fresh fruit and fish oils may offer some protection against such damage.^{81–89} This may help to explain some of the increased susceptibility related to social class. Antioxidants in wine could hypothetically have a protective effect on the lungs, judging from its effects on overall mortality and lung cancer.^{90 91} However, this association, which would influence the association between socioeconomic status and lung disease, has not so far been addressed in

detail. In fact, a risk factor of marginal importance is heavy alcohol consumption which is associated with airflow limitation.⁹²

Other possible contributory explanations that need consideration include selective drift—that is, that subjects with chronic lung diseases are downwardly mobile. In recent years a number of studies have attempted to clarify these causal relations.^{13 93} The socioeconomic gradient seems to be only slightly affected by selective drift or access to health care.^{94 95} In addition, a gradient is also found for education which, in contrast to income and social class, precedes morbidity.

Conclusion

The socioeconomic gradient in COPD is as great, if not greater, than in any other disease. The impact of socioeconomic status on symptoms, lung function, and other indices of COPD morbidity and mortality is second only to smoking. The association is important because it represents risk factors that are at least partly modifiable. With increasing socioeconomic differences it is becoming increasingly important to treat socioeconomic status in COPD not just as a “nuisance” parameter to adjust for, but as an independent risk factor whose constituents need to be disentangled.

- 1 Speizer FE, Tager IB. Epidemiology of chronic mucus hypersecretion and obstructive airways disease. *Epidemiol Rev* 1979;1:124–42.
- 2 Marmot MG, Shipley MJ, Rose G. Inequalities in death: specific explanations of a general pattern? *Lancet* 1984;i:1003–6.
- 3 Marmot MG, McDowall ME. Mortality decline and widening social inequalities. *Lancet* 1986;ii:274–6.
- 4 Marmot MG, Shipley MJ. Do socioeconomic differences in mortality persist after retirement? 25 year follow up of civil servants from the first Whitehall study. *BMJ* 1996;313:1177–80.
- 5 Marmot M, Feeney A. General explanations for social inequalities in health. *Lancet* 1997;350:207–28.
- 6 Marmot M, Ryff CD, Bumpass LL, et al. Social inequalities in health: next questions and converging evidence. *Soc Sci Med* 1997;44:901–10.
- 7 Department of Health and Social Security. Inequalities in health: report of a working group chaired by Sir Douglas Black. London: DHSS, 1980.
- 8 Kaplan GA, Keil JE. Socioeconomic factors and cardiovascular disease: a review of the literature. *Circulation* 1993;88:1973–98.
- 9 Ingerslev O, Madsen M, Andersen O. Social inequality and mortality in Denmark (in Danish). Sundhedsministeriets Middellevetidsudvalg. Sundhedsministeriet, 1994.
- 10 Osler M, Klebak S. Social deprivation and health in a Danish county. *Scand J Soc Med* 1999;26:289–92.
- 11 Bjerregaard P, Juel K. Life expectancy and mortality in Denmark 1950–1990 (in Danish). København: DIKE, 1993: 1.

- 12 Mackenbach JP, Kunst AE, Cavelaars AEJM, *et al*, the EU Working Group on Socioeconomic Inequalities in Health. Socioeconomic inequalities in morbidity and mortality in western Europe. *Lancet* 1997;**349**:1655-9.
- 13 Kaplan GA, Lynch JW. Editorial: whether studies on the socioeconomic foundations of population health? *Am J Public Health* 1997;**87**:1409-11.
- 14 Stebbings JHJ. Chronic respiratory disease among non-smokers in Hagerstown, Maryland. III. Social class and chronic respiratory disease. *Environ Res* 1971;**4**:213-32.
- 15 Lebowitz MD. The relationship of socio-environmental factors to the prevalence of obstructive lung diseases and other chronic conditions. *J Chronic Dis* 1977;**30**:599-611.
- 16 Cohen BH, Ball WCJ, Brashears S, *et al*. Risk factors in chronic obstructive pulmonary disease (COPD). *Am J Epidemiol* 1977;**105**:223-32.
- 17 Higgins MW, Keller JB, Metzner HL. Smoking, socioeconomic status and chronic respiratory disease. *Am Rev Respir Dis* 1977;**116**:403-10.
- 18 Krzyzanowski M, Jedrychowski W, Wysocki M. Factors associated with the change in ventilatory function and the development of chronic obstructive pulmonary disease in a 13-year follow-up of the Cracow Study. Risk of chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1986;**134**:1011-9.
- 19 Burr ML, Holliday RM. Why is chest disease so common in South Wales? Smoking, social class, and lung function: a survey of elderly men in two areas. *J Epidemiol Community Health* 1987;**41**:140-4.
- 20 Krzyzanowski M, Kauffmann F. The relation of respiratory symptoms and ventilatory function to moderate occupational exposure in a general population. Results from the French PAARC study of 16 000 adults. *Int J Epidemiol* 1988;**17**:397-406.
- 21 Lebowitz MD, Holberg CJ, Martinez FD. A longitudinal study of risk factors in asthma and chronic bronchitis in childhood. *Eur J Epidemiol* 1990;**6**:341-7.
- 22 Bakke PS, Hanoa R, Gulsvik A. Educational level and obstructive lung disease given smoking habits and occupational airborne exposure: a Norwegian community study. *Am J Epidemiol* 1995;**141**:1080-8.
- 23 Hole DJ, Watt GC, Davey Smith G, *et al*. Impaired lung function and mortality risk in men and women: findings from the Renfrew and Paisley prospective population study. *BMJ* 1996;**313**:711-5.
- 24 Demissie K, Ernst P, Hanley JA, *et al*. Socioeconomic status and lung function among primary school children in Canada. *Am J Respir Crit Care Med* 1996;**153**:719-23.
- 25 Prescott E, Lange P, Vestbo J, The Copenhagen City Heart Study Group. Socioeconomic status, lung function and admission to hospital for COPD. *Eur Respir J* 1999 (in press).
- 26 Rasmussen FV, Borchsenius L, Winslow JB. Associations between housing conditions, smoking habits and ventilatory function in men with clean jobs. *Scand J Respir Dis* 1978;**59**:264-76.
- 27 Kerstjens HA, Rijcken B, Schouten JP, *et al*. Decline of FEV₁ by age and smoking status: facts, figures and fallacies. *Thorax* 1997;**52**:820-7.
- 28 Stein CE, Kumaran K, Fall CH, *et al*. Relation of fetal growth to adult lung function in south India. *Thorax* 1997;**52**:895-9.
- 29 Shaheen SO, Barker DJ, Holgate ST. Do lower respiratory tract infections in early childhood cause chronic obstructive pulmonary disease? *Am J Respir Crit Care Med* 1995;**151**:1649-52.
- 30 Shaheen S, Barker DJ. Early lung growth and chronic airflow obstruction (editorial). *Thorax* 1994;**49**:533-6.
- 31 Shaheen SO, Barker DJ, Shiell AW, *et al*. The relationship between pneumonia in early childhood and impaired lung function in late adult life. *Am J Respir Crit Care Med* 1994;**149**:616-9.
- 32 Coggon D, Barker DJ, Inskip H, *et al*. Housing in early life and later mortality. *J Epidemiol Community Health* 1993;**47**:345-8.
- 33 Barker DJ. *The fetal and infant origins of adult disease*. London: BMJ Publications, 1992.
- 34 Barker DJ, Godfrey KM, Fall C, *et al*. Relation of birth weight and childhood respiratory infection to adult lung function and death from chronic obstructive airways disease. *BMJ* 1991;**303**:671-5.
- 35 Barker DJ. The foetal and infant origins of inequalities in health in Britain. *J Public Health Med* 1991;**13**:64-8.
- 36 Barker DJ, Godfrey KM, Fall C, *et al*. Relation of birth weight and childhood respiratory infection to adult lung function and death from chronic obstructive airways disease. *BMJ* 1991;**303**:671-5.
- 37 Barker DJ, Osmond C. Childhood respiratory infection and adult chronic bronchitis in England and Wales. *BMJ* 1986;**293**:1271-5.
- 38 Chan KN, Elliman A, Bryan E, *et al*. Respiratory symptoms in children of low birthweight. *Arch Dis Child* 1989;**64**:1294-304.
- 39 Chan KN, Noble-Jamieson CM, Elliman A, *et al*. Lung function in children of low birthweight. *Arch Dis Child* 1989;**64**:1284-93.
- 40 Seidman DS, Laor A, Gale R, *et al*. Is low birthweight a risk factor for asthma during adolescence? *Arch Dis Child* 1991;**66**:584-7.
- 41 Rona RJ, Gulliford MC, Chinn S. Effects of prematurity and intrauterine growth on respiratory health and lung function in childhood. *BMJ* 1993;**306**:817-20.
- 42 Galdes-Sebaldo M, Sheller JR, Groggaard J, *et al*. Prematurity is associated with abnormal airway function in childhood. *Pediatr Pulmonol* 1989;**7**:259-64.
- 43 Mansell AL, Driscoll JM, James LS. Pulmonary follow-up of moderately low birthweight infants with and without respiratory distress syndrome. *J Pediatr* 1987;**110**:111-5.
- 44 Lechner AJ, Winston DC, Bauman JE. Lung mechanisms, cellularity and surfactant after prenatal starvation in guinea pigs. *J Appl Physiol* 1986;**60**:1610-4.
- 45 Goswami T, Vu ML, Srivastava U. Quantitative changes in DNA, RNA and protein content of the various organs of the young of undernourished female rats. *J Nutr* 1974;**104**:1257-64.
- 46 Faridy EE. Effect of maternal malnutrition on surface activity of fetal lungs in rats. *J Appl Physiol* 1975;**39**:535-40.
- 47 Das RM. The effects of intermittent starvation on lung development in suckling rats. *Am J Pathol* 1984;**117**:326-32.
- 48 Sahebhami H, MacGee J. Effects of starvation on lung mechanics and biochemistry in young and old rats. *J Appl Physiol* 1985;**58**:778-84.
- 49 Kalenga M, Eeckhout Y. Effects of protein deprivation from the neonatal period on lung collagen and elastin in the rat. *Pediatr Res* 1989;**26**:125-7.
- 50 Matsui R, Thurlbeck WM, Fujita Y, *et al*. Connective tissue, mechanical and morphometric changes in the lungs of weanling rats fed a low protein diet. *Pediatr Pulmonol* 1989;**7**:159-66.
- 51 Joseph KS, Kramer MS. Review of the evidence on fetal and early childhood antecedents of adult chronic disease. *Epidemiol Rev* 1996;**18**:158-74.
- 52 Martinez FD, Wright AL, Holberg CJ, *et al*. Maternal age as a risk factor for wheezing lower respiratory illnesses in the first year of life. *Am J Epidemiol* 1992;**136**:1258-68.
- 53 Hanrahan JP, Tager IB, Segal MR, *et al*. The effect of maternal smoking during pregnancy on early infant lung function. *Am Rev Respir Dis* 1992;**145**:1129-35.
- 54 Pullan CR, Hey EN. Wheezing, asthma and pulmonary dysfunction 10 years after infection with respiratory syncytial virus in infancy. *BMJ* 1982;**284**:1665-9.
- 55 Sims DG, Downham MAPS, Gardner PS, *et al*. Study of 8 year-old children with a history of respiratory syncytial virus bronchiolitis in infancy. *BMJ* 1978;**1**:11-14.
- 56 Britton J, Martinez FD. The relationship of childhood respiratory infection to growth and decline in lung function. *Am J Respir Crit Care Med* 1996;**154**:S240-5.
- 57 Jedrychowski W, Krzyzanowski M. The effect of acute broncho-pulmonary infections on the FEV₁ change in 13-year follow-up. The Cracow study. *Eur J Epidemiol* 1990;**6**:20-8.
- 58 Krzyzanowski M, Sherrill DL, Lebowitz MD. Longitudinal analysis of the effects of acute lower respiratory illnesses on pulmonary function in an adult population. *Am J Epidemiol* 1990;**131**:412-22.
- 59 Brunekreef B, Dockery DW, Speizer FE, *et al*. Home dampness and respiratory morbidity in children. *Am Rev Respir Dis* 1989;**140**:1363-7.
- 60 Brunekreef B. Damp housing and adult respiratory symptoms. *Allergy* 1992;**47**:498-502.
- 61 Soliman MY, Rosenstreich D. Natural immunity to dust mites in adults with chronic asthma. *Am Rev Respir Dis* 1986;**134**:962-8.
- 62 Viegi G, Paoletti P, Carrozzi L, *et al*. Effects of home environment on respiratory symptoms and lung function in a general population sample in north Italy. *Eur Respir J* 1991;**4**:580-6.
- 63 Comstock GW, Meyer MB, Helsing KJ, *et al*. Respiratory effects of household exposures to tobacco smoke and gas cooking. *Am Rev Respir Dis* 1981;**124**:143-8.
- 64 Britten N, Davies JMC, Colley JRT. Early respiratory experience and subsequent cough and peak expiratory flow rate in 36 year old men and women. *BMJ* 1987;**294**:1317-20.
- 65 Strachan DP. Damp housing and childhood asthma: validation of reporting of symptoms. *BMJ* 1988;**297**:1223-6.
- 66 Speizer FE, Ferris BG Jr, Bishop YM, *et al*. Respiratory disease rates and pulmonary function in children associated with NO₂ exposure. *Am Rev Respir Dis* 1980;**121**:3-10.
- 67 Melia RJW, Florey CD, Chinn S. The relation between respiratory illness in primary school children and the use of gas for cooking. I. Results from a national survey. *Int J Epidemiol* 1979;**8**:333-8.
- 68 Florey CD, Melia RJW, Chinn S. The relation between respiratory illness in primary school children and the use of gas for cooking. III. Nitrogen dioxide, respiratory illness and lung infection. *Int J Epidemiol* 1979;**8**:347-53.
- 69 Keller MD, Lanese RR, Mitchell RI, *et al*. Respiratory illness in households using gas and electricity for cooking. I. Survey of incidence. *Environ Res* 1979;**19**:495-503.
- 70 Keller MD, Lanese RR, Mitchell RI, *et al*. Respiratory illness in households using gas and electricity for cooking. II. Symptoms and objective findings. *Environ Res* 1979;**19**:504-15.
- 71 Wright AL, Holberg C, Martinez FD. Relationship of parental smoking to wheezing and non-wheezing lower respiratory tract illnesses in infancy. *J Pediatr* 1991;**118**:207-14.
- 72 Lebowitz MD, Burrows B. Respiratory symptoms related to smoking habits of family adults. *Chest* 1976;**69**:48-50.
- 73 Harlap S, Davies AM. Infant admissions to hospital and maternal smoking. *Lancet* 1974;**i**:529-32.
- 74 Sherrill DL, Martinez FD, Lebowitz MD, *et al*. Longitudinal effects of passive smoking on pulmonary function in New Zealand children. *Am Rev Respir Dis* 1992;**145**:1136-41.

- 75 Gold DR, Wang X, Wypij D, *et al.* Effects of cigarette smoking on lung function in adolescent boys and girls. *N Engl J Med* 1996;**335**:931–7.
- 76 Osler M, Prescott E. Psychosocial, behavioural, and health determinants of successful smoking cessation: a longitudinal study of Danish adults. *Tobacco Control* 1998;**7**:262–7.
- 77 Freund KM, D'Agostino RB, Belanger AJ, *et al.* Predictors of smoking cessation: the Framingham study. *Am J Epidemiol* 1992;**135**:957–64.
- 78 Arber S. Comparing inequalities in women's and men's health: Britain in the 1990s. *Soc Sci Med* 1997;**44**:773–87.
- 79 Koskinen S, Martelin T. Why are socioeconomic mortality differences smaller among women than among men. *Soc Sci Med* 1994;**38**:1385–96.
- 80 Williams DR. Race/ethnicity and socioeconomic status: measurement and methodological issues. *Int J Health Serv* 1996;**26**:483–505.
- 81 Sridhar MK. Nutrition and lung health. *BMJ* 1995;**3**:1075–6.
- 82 Shahar E, Folsom AR, Melnick SL, *et al.* Dietary n-3 polyunsaturated fatty acids and smoking-related chronic obstructive pulmonary disease. Atherosclerosis Risk in Communities Study Investigators. *N Engl J Med* 1994;**331**:228–33.
- 83 Strachan DP, Cox BD, Erzincioğlu SW, *et al.* Ventilatory function and winter fresh fruit consumption in a random sample of British adults. *Thorax* 1991;**46**:624–9.
- 84 Schwartz J, Weiss ST. Relationship between dietary vitamin C intake and pulmonary function in the first national health and nutrition examination survey (NHANES I). *Am J Clin Nutr* 1994;**59**:110–4.
- 85 Shahar E, Folsom AR, Melnick SL, *et al.* Does dietary vitamin A protect against airway obstruction? *Am J Respir Crit Care Med* 1994;**150**:978–82.
- 86 Sharp DS, Rodriguez BL, Shahar E, *et al.* Fish consumption may limit the damage of smoking on the lung. *Am J Respir Crit Care Med* 1994;**150**:983–7.
- 87 Schwartz J, Weiss ST. Dietary factors and their relationship to respiratory symptoms: NHANES II. *Am J Epidemiol* 1990;**130**:67–76.
- 88 Britton J, Pavord I, Richards K, *et al.* The effects of dietary antioxidants on lung function in the general population. *Am Rev Respir Dis* 1993;**147**:369.
- 89 Meidema I, Feskens EJM, Heederik D, *et al.* Dietary determinants of long-term incidence of chronic nonspecific lung diseases. The Zutphen study. *Am J Epidemiol* 1993;**138**:37–45.
- 90 Grønbaek M, Deis A, Sørensen TIA, *et al.* Mortality associated with moderate intakes of wine, beer, or spirits. *BMJ* 1995;**310**:1165–9.
- 91 Prescott E, Grønbaek M, Becker U, *et al.* Alcohol and lung cancer: influence of type of alcoholic beverage. *Am J Epidemiol* 1999;**149**:463–70.
- 92 Lange P, Groth S, Mortensen J, *et al.* Pulmonary function is influenced by heavy alcohol consumption. *Am Rev Respir Dis* 1988;**137**:1119–23.
- 93 Lynch JW, Kaplan GA, Cohen RD, *et al.* Do cardiovascular risk factors explain the relation between socioeconomic status, risk of all-cause mortality, cardiovascular mortality, and acute myocardial infarction? *Am J Epidemiol* 1996;**144**:934–42.
- 94 Møller L, Kristensen TS, Hollnagel H. Social class and cardiovascular risk factors in Danish men. *Scand J Soc Med* 1991;**19**:116–26.
- 95 Blane D, Davey Smith G, Bartley M. Social selection: what does it contribute to social class differences in health? *Soc Sci Health Illness* 1993;**15**:1–15.