Home treatment of COPD exacerbations

Dirkje S Postma, Nick H T Ten Hacken, Huib A M Kerstjens, Gerard H Koëter

Department of Pulmonology, University Hospital, Groningen, The Netherlands

Introductory article

Home treatment of exacerbations of chronic obstructive pulmonary disease by an acute respiratory assessment service

J H Gravi, O A Al-Rawas, M M Cotton, U Flanigan, A Irwin, R D Stevenson

Background. Exacerbations of chronic obstructive pulmonary disease are a major cause of hospital admissions, but do not require intensive investigation or complex therapy. We investigated the suitability of home care for severe uncomplicated exacerbations. **Methods.** Over 3–5 years we assessed 962 patients with exacerbations of chronic obstructive pulmonary disease after referral to a hospital respiratory department by their family physicians. All patients had chest radiographs, oxygen-saturation or arterial-gas analysis, spirometry, and physical assessment. Unless admission was thought to be essential, patients were allowed home with a customised treatment package. Each patient was visited daily by a respiratory nurse who monitored progress and treatment compliance and provided education and reassurance. **Findings.** 145 (15%) of 962 required admission at initial referral and 115 (12%) were admitted later. 653 (68%) patients were managed entirely at home and 49 (5%) were referred inappropriately. One patient died at home. All patients had severe disease with a mean forced expiratory volume in 1 s of 1.02 l and 395 (41%) had required hospital admission in the previous year. **Interpretation.** After formal assessment in a hospital respiratory unit, many patients with exacerbations of chronic obstructive pulmonary disease can be treated at home by respiratory nurses. (Lancet 1998;351:1853–5)

Respiratory diseases are a major cause of morbidity and mortality in the community. Patients with moderate to severe chronic obstructive pulmonary disease (COPD) are prone to exacerbations and the frequency of these exacerbations increases with the severity of COPD.¹ Exacerbations are the main cause of visits to general practitioners and emergency departments, and of hospital admissions in affluent countries. Patients with COPD have been reported to have on average 1–4 exacerbations per year, and some of these exacerbations lead to hospitalisation.^{2,3} Hospital admissions for airway diseases such as asthma and COPD account for about 7% of all admissions to hospital.⁴ In addition, hospital mortality due to an acute exacerbation of COPD is approximately 10%, thereby exceeding current hospital mortality rates for myocardial infarction. The long term outcome after admission to hospital for an exacerbation is rather poor, with 40% mortality in one year.⁵ These figures vary from country to country depending on the health care system and the availability of ICU beds. Overall, the economic burden for society and the personal burden to an individual patient are very high. It is therefore not surprising that studies have been initiated to pay more attention to adequate treatment of exacerbations in and out of hospital, to improvement of management at home and to the prevention of these exacerbations. The article by Gravi and coworkers is an example of the increasing attention that is now being paid to the management of COPD.⁶

One of the best strategies for preventing admission to hospital is prevention of the exacerbation itself. In this review we will pay attention to the definition of an exacerbation, the causes and consequences of exacerbations, and the management, including available knowledge on prevention. We will try specifically to address the issue of whether or not exacerbations would be better treated at home.

Definition of exacerbation

Recent guidelines do not give a definition of exacerbations nor do they provide a classification of exacerbations of COPD based on clinical parameters.^{7,8} Although many classifications in the literature can be useful for clinical studies, they are not standardised and are rather difficult to use in clinical practice, especially for general practitioners in the home situation.

Mild exacerbations of COPD are generally accepted to represent an increase in symptoms, especially dyspnoea, not necessarily accompanied by increased cough and sputum production which may be more tenacious than usual. The increased symptomatology forces the patient to seek medical attention.⁹ A severe exacerbation is associated with acute respiratory failure especially in patients with impaired lung function, sometimes accompanied by CO₂ retention.^{10,11} It is clear that the latter cannot be easily diagnosed in the general practice setting and it cannot be treated in the home situation.

Table 1 Assessment of severity of exacerbation

Measurements at home and in hospital	Measurements in hospital
Knowledge of provoking factors	FEV ₁ <1 litre
History	Pao ₂ <8 kPa, Sao ₂ <90%
Previous condition	Paco ₂ ≥6 kPa
Duration/progression	Chest radiography
Sleeping/eating difficulties	White blood cell count
Symptoms	Haemoglobin
Change in cough	Sputum stain/culture
Change in sputum	Biochemistry
Increase in dyspnoea at rest	Electrocardiogram
Inability to speak one full sentence	
Temperature >38.5°C	
Ankle oedema	
Respiratory rate >25/min	
Heart rate >110/min	
Worsening cyanosis	
Use of accessory muscles	
Loss of alertness	

Factors which determine the choice between home management or inpatient care other than the severity of the exacerbation are the cause of the exacerbation (for example, severe pneumonia), concomitant diseases, and/or social factors.

Assessments of severity of exacerbation

Proper management of an acute exacerbation requires information on the previous condition of the patient when stable. Next, a precise assessment of the severity of the current exacerbation is needed. Table 1 shows the assessment of severity of an exacerbation.

PROVOKING FACTORS

Before assessing the severity of an exacerbation one must first consider the provoking factors (table 2). Common causes of acute exacerbations of COPD are thought to be primary infections (often viral) of the tracheobronchial tree. Their role has been discussed at length in recent years.^{11–19} There is general agreement that the major pathogens isolated from sputum during acute exacerbations of COPD are *Haemophilus influenzae*, *Streptococcus pneumoniae* and *Moraxella catarrhalis*.¹⁸ Several studies have now reported that these bacteria are also isolated from patients with COPD in a stable state. The mere presence of bacterial colonisation of the lower airway does not confirm the importance of bacterial infection in the pathogenesis of COPD exacerbations. It might also be a marker of disease severity since it has been shown that there is an association between bacterial colonisation of the lower airway and more severe airway obstruction.¹⁷ It may reflect a balance in which the impaired host defences are able to limit the numbers of bacteria but not eradicate

Table 2 Provoking and aggravating factors of an acute exacerbation of COPD

Bacterial infection	Inappropriate oxygen administration
Viral infection	Stopping treatment
Pneumonia	Drugs (tranquillisers, diuretics, etc)
Heart failure	Poor nutrition
Pulmonary embolus	Respiratory fatigue of the muscles
Pneumothorax	Gastro-oesophageal reflux and/or aspiration
Air pollution	Airway plugging due to excessive mucus production or inadequate expectoration

them.¹⁸ Thus, the role of bacterial infections in exacerbations of COPD remains controversial. A meta-analysis showed a minor but significant benefit from antibiotics which led to general advice to use antibiotics for exacerbations.¹⁹ However, a recent negative study carried out in general practice has weakened even further the very cautious recommendation for antibiotic use.¹² The role of viral infection in COPD is also still under scrutiny. The contribution of viral infections to COPD exacerbations has been variable,¹⁵ ranging from 4.4% to 63.3% depending on the population under study, the severity of exacerbation, and whether it is in hospital or general practice. Furthermore, only a few studies have used viral serology to diagnose the viral infection. Improved techniques for detection of viral infections will probably change these figures. Nevertheless, influenza vaccination is still recommended for patients with COPD since influenza is the most important cause of excess mortality.¹⁶

Other factors associated with exacerbations of COPD include pneumonia, pulmonary hypertension, heart failure, pulmonary embolism, and pneumothorax.²⁰ Outdoor air pollution has also been shown recently to increase the hospital admission rates.²¹ Poor nutrition, drugs (especially tranquillisers), aspiration, airway plugging by mucus, and respiratory muscle fatigue, especially in patients with malnutrition, may aggravate the exacerbation. Furthermore, stopping regular medication such as diuretics and/or bronchodilators at the patients own initiative may increase symptoms. Inappropriate oxygen administration may aggravate an exacerbation since many COPD patients with hypercapnia as well as hypoxia deteriorate when hypoxic respiratory drive is reduced by high oxygen administration.²⁰

PATIENT HISTORY

This is the most important means for deciding on the severity and causation of the exacerbation. The history allows comparison of the current status with the stable state and assessment of whether there has been a gradual increase in symptoms or an acute event. It may also help in diagnosis and management if the patient recognises the symptom pattern from previous experience and response to treatment.

Assessment of the severity of an exacerbation is also helped by detecting sleeping and/or eating difficulties as these also indicate a serious situation.

PHYSICAL SIGNS

Signs that indicate severity include inability to speak one sentence in full, central cyanosis, use of accessory respiratory muscles, paradoxical chest wall movements, inability to lie flat, oedema, and ultimately haemodynamic instability (table 1). A body temperature of >38.5°C, a respiratory rate of >25 breaths per minute, and a heart rate of 110 beats per minute or more are arbitrary cut off points taken to indicate a severe exacerbation.¹¹ It is questionable whether these are the correct parameters and levels of abnormality for assessment of severity. For instance, a recent study showed that mean body temperature at admission for a severe exacerbation was 36.4°C.¹⁰ In the introductory article on the management of COPD exacerbations to be discussed in this review Gravit *et al*⁶ did not use temperature in their algorithm for home or hospital management which may be correct.

CLINICAL INVESTIGATIONS

Blood gas analysis provides further evidence for the severity of an exacerbation.¹¹ Arterial oxygen tension (PaO_2) of <8.0 kPa (60 mm Hg) and/or arterial oxygen saturation (SaO_2) of $<90\%$ when breathing air indicate respiratory failure and the need for evaluation and treatment. Changes must, where possible, be interpreted in conjunction with baseline values for an individual patient. Generally, however, a PaO_2 below 6.7 kPa (50 mm Hg), PaCO_2 above 9.3 kPa (70 mm Hg), and pH <7.3 signifies a life threatening episode possibly requiring ICU management. Gravil *et al*⁶ did not use SaO_2 as a parameter for assessing the severity of an exacerbation. Whether this influences their results is not certain since the effect of SaO_2 on re-admission was not evaluated.

In a home based setting a general practitioner cannot use blood gas analysis and has to rely on cyanosis. However, this is not a reliable estimate of hypoxaemia. When present, and new for the particular patient, this signifies severity. When absent or unnoticed, hypoxaemia may still be present and the severity of the exacerbation will be underestimated. The European Respiratory Society guidelines therefore advise that, if the severity of an episode is in doubt, assessment should take place in hospital.⁸

Other aggravating factors should be sought such as congestive heart failure or pneumonia.²⁰ Thus, chest radiography is necessary when considering a severe exacerbation. Furthermore, sputum bacteriology may help, the white blood cell count and differential cell count may point to a viral or bacterial infection, and a high haemoglobin measurement may suggest long-standing hypoxaemia. An electrocardiogram (ECG) may help in establishing right ventricular hypertrophy, arrhythmias or ischaemia and even pulmonary embolism, especially if a previously normal ECG is available.

Lung function measurements are also recommended in guidelines on the assessment of the severity of an exacerbation.^{7,8} Airways obstruction is the *conditio sine qua non* in the definition of COPD, and obstruction is the factor that is best related to disability and mortality. For instance, in patients with a forced expiratory volume in one second (FEV_1) of <0.75 l the mortality rate at one year is approximately 30%, rising to 90% at 10 years.²² Fortunately, airways obstruction as assessed by FEV_1 is a very reliable measurement. Airway obstruction is measured as a reduction in FEV_1/FVC ratio or as a reduced FEV_1 to slow vital capacity ratio. Measurement of FEV_1 and VC are increasingly accessible to non-respiratory physicians. However, in many individuals with a severe exacerbation spirometric testing may not be possible. Also, it is not clear how far the level of reduction in FEV_1 signifies severity of an exacerbation or its outcome. Some studies have shown that mild exacerbations are not accompanied by a reduction in FEV_1 ²³ while others found decreased FEV_1 in both mild exacerbations¹² and moderate to severe exacerbations.¹³ A recent study showed that FEV_1 levels during an exacerbation did not relate to re-admission within 12 months after the first admission to hospital with an exacerbation of COPD.²⁴

SUBJECTIVE MEASURES

Doctors react to several components of disease severity. Subjectively, they assess frailty and mental state. Whether it is correct to use these criteria for making a decision to either let a patient stay at home or refer to

Figure 1 Exacerbation scores for patients with COPD who were admitted immediately to the hospital because of an exacerbation, referred home for treatment and admitted afterwards to the hospital, or referred home for treatment and staying home. Reproduced from Gravil *et al*⁶ with permission.

hospital is questionable. These will depend on interpretation and commitment of the general practitioner. The same is true for doctors and nurses working in the outpatient clinic. Gravil *et al*⁶ used the degree of disability or frailty in their algorithm for deciding whether or not to admit a patient to hospital in their study on exacerbation assessment by nurses and doctors at the outpatient clinic. They used these even though the grading of severity of the exacerbation was based on respiratory rate, heart rate, PaO_2 , abnormal chest radiograph, serious co-morbid condition, altered mental state, and living alone. Unfortunately, the score used did not predict for whether patients were likely to be re-admitted after being sent home—53% of those without an exacerbation and 61.7% of those re-admitted when first sent home had a score of 1 whereas 87.6% of those admitted immediately had this level of severity, mean score values being 0.77, 0.92, and 2.11, respectively. When looking at figure 1 in their article (fig 1), those sent home and re-admitted were indiscernible with respect to their severity score, but only those admitted immediately had scores of 4 or more. There is room for improvement of severity scores in this respect, and the use of this type of severity score in clinical practice is not yet advised.

Predictive factors for hospital admission for exacerbations of COPD

Some patients are prone to frequent exacerbations which have considerable impact on their daily living activities. It is not clear, however, which factors determine the development and severity of an exacerbation. Predictors for admission to hospital are variously reported in the available studies. Some studies have assessed the effect of treatment on the reduction of frequency of hospital admissions. Hospital needs are reduced when home oxygen is instituted in patients with hypoxaemia.²⁵ One

retrospective study showed that basal body weight, decline of FEV₁ and the rate of deterioration of blood gas tensions were related to the necessity for admission to ICU.²⁶ However, the study was retrospective in nature and included only 16 individuals who were all chronically hypercapnic. A recent prospective study investigated 64 patients with moderate to severe COPD with mean FEV₁ values of 39 (20)% predicted.²⁷ Patients were all followed with three monthly visits and admitted to hospital when they had an acute exacerbation defined by increased symptoms of dyspnoea, cough, or sputum production plus one or more of the following: inadequate response of symptoms to outpatient management which consisted of an increase in the daily doses of inhaled bronchodilators, antibiotics, and a short course of oral steroids; inability to walk between rooms (patient previously mobile); inability to eat or sleep due to dyspnoea; high risk co-morbidity, either pulmonary (e.g. pneumonia) or non-pulmonary; prolonged (>15 days) progressive symptoms before emergency visit; altered mental state; worsening hypoxaemia; new or worsening hypercapnia. Furthermore, patients were admitted to hospital when co-morbid conditions worsened the pulmonary symptoms or when they had unresponsive cor pulmonale. Twenty nine patients were admitted to hospital for an exacerbation, 62% of these because of worsening hypoxaemia, 59% because of an inadequate response to outpatient management, and 52% with severe dyspnoea (inability to walk, eat or sleep); 66% of these subjects had three or more of the listed criteria for hospital admission. The conclusion of the study was that only PaCO₂ and mean pulmonary arterial pressure were independently related to admission to hospital. Markers of airflow limitation (such as FEV₁) and hypoxaemia were not retained as independent predictors of hospitalisation. However, it is possible that this cohort was too small to detect any influence of the latter two factors on the risk of hospital admission. For instance, a PaO₂ below 8.0 kPa (60 mm Hg) was only present in 15 patients, and even then it was already corrected by oxygen therapy in many individuals. Thus, the best predictors of mortality in COPD—namely, the severity of airflow limitation, advancing age and hypoxaemia²²—were not of use for predicting hospital admission. However, as already mentioned, the study group was small. Notwithstanding this, hypercapnia was an important predictor, as expected, yet only 13 patients had a PaCO₂ higher than 6.66 kPa (50 mm Hg). Hypercapnia has also been shown to be a good predictor of mortality after discharge from the ICU in patients older than 65 years. Interestingly, Costello *et al*²⁸ have shown that many patients admitted to the ICU with hypercapnia associated with an exacerbation normalised their hypercapnia. However, the patients who remained hypercapnic had a worse long term prognosis than those with reversible hypercapnia.

The best independent predictor for admission to hospital in the study by Kessler *et al*²⁷ was pulmonary hypertension, a marker of the deleterious effects of longstanding alveolar hypoxia on the pulmonary circulation. The authors speculated that home mechanical ventilation might reduce the number of hospital admissions by reducing hypercapnia, as was shown by Leger *et al* in a previous study with supervision of nasal intermittent positive pressure ventilation.²⁹ However, there are also negative studies of home mechanical ventilation and its role is beyond the scope of this review. Home care supervision programmes alone have also been shown to be successful in reducing the need for hospitalisation in these patients.³⁰

One study has looked into risk factors predisposing to frequent exacerbations in patients with moderately severe COPD and an FEV₁ of 40 (19)% predicted.³¹ The results showed that daily cough or wheeze and bronchitic symptoms increased the risk for frequent exacerbations (3–8 in one year follow up) but not daily sputum production or dyspnoea. In a multivariate analysis patients with frequent past exacerbations, daily wheeze, and bronchitic symptoms were at the highest risk for frequent exacerbations. Moreover, a worse disease specific health status at the last clinic visit in a stable state, assessed with the St George's Respiratory Questionnaire (SGRQ), was strongly related to a higher number of exacerbations over the one year study period. The authors found no relationship between the SGRQ outcome and the frequency of exacerbations and the need for hospital admission. One recent study found a relationship between poor quality of life and re-admission to hospital.²⁴ Reasons for admissions to hospital are complex and patients are frequently admitted late in the evolution of the disease, which affects quality of life to a large extent. Nevertheless, it seems reasonable to anticipate that a reduction in the frequency of exacerbations may lead to improvement in quality of life in individuals with COPD. It has been shown that poor quality of life is associated with greater use of health care resources³² and that, when quality of life improves even though lung function does not improve, consultations and hospital admissions decrease.³³ Whether improvement in quality of life will also lead to a reduction in exacerbations is, however, questionable.

Further studies are needed to determine the extent to which nurse support and education may contribute to recovery or prevention of exacerbations. This effect is uncertain, but support at a moment of uncertainty for both the patient and his environment would seem important. Appropriate pharmacological treatment and oxygen therapy may further enhance the outcome of COPD exacerbations, which can also be assisted by nurses. The study by Gravit *et al*⁶ is the first to assess whether home treatment, in association with supervision and training by nurses, may be beneficial for treatment of COPD exacerbations. Many practitioners need to decide between home and hospital management of COPD exacerbations on a daily basis, and further studies may help us decide which is the best choice and for which patients to reduce the number of hospitalisations with better outcomes and lower costs.

Management of an exacerbation

OUTPATIENT MANAGEMENT

The objectives of outpatient or home management have been formulated by Siafakas and Bouros²⁰ as follows: to treat infection, if present; to remove excess bronchial secretions; to increase airflow; to improve respiratory muscle strength and thus facilitate cough; to avoid adverse events of treatment; to educate patients and their families on the signs of deterioration and the actions to be taken (written instruction). The European guidelines stress that it is important to monitor the patient with a moderate to severe exacerbation and to reassess the effects of treatment within 48 hours.⁸ The education and instructions to the patient and/or family are stressed.

In general, exacerbations are treated by institution or increase of bronchodilators, systemic corticosteroids, and oxygen when necessary. If infection is likely—for instance, in the presence of fever—antibiotics are started. Otherwise, following observation for 24–48

LEARNING POINTS

- * Exacerbations of COPD are a major cause of hospital admissions, yet the individual risk factors for hospital admission are not clearly known.
- * The role of nurse support and education in prevention of COPD exacerbations should be further studied.
- * Antibiotics are frequently prescribed in COPD but their usefulness in mild to moderate exacerbations has not been proved.
- * Frequent prescription of antibiotics induces β -lactamase resistance of bacteria in sputum in patients with COPD.

hours, antibiotics can be started if there is further deterioration. When a patient is known to have severe COPD with hypoxaemia, early antibiotics are appropriate. However, studies have not been done to assess whether this is the best strategy.

Short acting β agonists are generally the choice of bronchodilator and their frequency and dosage should be increased to a maximum. Anticholinergics are recommended and may offer greater benefit in combination with a β agonist. Their use during an exacerbation can be more efficacious when used via a spacer. Aminophylline use remains controversial for exacerbations and some studies have shown that the toxicity of β agonists is increased when combined with aminophylline, but not the efficacy.³⁴

A recent study in an outpatient setting has shown that oral corticosteroids may accelerate the recovery from an exacerbation, especially with hypoxaemia, and that they may improve spirometric parameters and reduce the treatment failure rate.⁹ Moreover, a study by Paggiaro and coworkers³⁵ and the recent results of the ISOLDE study show that inhaled corticosteroids may reduce the number of exacerbations. The same may be true for antioxidants. These may produce benefit by better clearance of sputum. This may be important since mucus production has been shown to be a risk factor for mortality due to pulmonary infections.³⁶

Initiation of oxygen therapy at home is unwarranted. It may lead to serious complications and must be avoided. If the patient is on long term oxygen therapy a thorough inspection of the apparatus and review of the dosage of oxygen given is recommended. This aspect requires further clinical evaluation.

Recent studies have advocated education as a measure to reduce consumption of health care facilities. Tougaard *et al.*³⁷ showed that this is the case in a group of patients with COPD of variable severity. Education on actions to be taken for an incipient exacerbation should include: (1) optimal compliance with medication; (2) avoidance of hypnotics and sedatives; (3) sufficient fluid intake; (4) small and frequent meals if eating is difficult; (5) self-clearance of sputum by adequate cough techniques (in cases of emphysema the forced expiration technique is not advised³⁸). The patient and family should have written instructions on the symptoms and signs indicating a worsening condition and the actions to be taken. In this respect there is a note of caution on prescription of antibiotics. A study in patients with COPD showed that increasing the number of antibiotic courses was associated with a higher incidence of β -lactamase producing bacteria.³⁹ Further, the number of COPD patients in this study admitted to hospital for an exacerbation of COPD was higher in the group who had β -lactamase positive strains of *Haemophilus*

influenzae and *Moraxella catarrhalis* in their sputum than in the group with β -lactamase negative strains. Antibiotics are probably instituted too frequently during exacerbations.

Referral or home care?

As with many other diseases, it is difficult to give a simple guide as to which patients with an exacerbation should be seen by the general physician, which should be seen at an emergency department and sent home, and which should be admitted to hospital. These decisions will probably vary by country and, unfortunately, also by billing systems. In general it is important to realise that criteria for patterns of care should be evidence based from controlled studies. So far there is little evidence on which to base the decision process. The best advice remains: when in doubt refer to an emergency department. When there is significant deterioration in mental status, increased use of accessory respiratory muscles, and paradoxical abdominal breathing referral is mandatory since any of these symptoms suggests respiratory failure.

Since the staging of exacerbation severity is still not formalised and does not give a clear indication of the best treatment, we still need more prospective studies in patients with COPD over a range of severity. This includes the large group of patients under the care of general practitioners, as well as more severe cases with shared care between general practitioners and specialists. Finally, the role of education has to be further explored. If the reduction in hospital use, as shown by Gravil and coworkers,⁶ can be translated into a reduction in the numbers of exacerbations, there will be important consequences for COPD patients, their families, nurses and doctors, and for the health economy.

- 1 Fletcher C, Peto R. The natural history of chronic airflow obstruction. *BMJ* 1977;1:1645-8.
- 2 Collet JP, Shapiro S, Ernst P, *et al.* Effect of an immunostimulating agent on acute exacerbations and hospitalization in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1997;156:1719-24.
- 3 Anthonisen NR, Manfreda J, Warren CPW, *et al.* Antibiotic therapy in exacerbations of chronic obstructive pulmonary disease. *Ann Intern Med* 1987;106:196-204.
- 4 Angus RM, Murray S, Lay JW, *et al.* Management of chronic airflow obstruction: differences in practice between respiratory and general physicians. *Thorax* 1994;88:493-7.
- 5 Connors AF, Dawson NV, Thomas C, *et al.* Outcome following acute exacerbation of severe chronic obstructive lung disease. *Am J Respir Crit Care Med* 1996;154:959-67.
- 6 Gravil JH, Al-Rawas OA, Cotton MM, *et al.* Home treatment of exacerbations of chronic obstructive pulmonary disease by an acute respiratory assessment service. *Lancet* 1998;351:1853-5.
- 7 American Thoracic Society. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease: ATS statement. *Am J Respir Crit Care Med* 1995;152:S77-S120.
- 8 Siafakas NM, Vermeire P, Pride NB, *et al.* Optimal assessment and management of chronic obstructive pulmonary disease (COPD). A consensus statement of the European Respiratory Society. *Eur Respir J* 1995;8:1398-420.

- 9 Thompson WH, Nielson CP, Carvalho P, *et al*. Controlled clinical trial of oral prednisone in outpatients with acute COPD exacerbation. *Am J Respir Crit Care Med* 1996;154:407–12.
- 10 Connors AF, Dawson NW, Thomas C, *et al* for the SUPPORT Investigators. Outcome following acute exacerbations of severe chronic obstructive lung disease. *Am J Respir Crit Care Med* 1996;154:959–67.
- 11 Hagedorn SD. Acute exacerbations of COPD. How to evaluate severity and treat the underlying cause. *Postgrad Med* 1992;91:105–12.
- 12 Sachs AP, Koëter GH, Groenier KG, *et al*. Changes in symptoms, peak expiratory flow and sputum flora during treatment with antibiotics of exacerbations in patients with chronic obstructive pulmonary disease in general practice. *Thorax* 1995;50:758–63.
- 13 Fletcher CM, Peto R, Tinker C, *et al*. The natural history of chronic bronchitis and emphysema: an eight-year study of early chronic obstructive lung disease in working men in London. New York: Oxford University Press, 1976: 272.
- 14 Wilson R, Dowling RB, Jackson AD. The biology of bacterial colonization and invasion of the respiratory mucosa. *Eur Respir J* 1996; 9:1523–30.
- 15 Schlick W. Selective indications for use of antibiotics: when and what? *Eur Respir Rev* 1992;9:187–92.
- 16 Rothbarth PH, Kempen BM, Sprenger JW. Sense and nonsense of influenza vaccination in asthma and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1995;151:1682–6.
- 17 Zalacain R, Sobradillo V, Amilibia J, *et al*. Predisposing factors to bacterial colonization in chronic obstructive pulmonary disease. *Eur Respir J* 1998;13:343–8.
- 18 Wilson R. Bacterial infection and chronic obstructive pulmonary disease. *Eur Respir J* 1998;13:233–5.
- 19 Saint S, Bent S, Vittinghoff E, *et al*. Antibiotics in chronic obstructive pulmonary disease exacerbations: a meta-analysis. *JAMA* 1995;273: 957–60.
- 20 Siafakas NM, Bouros D. Management of acute exacerbation of chronic obstructive pulmonary disease. In: *Management of chronic obstructive pulmonary disease, A European Respiratory Monograph*. Vol 3. 1998: 264–277.
- 21 Stedman JR, Anderson HR, Atkinson RW, *et al*. Emergency hospital admissions for respiratory disorders attributable to summer time ozone episodes in Great Britain. *Thorax* 1997;52:958–63.
- 22 Hodgkin JE. Prognosis in chronic obstructive pulmonary disease. *Clin Chest Med* 1990;11:555–69.
- 23 Saetta M, Di Stefano A, Maestrelli P, *et al*. Airway eosinophilia in chronic bronchitis during exacerbations. *Am J Respir Crit Care Med* 1994;150:1646–52.
- 24 Osman LM, Godden DJ, Friend JAR, *et al*. Quality of life and hospital re-admission in patients with chronic obstructive pulmonary disease. *Thorax* 1997;52:67–71.
- 25 Petty TL. Long-term outpatient oxygen-therapy. In: Petty TL, ed. *Chronic obstructive pulmonary disease*. New York: Marcel Dekker, 1985: 375–88.
- 26 Vitacca M, Cline E, Porta R, *et al*. Acute exacerbation in patients with COPD: predictors of need for mechanical ventilation. *Eur Respir J* 1996;9:1487–93.
- 27 Kessler R, Faller M, Fourgaut G, *et al*. Predictive factors of hospitalization for acute exacerbation in a series of 64 patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;159:158–64.
- 28 Costello R, Deagen P, Fitzpatrick M, *et al*. Reversible hypercapnia in chronic obstructive pulmonary disease: a distinct pattern of respiratory failure with a favorable prognosis. *Am J Med* 1997;102:239–44.
- 29 Leger P, Bedicam JM, Cornette A, *et al*. Nasal intermittent positive pressure ventilation: long-term follow-up in patients with severe chronic respiratory insufficiency. *Chest* 1994;105:100–5.
- 30 Cline E, Vitacca M, Foglio K, *et al*. Long-term home care programmes may reduce hospital admissions in COPD with chronic hypercapnia. *Eur Respir J* 1996;9:1605–10.
- 31 Seemungal TAR, Donaldson GC, Paul EA, *et al*. Effect of exacerbation on quality of life in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;157:1418–22.
- 32 Traver GA. Measures of symptoms and life quality to predict emergent use of institutional health care resources in chronic obstructive airway disease. *Heart Lung* 1988;17:689–97.
- 33 Cox NJ, Hendricks JC, Binkhorst RA, *et al*. A pulmonary rehabilitation program for patients with asthma and mild chronic obstructive pulmonary diseases (COPD). *Lung* 1993;171:235–44.
- 34 Rice KL, Leatherman JW, Duane PG, *et al*. Aminophylline for acute exacerbations of chronic obstructive pulmonary disease. *Ann Intern Med* 1987;107:305–9.
- 35 Paggiaro PL, Dahle R, Bakran L, *et al*. Multicenter randomized placebo controlled trial of inhaled fluticasone propionate in patients with COPD. *Lancet* 1998;351:773–80.
- 36 Prescott E, Lange P, Vestbo J. Chronic mucus hypersecretion in COPD and death from pulmonary infection. *Eur Respir J* 1995;8:1333–8.
- 37 Tougaard L, Krone T, Sorknaes A, *et al* and the PASTMA group. Economic benefits of teaching patients with chronic obstructive pulmonary disease about their illness. *Lancet* 1992;339:1517–20.
- 38 Van der Schans CP, Piers DA, Beekhuis H, *et al*. Effect of forced expirations on mucus clearance in patients with chronic airflow obstruction: effect of lung recoil pressure. *Thorax* 1990;45:623–7.
- 39 Sportel JH, Koëter GH, Van Altena R, *et al*. Relation between beta-lactamase producing bacteria and patient characteristics in chronic obstructive pulmonary disease (COPD). *Thorax* 1995;50:249–53.