Decreased bronchodilating effect of salbutamol in relieving methacholine induced moderate to severe bronchoconstriction during high dose treatment with long acting β_2 agonists

H J van der Woude, T H Winter, R Aalbers

Abstract

Background—In vitro the long acting β_2 agonist salmeterol can, in contrast to formoterol, behave as a partial agonist and become a partial antagonist to other β_2 agonists. To study this in vivo, the bronchodilating effect of salbutamol was measured during methacholine induced moderate to severe bronchoconstriction in patients receiving maintenance treatment with high dose long acting β_2 agonists.

Methods—A randomised double blind crossover study was performed in 19 asthmatic patients with mean forced expiratory volume in one second (FEV₁) of 88.4% predicted and median concentration of methacholine provoking a fall in FEV₁ of 20% or more (PC₂₀) of 0.62 mg/ml at entry. One hour after the last dose of 2 weeks of treatment with formoterol (24 μ g twice daily by Turbuhaler), salmeterol (100 μ g twice daily by Diskhaler), or placebo a methacholine provocation test was performed and continued until there was at least a 30% decrease in FEV₁. Salbutamol (50 μ g) was administered immediately thereafter, followed by ipratropium bromide (40 μ g) after a further 30 minutes. Lung function was monitored for 1 hour after provocation.

Results—There was a significant bronchodilating and bronchoprotective effect after 2 weeks of active treatment. The dose of methacholine needed to provoke a fall in FEV₁ of $\geq 30\%$ was higher after pretreatment with formoterol (2.48 mg) than with salmeterol (1.58 mg) or placebo (0.74 mg). The difference between formoterol and salmeterol was statistically significant: 0.7 doubling dose steps (95% CI 0.1 to 1.2, $p=0.016$). The immediate bronchodilating effect of subsequently administered salbutamol was significantly impaired after pretreatment with both drugs ($p<0.0003$ for both). Three minutes after inhaling salbutamol the increase in FEV₁ relative to the pre-methacholine baseline was 15.8%, 7.3%, and 5.5% for placebo, formoterol and salmeterol, respectively (equivalent to increases of 26%, 14%, and 12%, respectively, from the lowest FEV₁ after methacholine). At 30 minutes significant differences remained, but 1 hour after completing the methacholine challenge FEV₁ had returned to

baseline values in all three treatment groups.

Conclusion—Formoterol has a greater intrinsic activity than salmeterol as a bronchoprotective agent, indicating that salmeterol is a partial agonist compared with formoterol in contracted human airways in vivo. Irrespective of this, prior long term treatment with both long acting β_2 agonists reduced the bronchodilating effect of an additional single dose of salbutamol equally, indicating that the development of tolerance or high receptor occupancy overshadowed any possible partial antagonistic activity of salmeterol. Patients on regular treatment with long acting β_2 agonists should be made aware that an additional single dose of a short acting β_2 agonist may become less effective.

(Thorax 2001;56:529–535)

Keywords: formoterol; salmeterol; salbutamol; asthma; tolerance

Inhaled β_2 agonists are extensively used in the treatment of asthma due to their excellent bronchodilating and bronchoprotective effects.^{1–3} As the duration of action of the two newest β_2 agonists, salmeterol and formoterol, is longer than of the older β_2 agonists, they are classified as long acting β_2 agonists. Both salmeterol and formoterol have an established position in the maintenance treatment of moderate to severe asthma, with short acting β_2 agonists being used additionally “as needed”.³

Salmeterol and formoterol differ pharmacologically in that formoterol has a much faster onset of action than salmeterol^{4–6} and, in vitro, salmeterol is a partial agonist on the β_2 receptor while formoterol is virtually a full agonist.^{7,8} During strong cholinergic bronchoconstriction, salmeterol has been shown to behave as an antagonist towards subsequently administered β_2 agonists.^{7,8} In vivo, such cholinergic bronchoconstriction may exist during an acute asthma attack.

Many studies have focused on tachyphylaxis and tolerance during treatment with β_2 agonists.^{9–18} This phenomenon appears to have a stronger effect after continued treatment of the β_2 agonist than after a single dose.

In this study we have investigated whether formoterol and salmeterol, given in high doses, behave as antagonists to subsequently given

Department of
Pulmonology, Martini
Hospital, 9700 RM
Groningen, The
Netherlands
H J van der Woude
T H Winter
R Aalbers

Correspondence to:
Dr R Aalbers
r.aalbers@mzh.nl

Received 26 May 2000
Returned to authors
10 December 2000
Revised version received
13 March 2001
Accepted for publication
4 April 2001

Table 1 Characteristics of patients

Sex (female/male)	15/4
Age (years)	38 (7), (24–49)
FEV ₁ (l)	2.88 (0.66), (1.74–3.98)
FEV ₁ (% predicted)	88.4 (16.1), (63–111)
PC ₂₀ methacholine (mg/ml)	0.62 (0.06–4.30)
PD ₃₀₊ methacholine (mg)	0.77 (0.18–3.18)
Inhaled steroid dose (µg/day)	500 (400–1600)

Values are presented as absolute numbers or as mean (SD), (range) except PC₂₀, PD₃₀₊, and steroid dose where a median (range) is given.

FEV₁ = forced expiratory volume in 1 second; PC₂₀ = provocative concentration of methacholine causing a decrease in FEV₁ of 20%; PD₃₀₊ = provocative dose of methacholine causing decrease in FEV₁ of ≥30%.

salbutamol. The effect of β_2 agonist induced reversal of methacholine induced bronchoconstriction (inducing a fall in FEV₁ of at least 30% from baseline) was used as a model.⁴ The assumption was that, after strong cholinergic stimulation, a difference in the ability of salbutamol to reverse the bronchoconstriction may develop depending on the type of additional β_2 agonist pretreatment, creating an in vivo analogy of the in vitro study by Molimard *et al.*⁸ In contrast to Molimard's model which used single dose administration, high dose maintenance treatment with the two long acting β_2 agonists was applied which has more relevance to the clinical situation.

Methods

PATIENTS

Patients diagnosed as having asthma according to the American Thoracic Society guidelines were invited to participate in the study.¹ The inclusion criteria included age 18–45 years; forced expiratory volume in one second (FEV₁) >1.5 l and >60% predicted¹⁹; concentration of methacholine causing a decrease in FEV₁ of 20% (PC₂₀) 4 mg/ml maximum; and decrease in FEV₁ of at least 30% during methacholine provocation. Exclusion criteria included concomitant diseases or conditions that might affect the study; use of long acting β_2 agonists, oral antihistamines, or oral bronchodilators for 24 hours before the enrolment visit until completion of the study; change in dose of inhaled corticosteroids or use of oral steroids in the 6 weeks before the study; and pregnancy. All patients had mild to moderate asthma with obvious bronchial hyperresponsiveness and were being treated with an inhaled glucocorticosteroid (continued unchanged during the study) and a bronchodilator on an “as needed” basis (replaced with ipratropium bromide during the study).

Table 2 Mean (SD) forced expiratory volume in one second (FEV₁) in litres on the three test days before and after the morning dose, at reaching a methacholine induced decrease in FEV₁ of ≥30%, at 3 and 30 minutes after salbutamol inhalation, and at 30 minutes after additional ipratropium bromide inhalation

	Pre-dose	Post-dose (baseline)	After methacholine (t = 0 min)	After salbutamol (t = 3 min)	After salbutamol (t = 30 min)	After additional ipratropium (t = 60 min)
Formoterol (n=17)	2.82 (0.72)	2.99 (0.74)	1.73 (0.52)	1.94 (0.54)*	2.50 (0.71)*	2.90 (0.75)
Salmeterol (n=16)	2.80 (0.69)	2.98 (0.73)	1.83 (0.56)	2.00 (0.54)*	2.57 (0.65)*	2.94 (0.77)
Placebo (n=16)	2.65 (0.67)	2.67 (0.65)	1.68 (0.46)	2.10 (0.56)	2.68 (0.67)	2.94 (0.75)

*p<0.05 compared with placebo (significance only tested at 3 and 30 minutes).

The study was approved by the medical ethics committee of the Martini Hospital, Groningen, the Netherlands and was conducted according to Good Clinical Practice Guidelines. Written informed consent was obtained from all patients prior to enrolment.

STUDY DESIGN

This double blind, randomised, crossover study used a double dummy technique for administering study drugs. The enrolment visit and each of the three test days were separated by three treatment periods of 2 weeks.

Inhalers with formoterol (blinded Oxis Turbuhaler, 12 µg formoterol fumarate per metered dose mixed with lactose, equivalent to 9 µg delivered dose), salmeterol (blinded Ser-event Diskhaler, 50 µg salmeterol xinafoate per dose mixed with lactose), and placebo (lactose) were provided by AstraZeneca R&D Lund. Two doses from each of two inhalers were administered twice daily over 2 weeks, starting in the evening after a visit day and using the last dose on the morning of the clinic visit, after measuring FEV₁. The daily dosages were thus 48 µg formoterol and 200 µg salmeterol. The 1:4 relationship has previously been reported to result in approximately equal bronchodilation; the high dose was applied to ensure maximal bronchodilation.⁶ Ipratropium bromide was allowed as rescue medication (40 µg Atrovent capsules administered via Inhalator Ingelheim, Boehringer Ingelheim, the Netherlands). No washout period was deemed necessary since it was expected that a new steady state situation would be attained within 1 week.⁹ Before each visit the patients had not taken inhaled ipratropium bromide, inhaled glucocorticosteroids, or cromoglycate in the previous 6 hours, nor caffeine containing beverages in the previous 2 hours. They also refrained from exercise before the tests.

A test day was postponed if morning FEV₁ was below 80% or above 120% of the baseline FEV₁ at enrolment. In the morning of the three test days pre-dose lung function was measured, followed by the last dose of study medication. One hour later lung function was again measured—referred to hereafter as the post-dose and “baseline” lung function—and the methacholine provocation test started. The provocation tests were scheduled within 1 hour of the time at which they were performed during the enrolment visit. The same technician performed all tests. At the initial visit the patients practised their inhalation technique with empty inhalers in accordance with the manufacturers' instructions. At the screening visit and 1 hour after the morning dose on the three test days a standard methacholine provocation test was performed and continued until a reduction of at least 30% in FEV₁ was achieved.⁴ The challenges were performed using a Wiesbadener Doppelinhalator giving an output of 0.2 ml in 2 minutes. After saline, doubling concentrations of methacholine bromide from 0.125 mg/ml to 64 mg/ml dissolved in saline were inhaled at 5 minute intervals during 2 minutes of tidal breathing. FEV₁ was measured with a dry spirometer (Schiller

Figure 1 Mean (SE) percentage predicted forced expiratory volume in one second (FEV_1) before and one hour after the morning dose, at reaching a methacholine induced decrease in FEV_1 of $\geq 30\%$, during 30 minutes following inhalation of $50 \mu\text{g}$ salbutamol, and after an additional 30 minutes following inhalation of $40 \mu\text{g}$ ipratropium bromide. Pretreatment for 2 weeks with formoterol $24 \mu\text{g}$ twice daily via Turbuhaler, salmeterol $100 \mu\text{g}$ twice daily via Diskhaler, or placebo twice daily.

SP-100, Schiller, Baar, Switzerland) which was calibrated daily. Pre-dose and post-dose (baseline) FEV_1 were measured in triplicate and the highest FEV_1 values were used in the analyses. The baseline FEV_1 (after the study drug but before saline) was taken as the reference value for the calculation of changes in FEV_1 . Single FEV_1 assessments were made 30 and 90 seconds after completing each 2 minute nebulisation and the lowest value was used for the analyses. The methacholine provocation test was stopped when FEV_1 decreased by 30% or more compared with baseline. One dose of salbutamol ($50 \mu\text{g}$ via Turbuhaler, Astra Draco, Lund, Sweden) was inhaled within 1 minute of reaching a decrease in FEV_1 of $>30\%$. Single assessments of FEV_1 were made at 1, 3, 5, 10, 15, 20, 25 and 30 minutes after inhaling salbutamol (0 min). A single dose of ipratropium bromide ($40 \mu\text{g}$) was then inhaled and lung function was assessed in triplicate after a further 30 minutes. If dyspnoea became too severe, either during or within 60 minutes of the provocation test, as judged by the patient or the technician, a bronchodilator was administered and the test halted and an inhaled bronchodilator was then administered freely. This test was repeated on another day.

Compliance was assessed by counting either the number of used and unused Diskhaler doses returned or the number of remaining clicks in the Turbuhaler. Adverse events were recorded at the start of each test day.

DATA ANALYSIS

The primary parameter in the statistical analysis was the increase in FEV_1 from 0 to 3 minutes after inhaling salbutamol, expressed as % of the baseline value (post-dose FEV_1). Secondary parameters (FEV_1 at 30 and at 60 minutes, PC_{20} methacholine, PD_{30+} methacholine, and the recovery time) were statistically analysed to a limited extent to prevent false positive conclusions. The PC_{20} was calculated by linear interpolation of the log methacholine concentration (non-cumulative) versus the percentage fall in FEV_1 from baseline. The

Figure 2 Methacholine dose needed to provoke a fall in FEV_1 of $\geq 30\%$ (PD_{30+}) measured about one hour after the last dose of treatment for 14 days with either formoterol $24 \mu\text{g}$ twice daily via Turbuhaler, salmeterol $100 \mu\text{g}$ twice daily via Diskhaler, or placebo. Log transformed data. (A) Individual data (horizontal lines represent mean values); (B) mean values with 95% confidence intervals.

cumulative methacholine dose administered to induce the PD_{30+} was calculated from the nebuliser output, arbitrarily assuming that all methacholine nebulised during 2 minutes penetrated into the lungs. The recovery time was calculated from the data points immediately before and after FEV_1 returned to 85% of baseline by linear interpolation. In cases where FEV_1 did not return to 85% of baseline within 60 minutes, an arbitrary time of 75 minutes was used in the analyses. For the statistical analysis of PC_{20} , PD_{30+} and the recovery time a log transformation was made. If the lowest methacholine dose caused a decrease in FEV_1 of 20% or more, PC_{20} was arbitrarily set at half the lowest concentration—that is, 0.0625 mg/ml .

Analysis of covariance (ANCOVA) was used to determine treatment differences with patient, period, and treatment as factors. The observed maximal percentage decreases in FEV_1 and PD_{30+} were used as covariates since both the magnitude of the fall in FEV_1 and the methacholine dose could be expected to have an influence on the recovery of FEV_1 at 3 minutes and the recovery time. When a covariate was not statistically significant it was excluded from the analyses. A two sided p value of <0.05 was considered significant. When treatment was statistically significant, 95% confidence intervals (95% CI) were calculated from the least squares means. Differences in PC_{20} and PD_{30+} between treatments are expressed as doubling dose steps, originating from analyses from base-2 log transformed data.

Table 3 Methacholine provocation tests and recovery time of FEV₁

	PC ₂₀ methacholine (mg/ml)	PD ₃₀₊ methacholine (mg)	Recovery time (min)
Formoterol (n=17)	2.23 (1.6 to 3.1)*	2.48 (0.38–25.6)*#	30.6 (21 to 45)*
Salmeterol (n=16)	1.59 (1.1 to 2.3)*	1.58 (0.28–6.4)*	25.8 (17 to 39)*
Placebo (n=16)	0.81 (0.57 to 1.1)	0.74 (0.08–4.8)	6.9 (4.5 to 10)

Values are geometric mean (95% CI) for PC₂₀ and recovery time, and geometric mean (range) for PD₃₀₊. Values for recovery time above 60 minutes arbitrarily set at 75 minutes.

PC₂₀ = concentration of methacholine provoking a decrease in FEV₁ of 20%; PD₃₀₊ = cumulative dose of methacholine provoking a decrease in FEV₁ of ≥30%.

*p<0.05 compared with placebo; #p<0.05 compared with salmeterol.

Figure 3 Increase in FEV₁ three minutes after inhalation of 50 µg salbutamol via Turbuhaler as percentage of the lowest FEV₁. Salbutamol was inhaled immediately after reaching a methacholine induced decrease in FEV₁ of ≥30% about one hour after the last dose of either formoterol 24 µg twice daily via Turbuhaler, salmeterol 100 µg twice daily via Diskhaler, or placebo via both devices for 2 weeks. (A) Individual values; (B) mean values and 95% confidence intervals.

It was estimated that 20 enrolled subjects were needed to give at least 15 subjects who could be fully evaluated. The effect at 3 minutes was considered to be the primary parameter in the analysis. A power calculation, using the results of a previous study,⁴ indicated that a difference of 4% could be detected in the increase in FEV₁ after 3 minutes with a significance level of 5% and a power of 80%.

Results

Nineteen patients were enrolled in the study and randomised. Four patients were withdrawn because of concurrent airway infections; all four were comparable to the other 15 patients in baseline lung function and PC₂₀ methacholine. The provocation tests were well tolerated and there was no need for an additional bronchodilator. Table 1 shows the patient

characteristics; the patients can be regarded as having mild to moderate asthma but with pronounced bronchial hyperresponsiveness.

The absolute FEV₁ values on test days are shown in table 2 and the percentage predicted FEV₁ during the entire test day is shown in fig 1. At the start of the test day, approximately 12 hours after inhaling the previous evening dose, pre-dose FEV₁ values in the formoterol and salmeterol periods were 0.15 l and 0.12 l higher, respectively, than in the placebo period. One hour after the morning dose the differences were 0.32 l and 0.26 l (approximately 10%, statistical significance not tested). Compared with placebo, more methacholine had to be administered following active treatment before FEV₁ decreased by 20% and by ≥30% from baseline. The PC₂₀ values after formoterol and salmeterol administration differed by 1.5 doubling dose steps (95% CI 0.8 to 2.1) and 1.0 doubling dose step (95% CI 0.3 to 1.7), respectively, both differences being statistically significant. The values between active treatments did not differ significantly (0.5 doubling dose steps, 95% CI -0.2 to 1.1). PD₃₀₊ differed in a similar way; geometric mean PD₃₀₊ was 2.48 mg after formoterol treatment, 1.58 mg after salmeterol pretreatment, and 0.74 mg after placebo pretreatment (fig 2). The difference in PD₃₀₊ of 0.7 doubling dose steps between the two active treatments was statistically significant (95% CI 0.1 to 1.2, p=0.016).

Apart from the difference in methacholine dose, the provocation tests on the three test days were comparable with a mean overall decrease in FEV₁ of 39.3% from baseline (ranging from 29.8% to 61.3% on individual days).

Three minutes after inhaling salbutamol FEV₁ increased by a mean (SD) of 15.8 (9.7)% of the baseline (pre-challenge) value after pretreatment with placebo compared with 7.3 (8.4)% after pretreatment with formoterol and 5.5 (8.9)% after pretreatment with salmeterol (p=0.0003 for both treatments, ANCOVA). The covariate actual % decrease in FEV₁ was statistically significant (p=0.014) whereas the covariate PD₃₀₊ was not (p=0.12). The difference in FEV₁ between pretreatment with placebo and with active drug was 11.9% (95% CI 6.1 to 17.7) for formoterol and 11.3% (95% CI 5.7 to 16.9) for salmeterol (both p=0.0003). The difference between the two active pretreatments was not significant (95% CI -6.2 to 5.0, p=0.82). In a post hoc analysis comparing only the two active treatments and taking PD₃₀₊ as covariate, the difference between the two long acting β₂ agonists remained not significant, nor was the covariate significant. The percentage increase in FEV₁ from the lowest FEV₁ after methacholine was 14 (19)%, 12 (21)%, and 26 (17)% after pretreatment with formoterol, salmeterol, and placebo, respectively (fig 3). Thirty minutes after inhaling salbutamol the increase in FEV₁ was still significantly smaller after both active pretreatments than after placebo (p=0.0001, ANCOVA), with again no significant difference between the two active treatments. Following all three pretreatments the mean FEV₁ returned to

the same value within 30 minutes after the inhalation of ipratropium bromide. The recovery time is shown in table 3. After pretreatment with both long acting β_2 agonists recovery was significantly slower than after placebo pretreatment ($p=0.0001$), but the difference between formoterol and salmeterol was not significant ($p=0.55$). Geometric mean recovery times were 35, 28, and 7 minutes after formoterol, salmeterol and placebo, respectively.

Compliance with drug intake was more than 80% for both inhalers in all three study periods. Adverse events were reported by seven of 18 patients treated with formoterol, five of 17 treated with salmeterol, and seven of 18 following treatment with placebo. Besides respiratory symptoms and airway infections, five patients reported headache, four tremor, and one palpitations.

Discussion

This study shows that the magnitude of effect of salbutamol is equally decreased and its onset equally delayed following pretreatment with both formoterol and salmeterol in patients using inhaled glucocorticosteroids. This observation is not due to the difference in partial agonist properties of the two long acting β_2 agonists since no difference between the two drugs was observed. However, despite using equipotent doses—which was also shown by similar and probably maximal bronchodilatation with the two drugs—formoterol had a stronger bronchoprotective effect than salmeterol when more than 20% bronchoconstriction was induced. This confirms *in vitro* findings that formoterol has a greater intrinsic activity and is a fuller agonist than salmeterol, especially in contracted bronchi.^{7,8} A similar stronger bronchoprotective effect of formoterol was also shown recently in a clinical study using high single doses.²⁰

To the best of our knowledge, this study is the first to compare the bronchodilator response of salbutamol during regular treatment with the two long acting β_2 agonists formoterol and salmeterol in the same patients using methacholine provocation to mimic a moderate to severe asthma attack.

We used high doses of formoterol and salmeterol and treatment lasted for 14 days to provide maximum stimulation of the β_2 receptor and to establish a situation in which tolerance was likely.⁹ The methacholine provocation test was performed 1 hour after administration of the last dose of formoterol and salmeterol, when the effects of formoterol and salmeterol were expected to be maximal and when a potential antagonistic property, reflected as a difference in response to the additional salbutamol, would be most apparent. Salbutamol was chosen to reverse methacholine induced bronchoconstriction rather than fenoterol because salbutamol is the most widely used rescue medication, despite the fact that fenoterol is a fuller agonist.^{7,8} The dose of salbutamol used may appear to be low but, when inhaled via a Turbuhaler, a 50 μ g dose of salbutamol has been reported to be equipotent to 200 μ g by Rotahaler or Diskhaler²¹ or 100 μ g

by metered dose inhaler²² and has been shown to be effective during acute bronchoconstriction.⁴ A single dose was used to increase the likelihood of detecting small differences between the two long acting β_2 agonists.

At the time of salbutamol inhalation the mean FEV₁ had decreased by 39%, corresponding to a moderate to severe asthma attack. After pretreatment with placebo there was a rapid onset of salbutamol action as shown in a previous study.⁴ However, a decreased response to inhaled salbutamol, expressed both in magnitude and as recovery time, was observed after both active treatments with no difference between the two treatments.

The observed lower sensitivity to inhaled salbutamol may have several causes. Firstly, there is evidence that continuous treatment with long acting β_2 agonists leads to a reduction in their bronchoprotective effect to a number of bronchoconstricting stimuli.^{23–25} This loss of bronchoprotective effect is due to β_2 receptor downregulation (tolerance) which is associated with subsensitivity to salbutamol, as shown in this study. This tolerance can only have been induced by the long acting β_2 agonists since no additional β_2 rescue medication was used during the study period. There is, however, controversy about the development of the diminished bronchodilator effect of salbutamol after long term treatment with long acting β_2 agonists, as seen in our study. Some studies have found that bronchodilator subsensitivity to cumulative doses of salbutamol may be induced after continuous treatment with long acting β_2 agonists, as shown by an increase in FEV₁ from baseline,^{11–13,15} with no induction of subsensitivity after long term treatment with formoterol.¹³ Others have determined the absolute values of lung function^{16–18} and found no reduction in the sensitivity to the bronchodilating effects of salbutamol. Probable explanations are differences in baselines^{16–18} and in the time of the last dose of long acting β_2 agonist (within 12 hours^{17,18} or after 12 hours^{11–13,15,16}). Also, unlike this study, concomitant β_2 agonist rescue medication was used during placebo treatment in some studies^{17,18} and some remaining β_2 receptor stimulation may have induced the reduced sensitivity. In none of these studies was the methacholine provocation test model used. The justification for distinguishing tolerance between the bronchodilator and bronchoprotective response has been questioned. The methacholine provocation test has recently been used to show that continuous treatment with short acting β_2 agonists induces bronchodilator tolerance,²⁶ something that other methods had previously failed to do.^{27,28} In this situation of increased bronchomotor tone, relaxation of airway smooth muscle may require more β_2 receptor activity and tolerance is easier to demonstrate.²⁶ It seems likely that continuous treatment with long acting β_2 agonists will also lead to tolerance in bronchodilating response, similar to the loss of bronchoprotective effect seen previously.

Secondly, in the present study the methacholine provocation test was performed at the peak of the agonist response. This may have obscured the effects of salbutamol because of

the high degree of receptor occupancy induced. It was previously been shown that inhaled formoterol (72 µg) and salmeterol (300 µg) antagonised extrapulmonary β_2 receptor mediated responses to inhaled fenoterol,²⁹ while a single, much lower dose (formoterol 12 µg, salmeterol 25 µg) had no such significant effects.³⁰ This indicates that, in the latter situation, a lower degree of receptor occupancy occurred and this did not interfere with fenoterol.

Thirdly, after active treatment a higher dose of methacholine was administered which may have led to a slower recovery in FEV₁. When tested in the ANCOVA, however, the PD₅₀₊ was not a significant covariate.

The relative contribution of the partial agonist property, β_2 receptor downregulation, and receptor occupancy to the observed diminished response to salbutamol could not be assessed in this study. As we did not study spontaneous recovery with placebo as rescue medication, the exact additional effect of salbutamol could not be determined. Furthermore, we did not investigate the effect of salbutamol after a single dose of the two long acting β_2 agonists simultaneously. These measurements may have provided additional information with regard to the partial antagonistic properties and the development of tolerance. It is possible that the effects of receptor occupancy and tolerance may have overshadowed the effect of partial antagonism in the present study. The almost identical recovery time following the two active treatments may have been a consequence of this and may be a reason why no clinically relevant contribution resulting from the partial β_2 antagonistic property of salmeterol (to the additionally inhaled salbutamol) during maintenance treatment could be demonstrated.

These results differ from in vitro findings where salmeterol appeared to antagonise the relaxation of cholinergically contracted human bronchi induced by other β_2 agonists such as salbutamol.⁷⁻⁸ The in vitro study also differed from our in vivo study in that the bronchi were contracted first, then incubated with salmeterol, formoterol or placebo and the dose response curve to salbutamol produced while, in our study, the bronchi were contracted after dosing with salmeterol, formoterol, and placebo. The present study more closely mimics the clinical situation where patients are using long acting β_2 agonists and, despite this, experience an asthma attack; this is clinically more relevant than β_2 agonist naive patients experiencing bronchoconstriction and then being treated with two simultaneous β_2 agonists for reversal.

Additional studies are needed to assess interactions between long acting and short acting β_2 agonists after a longer interval such as 12 hours from administration of the long acting β_2 agonists when the bronchodilation and bronchoprotection of the long acting drug is submaximal. It would be expected that, at this time, patients are more vulnerable to an attack of bronchospasm requiring additional salbutamol rescue therapy.

This study shows that, after pretreatment with long acting β_2 agonists, the bronchodilating effect of a single dose of salbutamol is diminished, both in magnitude and onset. With a higher dose of salbutamol this phenomenon may have been missed. Indeed, in a recent study a higher dose of salbutamol was found to overcome terbutaline induced tachyphylaxis in a methacholine provocative test.²⁶ However, a high dose of salbutamol did not overcome bronchoprotective subsensitivity in asthmatic patients receiving a single dose³¹ or regular treatment with salmeterol or formoterol,²⁵ although in these latter studies some interaction caused by receptor occupancy may have occurred.

The present study shows that formoterol has a greater intrinsic activity than salmeterol. Theoretically, stronger β_2 receptor stimulation may lead to a stronger tendency towards desensitisation. If this occurred, it would have resulted in a decreased effect on the FEV₁ 12 hours after the previous dose and/or a poorer immediate response in FEV₁ and PC₂₀ after inhalation of the study medication. It would also have led to a diminished effect of salbutamol. However, we found small and insignificant differences between the two long acting β_2 agonists on FEV₁ before methacholine provocation, a significantly stronger protective effect against methacholine after formoterol compared with salmeterol, and an equally diminished response to inhaled salbutamol. These results indicate that formoterol does not cause a stronger desensitisation than salmeterol or, if it does, any difference is counterbalanced by the antagonist activity of salmeterol.

In summary, the present study did not confirm in vivo the observed in vitro findings of differences in (partial) antagonistic activities between salmeterol and formoterol. However, a higher intrinsic activity of formoterol was demonstrated in contracted bronchi which indicates that salmeterol is a partial agonist compared with formoterol in human airways in vivo in moderate to severe asthma attacks. The efficacy of an additional single dose β_2 agonist is decreased in a situation of moderate to severe methacholine induced bronchoconstriction after pretreatment with formoterol and salmeterol. This suggests that patients on regular long acting β_2 agonist treatment should be aware that a single dose of salbutamol may become less effective during an acute asthma attack.

The authors thank M Boorsma for help in designing, monitoring and analysing the study, AstraZeneca R&D Lund for preparing blinded medication, Professor J Zaagsma for critically reviewing the manuscript and G D Nossent, M P Politiek, L H Steenhuis, W J Snoek, C G Tol, and S H Wills for contributions to the clinical part of the study.

This study was supported by a research grant from AstraZeneca, the Netherlands.

- 1 American Thoracic Society. Standards for the diagnosis and care of patients with COPD and asthma. *Am Rev Respir Dis* 1987;136:225-44.
- 2 Global Initiative for Asthma. *Global strategy for asthma management and prevention*. Publication no. 95-3659. Washington: National Heart, Lung, and Blood Institute, 1995.
- 3 British Thoracic Society. The British guidelines on asthma management. 1995 review and position statement. *Thorax* 1997;52(Suppl 1):S1-21.

- 4 Politiek MJ, Boersma M, Aalbers R. Comparison of formoterol, salbutamol and salmeterol in methacholine-induced severe bronchoconstriction. *Eur Respir J* 1999;13:988-92.
- 5 Van Noord JA, Smeets JJ, Raaijmakers JAM, et al. Salmeterol versus formoterol in patients with moderately severe asthma: onset and duration of action. *Eur Respir J* 1996;9:1684-8.
- 6 Palmqvist M, Persson G, Lazer L, et al. Inhaled dry-powder formoterol and salmeterol in asthmatic patients: onset of action, duration of effect and potency. *Eur Respir J* 1997;10:2484-9.
- 7 Källström BL, Sjöberg J, Waldeck B. The interaction between salmeterol and β_2 -adrenoceptor agonists with higher efficacy on guinea pig trachea and human bronchus in vivo. *Br J Pharmacol* 1994;113:687-92.
- 8 Molimard M, Naline E, Zhang Y, et al. Long- and short-acting β_2 adrenoceptor agonists: interactions in human contracted bronchi. *Eur Respir J* 1998;11:583-8.
- 9 Cockcroft DW, Swystun VA. Functional antagonism: tolerance produced by inhaled β_2 agonists. *Thorax* 1996;51:1051-6.
- 10 Newnham DM, Grove A, McDevitt DG, et al. Subsensitivity of bronchodilator and systemic β_2 adrenoceptor responses after regular twice daily treatment with formoterol dry powder in asthmatic patients. *Thorax* 1995;50:497-504.
- 11 Fuglsang G, Vikre-Jørgensen J, Angertoft L, et al. Effect of salmeterol treatment on nitric oxide level in exhaled air and dose-response to terbutaline in children with mild asthma. *Pediatr Pulmonol* 1998;25:314-21.
- 12 Grove A, Lipworth B. Bronchodilator subsensitivity to salbutamol after twice daily salmeterol in asthmatic patients. *Lancet* 1995;346:201-6.
- 13 Aziz I, Hall IP, McFarlane LC, et al. β_2 adrenoreceptor regulation and bronchodilator sensitivity after regular treatment with formoterol in subjects with stable asthma. *J Allergy Clin Immunol* 1998;101:337-41.
- 14 Tan KS, Grove A, McLean A, et al. Systemic corticosteroid rapidly reverses bronchodilator subsensitivity induced by formoterol in asthmatic patients. *Am J Respir Crit Care Med* 1997;156:28-35.
- 15 Lipworth BJ, Aziz I. Bronchodilator response to albuterol after regular formoterol and effects of acute corticosteroid administration. *Chest* 2000;117:156-62.
- 16 Langley SJ, Masterson CM, Batty EP, et al. Bronchodilator response to salbutamol after chronic dosing with salmeterol or placebo. *Eur Respir J* 1998;11:1081-5.
- 17 Wilding P, Clark M, Thompson Coon J, et al. Effect of long term treatment with salmeterol on asthma control: a double blind, randomised crossover study. *BMJ* 1997;314:1441-7.
- 18 Nelson HS, Berkowitz RB, Tinkelman DA, et al. Lack of subsensitivity to albuterol after treatment with salmeterol in patients with asthma. *Am J Respir Crit Care Med* 1999;159:1556-61.
- 19 Quanjer PH, Tammeling GJ, Cotes JE, et al. Lung function and forced ventilatory flows. Report working party standardisation of lung function tests. European Community for Steel and Coal. *Eur Respir J* 1993;6(Suppl 16):5-40.
- 20 Palmqvist M, Ibsen T, Mellén A, et al. Comparison of the relative efficacy of formoterol and salmeterol in asthmatic patients. *Am J Respir Crit Care Med* 1999;160:244-9.
- 21 Mahadew Singh JV, Hamersma WBGJ, Scheurs AJM. Relative efficacy of three different inhalers containing salbutamol in patients with asthma. *Eur J Clin Pharmacol* 1996;50:467-9.
- 22 Löfdahl CG, Andersson L, Bondesson E, et al. Differences in bronchodilating potency of salbutamol in Turbuhaler as compared with a pressurized metered-dose inhaler formulation in patients with reversible airway obstruction. *Eur Respir J* 1997;10:2474-8.
- 23 Yates DH, Sussman HS, Shaw MJ, et al. Regular formoterol treatment in mild asthma: effect on bronchial responsiveness during and after treatment. *Am J Respir Crit Care Med* 1995;152:1170-4.
- 24 Cheung D, Timmers MG, Zwinderman AH, et al. Long-term effects of long-acting β_2 adrenoreceptor agonist, salmeterol, on airway hyperresponsiveness in patients with mild asthma. *N Engl J Med* 1992;327:1198-203.
- 25 Lipworth BJ, Aziz I. A high dose of albuterol does not overcome bronchoprotective subsensitivity in asthmatic subjects receiving regular salmeterol or formoterol. *J Allergy Clin Immunol* 1999;103:88-92.
- 26 Hancox RJ, Aldridge RE, Cowan JO, et al. Tolerance to beta-agonists during acute bronchoconstriction. *Eur Respir J* 1999;14:283-7.
- 27 Tattersfield AE. Tolerance to beta agonists. *Bull Eur Physiopathol Respir* 1985;21:1-5s.
- 28 Grove A, Lipworth BJ. Tolerance with β_2 adrenoreceptor agonists: time for reappraisal. *Br J Clin Pharmacol* 1995;39:109-18.
- 29 Grove A, Lipworth BJ. Evaluation of the β_2 adrenoreceptor agonist/antagonist activity of formoterol and salmeterol. *Thorax* 1996;51:54-8.
- 30 Grove A, Lipworth BJ. Effects of prior treatment with salmeterol and formoterol on airway and systemic β_2 responses to fenoterol. *Thorax* 1996;51:585-9.
- 31 Aziz I, Lipworth BJ. In vivo effect of albuterol on methacholine-contracted bronchi in conjunction with salmeterol and formoterol. *J Allergy Clin Immunol* 1999;103:816-22.