- 5 Byington CL, Samore MH, Stoddard GJ, *et al*. Temporal trends of invasive disease due to *Streptococcus pneumoniae* among children in the intermountain west: emergence of nonvaccine serogroups. *Clin Infect Dis* 2005;**41**:21–9.

Repeated tuberculin testing does not induce false positive ELISPOT results

The Enzyme Linked ImmunoSpot (ELISPOT) is a new rapid T cell based blood test (otherwise known as an interferon- γ assay) for the diagnosis of latent tuberculosis infection.^{1–3} The commercially available form of the assay, T-SPOT® TB (Oxford Immunotec, Abingdon, UK) has European regulatory approval as an in vitro diagnostic test and is increasingly being used in clinical practice. The test is based on the enumeration of interferon- γ producing T cells which are specific for two highly antigenic proteins, early secretory antigenic target-6 (ESAT-6) and culture filtrate protein 10 (CFP-10).¹ These proteins are expressed by *Mycobacterium tuberculosis* but are absent from *M bovis* BCG vaccine. Hence, the test does not give false positive results in BCG vaccinated individuals.^{1–3}

ESAT-6 and CFP-10 are, however, contained within tuberculin purified protein derivative (PPD). Since ELISPOT is a highly sensitive method for measuring even low numbers of antigen specific T cells,⁴ concerns have been raised as to whether repeated tuberculin skin tests might induce T cell responses to these specific antigens, resulting in false positive ELISPOT results.

As T-SPOT® TB enters clinical practice, it may initially be used by some people in conjunction with the tuberculin skin test. It is therefore important to know whether false positive ELISPOT results are induced by tuberculin testing. The following results strongly suggest that this is not the case.

The results reported here are from a 2 year follow up of a group of people with potential point source exposure to multidrug resistant tuberculosis on a maternity unit in Modena University Hospital, Italy.⁵ Forty four BCG unvaccinated subjects were negative at initial screening by tuberculin skin test and ELISPOT, 3 months after the point source exposure ceased. All participants had negative results on serological testing for HIV infection. Tuberculin skin tests were administered and read by two experienced chest physicians using 5 units of PPD-S injected intradermally about 2 hours after blood was drawn for ELISPOT assays. The ELISPOT assays were performed and scored, as previously described,⁵ by two technicians without knowledge of personal identifiers. All these individuals underwent repeated testing by skin test and ELISPOT at 9, 15 and 24 months after the point exposure. At 24 months all 44 individuals remained ELISPOT negative, although three had become positive with the tuberculin skin test (fig 1). Thus, inoculation of three PPD skin tests over a 21 month period in 44 initially ELISPOT negative individuals did not induce any false positive ELISPOT results.

These results show that repeated tuberculin skin testing over time does not induce a T cell response to ESAT-6 or CFP-10 resulting in false positive ELISPOT results. Our findings suggest that this new interferon- γ blood assay could be used in association with the standard PPD skin test without any reduction in its high diagnostic specificity. Given the

Figure 1 Time course of development of positive Mantoux results in the three participants who became tuberculin skin test (TST) positive as a result of repeated skin testing.

high sensitivity of the ELISPOT assay for detecting even low numbers of antigen specific T cells, the absence of a detectable response to ESAT-6 and CFP-10 suggests that T cells specific for these antigens were not induced by repeated inoculation of PPD. This is consistent with the observation that ESAT-6 has very poor immunogenicity when administered as a candidate vaccine, unless inoculated with powerful adjuvants.⁶ This is in stark contrast to its potent immunogenicity when presented to the immune system during natural *M tuberculosis* infection; indeed, ESAT-6 is the strongest known target of T cell responses during tuberculosis infection.

Our results also suggest that T-SPOT® TB could be especially useful in distinguishing true latent tuberculosis infection from false positive tuberculin skin test results that have arisen through “boosting”. Boosting occurs in people who undergo repeated tuberculin skin tests (such as healthcare workers) and causes false positive skin test results in uninfected people. This phenomenon is a major problem in tuberculosis screening programmes for healthcare workers, prisoners, and other groups at persistent risk of tuberculosis exposure, and was almost certainly the reason why three individuals in our study developed positive skin test results after repeated testing. Our findings suggest that T-SPOT® TB will maintain its high specificity even in individuals with false positive skin test results due to boosting from repeated tuberculin testing. Thus, use of T-SPOT® TB could enhance our ability to screen for latent tuberculosis infection even in populations who have already been repeatedly screened by the skin test.

L Richeldi

Respiratory Disease Clinic, University of Modena and Reggio Emilia, and Azienda Ospedaliera Policlinico di Modena, Modena, Italy

K Ewer

Nuffield Department of Clinical Medicine, University of Oxford, John Radcliffe Hospital, Oxford, UK

M Losi, P Roversi, I M Fabbri

Respiratory Disease Clinic, University of Modena and Reggio Emilia, and Azienda Ospedaliera Policlinico di Modena, Modena, Italy

A Lalvani

Nuffield Department of Clinical Medicine, University of Oxford, John Radcliffe Hospital, Oxford, UK

Correspondence to: Dr A Lalvani, Nuffield Department of Clinical Medicine, University of Oxford, John Radcliffe Hospital, Oxford OX3 9DU, UK; a.jit.lalvani@ndm.ox.ac.uk

The study was approved by the Modena research ethics committee and each study participant provided written informed consent.

doi: 10.1136/thx.2005.049759

This work was supported by the Wellcome Trust and Azienda Ospedaliera Policlinico di Modena.

Competing interests: AL is a named inventor on patents relating to T cell based diagnosis filed by the University of Oxford. Regulatory approval and commercialisation of ELISPOT (T-SPOT TB) has been undertaken by a spin out company of the University of Oxford (Oxford Immunotec Ltd), in which AL has a share of equity and to which he acts as scientific advisor in a non-executive capacity. KE is a named inventor on a patent application relating to the application of ELISPOT filed by the University of Oxford. The University of Oxford has a share of equity in Oxford Immunotec Ltd.

References

- 1 Lalvani A, Pathan AA, McShane H, *et al*. Rapid detection of *Mycobacterium tuberculosis* infection by enumeration of antigen-specific T cells. *Am J Respir Crit Care Med* 2001;**163**:824–8.
- 2 Lalvani A, Pathan AA, Durkan H, *et al*. Enhanced contact tracing and spatial tracking of *Mycobacterium tuberculosis* infection by enumeration of antigen-specific T cells. *Lancet* 2001;**357**:2017–21.
- 3 Ewer K, Deeks J, Alvarez L, *et al*. Comparison of T-cell-based assay with tuberculin skin test for diagnosis of *Mycobacterium tuberculosis* infection in a school tuberculosis outbreak. *Lancet* 2003;**361**:1168–73.
- 4 Lalvani A, Brookes R, Hambleton S, *et al*. Rapid effector function in CD8+ memory T cells. *J Exp Med* 1997;**186**:859–65.
- 5 Richeldi L, Ewer K, Losi M, *et al*. T cell-based tracking of multidrug resistant tuberculosis infection after brief exposure. *Am J Respir Crit Care Med* 2004;**170**:288–95.
- 6 Brandt L, Elhay M, Rosenkrands I, *et al*. ESAT-6 subunit vaccination against *Mycobacterium tuberculosis*. *Infect Immun* 2000;**68**:791–5.

Clinical importance of the Step 3 choice in asthma

We read with interest the meta-analysis by Masoli *et al*¹ which aimed to further guide clinicians in their choice between addition of long acting β_2 agonists (LABA) or use of higher doses of inhaled corticosteroids (ICS) in patients with symptomatic asthma. The pooled odds of at least one moderate or severe exacerbation was 1.35 times higher in those receiving a higher dose of ICS than in those treated with LABA.

Unfortunately, it is difficult to draw any meaningful conclusion as to the clinical relevance of these findings or to compare at a glance the results with those of the previous MIASMA study² because of differences in the summary statistics presented. For clinicians to understand the clinical context of these two studies, it is helpful to calculate the number needed to treat (NNT), as was done in the original MIASMA study.

Of the 2312 patients randomised to LABA treatment included in the newer study, 184 experienced one or more moderate or severe exacerbations (an incidence of 79.6 per 1000 patients) compared with 243 of the 2264 patients randomised to high dose ICS treatment (an incidence of 107.3 per 1000 patients). These incidences give an attributable risk reduction of 27.7 per 1000 patients which represents an NNT of 37, meaning that for every 37 patients receiving LABA in preference to high dose ICS, one less will experience an exacerbation. The corresponding

NNT from the MIASMA study was 41, so the findings were consistent.

Pointedly, neither of these studies looked at any inflammatory outcomes. Although adding a LABA may reduce exacerbations in a complementary manner to ICS, this is likely to be due to stabilising airway smooth muscle rather than potentiating the anti-inflammatory activity of the ICS. For example, in a study of inflammatory markers,³ doubling the dose of fluticasone from 250 µg/day to 500 µg/day reduced exhaled nitric oxide and adenosine monophosphate hyperresponsiveness more effectively than adding salmeterol to the 250 µg dose. In other words, while adding salmeterol in preference to a higher dose of ICS might reduce exacerbations and exhibit putative steroid sparing activity, this will occur at the expense of worsening anti-inflammatory control. Without monitoring inflammation in patients who are asymptomatic on ICS/LABA combination inhalers, clinicians may be lulled into a false sense of security and overlook potential long term damage from untreated airway inflammation.

M L Barnes, B J Lipworth

Asthma & Allergy Research Group, Department of Medicine and Therapeutics, Ninewells Hospital & Medical School, University of Dundee, Dundee DD1 9SY, UK

Correspondence to: Mr M L Barnes, Asthma & Allergy Research Group, Department of Medicine and Therapeutics, Ninewells Hospital & Medical School, University of Dundee, Dundee DD1 9SY, UK; mbarnes@rcsed.ac.uk

Sponsors: none.

Competing interests: The Asthma & Allergy Research Group have been financially supported by AstraZeneca, GlaxoSmithKline, Neolab, IVAX, Altana, Schering-Plough, and Merck for postgraduate lectures and meeting attendance, educational support and clinical trials.

References

- 1 Masoli M, Weatherall M, Holt S, *et al*. Moderate dose inhaled corticosteroids plus salmeterol versus higher doses of inhaled

corticosteroids in symptomatic asthma. *Thorax* 2005;**60**:730–4.

- 2 Shrewsbury S, Pyke S, Britton M. Meta-analysis of increased dose of inhaled steroid or addition of salmeterol in symptomatic asthma (MIASMA). *BMJ* 2000;**320**: 1368–73.
- 3 Currie GP, Bates CE, Lee DK, *et al*. Effects of fluticasone plus salmeterol versus twice the dose of fluticasone in asthmatic patients. *Eur J Clin Pharmacol* 2003;**59**:11–5.

Authors' reply

We appreciate the opportunity to respond to the issues raised by Barnes and Lipworth. However, with regard to calculating the number needed to treat (NTT), it is not clear that clinicians necessarily find this a useful measurement.¹ Most meta-analysis techniques use a weighted pooled outcome measurement that takes into account the different sample sizes and/or variances of each individual study measurement. The crude simple sum of events in both treatment groups that Barnes and Lipworth have suggested using does not. When the weighted technique is applied to the whole data set, under a fixed effects model this gives a pooled NTT of 58.4 (95% CI 32.6 to 278.3)—nearly double the number calculated by the crude method.

NNT refers to a specific time and this calculation does not take account of the fact that nearly half the studies ran for 12 weeks and the other half for 24 weeks (one for 26 weeks). The NTT for the 12 week studies was 75.5 (95% CI for the probability difference crosses zero) and for the 24 week studies it was 35.4 (95% CI 18.2 to 619.9). The point estimates for the two groups of studies are concordant in that 2×35.4 is close to 75. All but one of the studies analysed for exacerbations in the original MIASMA paper² ran for 24 weeks (the other study ran for 26 weeks) so that, if only the 24 week studies are used, our paper and the MIASMA paper agree.

Barnes and Lipworth also raise the issue of whether surrogate markers of airways inflammation such as exhaled nitric oxide

and adenosine monophosphate responsiveness are preferable to clinical measures such as severe exacerbations, lung function, night wakenings, and rescue β agonist use. The advantage of these clinical measures is that they represent relevant validated methods to assess long term asthma control and the risk of morbidity and mortality; this is not the case with the surrogate inflammatory markers. For this reason we consider that the findings from our meta-analysis should provide clinicians with greater confidence when deciding the dose of inhaled corticosteroid at which to consider adding salmeterol at Step 3 in the asthma guidelines.

M Weatherall, M Masoli, R Beasley

Medical Research Institute of New Zealand, P O Box 10055, Wellington, New Zealand

Correspondence to: Professor R Beasley, Medical Research Institute of New Zealand, P O Box 10055, Wellington, New Zealand; richard.beasley@mrnz.ac.nz

References

- 1 Smeeth L, Haines A, Ebrahim S. Numbers needed to treat derived from meta-analyses: sometimes informative, usually misleading. *BMJ* 1999;**318**:1548–51.
- 2 Shrewsbury S, Pyke S, Britton M. Meta-analysis of increased dose of inhaled steroid or addition of salmeterol in symptomatic asthma (MIASMA). *BMJ* 2000;**320**:1368–73.

ERRATUM

The name of the last author was missed from abstract number S40, *Thorax* 2005; **60**(suppl II):ii16. The correct listing of authors is: A Laverty¹, P Weller², A Jaffe¹ 1. Portex Respiratory Unit, Great Ormond Street Hospital for Children, London; 2. Centre for Measurement and Information in Medicine, City University, London.

The journal apologises for this error.