- pulmonary disease: more effective and less expensive. *Crit Care Med* 2000;**28**:2094–102.
- 128 **Plant PK**, Owen JL, Parrott S, *et al*. Cost effectiveness of ward based non-invasive ventilation for acute exacerbations of chronic obstructive pulmonary disease: economic analysis of randomised controlled trial. *BMJ* 2003;**326**:956.
 - 129 **Esteban A**, Anzueto A, Frutos F, *et al*. Characteristics and outcomes in adult patients receiving mechanical ventilation: a 28-day international study. *JAMA* 2002;**287**:345–55.
 - 130 **Nava S**, Ambrosino N, Cini E, Prato M, *et al*. Noninvasive mechanical ventilation in the weaning of patients with respiratory failure due to chronic obstructive pulmonary disease. A randomized, controlled trial. *Ann Intern Med* 1998;**128**:721–8.
 - 131 **Ferrer M**, Esquinas A, Arancibia F, *et al*. Noninvasive ventilation during persistent weaning failure: a randomized controlled trial. *Am J Respir Crit Care Med* 2003;**168**:70–6.
 - 132 **Esteban A**, Frutos-Vivar F, Ferguson ND, *et al*. Noninvasive positive-pressure ventilation for respiratory failure after extubation. *N Engl J Med* 2004;**350**:2452–60.
 - 133 **Donner CF**, Muir JF. Selection criteria and programmes for pulmonary rehabilitation in COPD patients. Rehabilitation and Chronic Care Scientific Group of the European Respiratory Society. *Eur Respir J* 1997;**10**:744–57.
 - 134 **Shepperd S**, Harwood D, Gray A, *et al*. Randomised controlled trial comparing hospital at home care with inpatient hospital care. II: Cost minimisation analysis. *BMJ* 1998;**316**:1791–6.
 - 135 **Cotton MM**, Bucknall CE, Dagg KD, *et al*. Early discharge for patients with exacerbations of chronic obstructive pulmonary disease: a randomized controlled trial. *Thorax* 2000;**55**:902–6.
 - 136 **Nicholson C**, Bowler S, Jackson C, *et al*. Cost comparison of hospital- and home-based treatment models for acute chronic obstructive pulmonary disease. *Aust Health Rev* 2001;**24**:181–7.
 - 137 **Davies L**, Wilkinson M, Bonner S, *et al*. "Hospital at home" versus hospital care in patients with exacerbations of chronic obstructive pulmonary disease: prospective randomised controlled trial. *BMJ* 2000;**321**:1265–8.
 - 138 **Ojoo JC**, Moon T, McGlone S, *et al*. Patients' and carers' preferences in two models of care for acute exacerbations of COPD: results of a randomised controlled trial. *Thorax* 2002;**57**:167–9.
 - 139 **Hernandez C**, Casas A, Escarabill J, *et al*. Home hospitalisation of exacerbated chronic obstructive pulmonary disease patients. *Eur Respir J* 2003;**21**:58–67.
 - 140 **Skwarska E**, Cohen T, Skwarski KM, *et al*. Randomized controlled trial of supported discharge in patients with exacerbations of chronic obstructive pulmonary disease. *Thorax* 2002;**55**:907–12.

LUNG ALERT

Asthma and natural killer T cells

▲ Akbari O, Faul JL, Hoyte EG, *et al*. CD4+ invariant T-cell-receptor+ natural killer T cells in bronchial asthma. *N Engl J Med* 2006;**354**:1117–29

The presence of CD4+ T helper cells in the airways of patients with asthma is well recognised, precipitating airway inflammation via cytokines such as interleukin-4, interleukin-5, and interleukin-13. Newly identified human natural killer T cells also express CD4. A subclass of this group, invariant natural killer T cells, possesses a restricted range of T cell receptors (V α 24-J α 18) which bind to glycolipid antigens presented by the class I MHC-like protein CD1d. Stimulation of such cells generates a potent cytokine response. The exact role of invariant natural killer T cells in humans is unknown. This study examined their role in asthma.

Twenty five patients were recruited, 14 with moderately severe asthma (10 taking inhaled corticosteroids), five with sarcoidosis, and six controls. None had received oral corticosteroids within 3 months of the study. Bronchoalveolar lavage (BAL), peripheral blood lymphocytes, and endobronchial biopsies were analysed.

Increased BAL lymphocyte counts with predominant CD4+ expression were seen in both patients with asthma and those with sarcoidosis. However, there were significantly more invariant natural killer T cells in those with asthma (mean 63%) than sarcoidosis (mean <1%). In asthma patients, natural killer T cells were over 100 times more abundant in BAL fluid than in peripheral blood. There was no significant difference in asthma between patients who were and were not receiving inhaled corticosteroids.

This study supports a role for invariant natural killer T cells in the pathogenesis of asthma and as a target for future therapy.

H E Davies

Specialist Registrar, Royal Berkshire Hospital, Reading, UK; hedavies@doctors.net.uk