IMAGES IN THORAX

MRI features of pleural endometriosis after catamenial haemothorax

Pleural endometriosis typically presents in young women within 72 h of onset of menstruation as catamenial pneumothorax, haemothorax or both. A 43-year-old multiparous woman with a history of pelvic endometriosis and myocardial infarction presented with acute onset of right hemithorax pain and dyspnoea coinciding with dysmenorrhoeic menstruation. A non-contrast enhanced chest CT scan (fig 1) showed a right haemopneumothorax with floating clots within the fluid, which was drained. A video-assisted thoracoscopy (VATS) guided random pleural biopsy sample was negative. Magnetic resonance imaging (MRI) of the chest coinciding with menses 4 months later showed an

oval-shaped lesion with homogeneous high signal intensity in T1- and T2-weighted images in the right posterior costodiaphragmatic recess (fig 2A, B). Owing to the relative contraindication for hormone therapy related to the coronary artery disease, the patient underwent a hysterectomy and oophorectomy for treatment of endometriosis. Six months later she was free of thoracic symptoms and an MRI scan showed the disappearance of the right pleural abnormality.

The diagnosis of thoracic endometriosis is usually based on clinical grounds but is often delayed. Cytological examination of the pleural fluid is usually not helpful and VATS is often negative.^{1,2} Although no

Learning points

- VATS is often non-diagnostic in patients with catamenial haemothorax or haemopneumothorax.
- The MRI appearance of a pleural-based lesion exhibiting homogeneous high signal in T1- and T2-weighted images is distinctive of pleural endometriosis.
- MRI after drainage of the pleural effusion and coinciding with menses could support the diagnosis of pleural endometriosis in women with catamenial haemothorax or haemopneumothorax.

Figure 1 Non-contrast enhanced chest CT scan at presentation showing (A) right haemopneumothorax in the lung parenchyma and (B) evidence of slightly hyperdense floating clots (arrows) within the pleural fluid.

Figure 2 Non-contrast enhanced MRI scan of the chest and upper abdomen showing an oval-shaped well-defined lesion in the right posterior costodiaphragmatic recess abutting the pleura. The lesion (arrow) exhibits homogeneous high signal intensity in (A) the coronal T1-weighted spin echo image and (B) the axial T2-weighted fast spin echo images without fat suppression.

pathological proof was available, the diagnosis of pleural endometriosis in our case was supported by the typical clinical presentation, the history of pelvic endometriosis, the MRI features distinctive of pleural endometriosis³ and the disappearance of the pleural MRI finding after surgical treatment of endometriosis.

Giulia Picozzi, Davide Beccani

Radiodiagnostic Section, Department of Clinical Physiopathology University of Florence, Italy

Florio Innocenti, Michela Grazzini

Respiratory Unit, Pistoia Hospital, Pistoia, Italy

Mario Mascacchi

Radiodiagnostic Section, Department of Clinical Physiopathology University of Florence, Italy

doi: 10.1136/thx.2006.071415

Correspondence to: Professor Mario Mascacchi, Radiodiagnostic Section, Department of Clinical Physiopathology, University of Florence, Viale Morgagni 50134, Florence, Italy; m.mascacchi@dfc.unifi.it

References

- 1 Joseph J, Sahani SA. Thoracic endometriosis syndrome: new observations from an analysis of 110 cases. *Am J Med* 1996;**100**:164–70.
- 2 Johnson MM. Catamenial pneumothorax and other thoracic manifestations of endometriosis. *Clin Chest Med* 2004;**25**:311–9.
- 3 Cassina PC, Hauser M, Kael G, et al. Catamenial hemoptysis: diagnosis with MRI. *Chest* 1997;**111**:1447–50.