15. **Jiang J**, Shi HZ, Liang QL, *et al*. Diagnostic value of interferon- γ in tuberculous pleurisy: a metaanalysis. *Chest* 2007;**131**:1133–41.
16. **Irwig L**, Macaskill P, Glasziou P, *et al*. Meta-analytic methods for diagnostic test accuracy. *J Clin Epidemiol* 1995;**48**:119–30.
17. **Vamvakas EC**. Meta-analyses of studies of the diagnostic accuracy of laboratory tests: a review of the concepts and methods. *Arch Pathol Lab Med* 1998;**122**:675–86.
18. **Suzuki S**, Moro-oka T, Choudhry NK. The conditional relative odds ratio provided less biased results for comparing diagnostic test accuracy in meta-analyses. *J Clin Epidemiol* 2004;**57**:461–9.
19. **Westwood ME**, Whiting PF, Kleijnen J. How does study quality affect the results of a diagnostic meta-analysis? *BMC Med Res Methodol* 2005;**5**:20.
20. **Egger M**, Davey Smith G, Schneider M, *et al*. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**:629–34.
21. **Porcel JM**, Light RW. Diagnostic approach to pleural effusion in adults. *Am Fam Physician* 2006;**73**:1211–20.
22. **Glas AS**, Lijmer JG, Prins MH, *et al*. The diagnostic odds ratio: a single indicator of test performance. *J Clin Epidemiol* 2003;**56**:1129–35.
23. **Deeks JJ**. Systematic reviews of evaluations of diagnostic and screening tests. In: Egger M, Smith GD, Altman DG, eds. *Systematic reviews in health care. Meta-analysis in context*. London: BMJ Publishing Group, 2001:248–82.
24. **Jaeschke R**, Guyatt G, Lijmer J. Diagnostic tests. In: Guyatt G, Rennie D, eds. *Users' guides to the medical literature. A manual for evidence-based clinical practice*. Chicago: AMA Press, 2002:121–40.
25. **Booth SN**, Lakin G, Dykes PW, *et al*. Cancer-associated proteins in effusion fluids. *J Clin Pathol* 1977;**30**:537–40.
26. **Pettersson T**, Klockars M, Froseth B. Neuron-specific enolase in the diagnosis of small-cell lung cancer with pleural effusion: a negative report. *Eur Respir J* 1988;**1**:698–700.
27. **Pettiti DB**. Approaches to heterogeneity in meta-analysis. *Stat Med* 2001;**20**:3625–33.

Pulmonary puzzle

A large chest wall tumour in an asymptomatic 15-year-old girl

CLINICAL PRESENTATION

A 15-year-old girl that had plain chest x ray as part of a routine medical examination was found to have a large left lower chest wall mass associated with rib destruction that was later shown to be painless and non-palpable (fig 1A). Chest CT revealed a 7×11 cm pleura based mass located in the left lower thorax and featured multiple rib destruction and downward depression of the left hemidiaphragm (fig 1B).

CT guided biopsy revealed a lesion with clusters of melanin-laden tumour cells. These tumour cells were positive for HMB-45 and S-100 protein immunohistochemically. The cytoplasmic pigments could be bleached by superoxidative agent and were positive by Fontana Masson stain, consistent with melanin pigments in nature. The skin overlying the tumour was intact, and the tumour was impalpable. Thorough skin examination over her whole body by an experienced dermatologist did not reveal any suspicious skin lesion. No other abnormality was demonstrated on whole body image studies.

QUESTION

What is the likely diagnosis and how should this be confirmed?

See page 94 for answer

This case is submitted by:

K-C Chen,¹ W-C Lin,² Y-C Lee,^{1,3} J-S Chen,¹ H-H Hsu¹

¹ Department of Surgery, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan; ² Department of Pathology, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan; ³ Department of Traumatology, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan

Correspondence to: Dr H-H Hsu, Department of Surgery, National Taiwan University Hospital, No 7, Chung-Shan South Road, Taipei, Taiwan; ntuhsf@yahoo.com.tw

Competing interests: None.

Thorax 2008;**63**:41. doi:10.1136/thx.2007.087981

Figure 1 (A) Chest plain film revealing a large, well defined left chest wall tumour with obvious rib destruction. (B) Thoracic CT revealing a large expansile tumour associated with osteolytic and sclerotic changes to a number of the patient's ribs.