6. **Stanescu D**, Sanna A, Veriter C, *et al*. Identification of smokers susceptible to development of chronic airflow limitation: a 13-year follow-up. *Chest* 1998;**114**:416–25.
7. **Vestbo J**, Hogg JC. Convergence of the epidemiology and pathology of COPD. *Thorax* 2006;**61**:86–8.
8. **Gevenois PA**, Scillia P, de Maertelaer V, *et al*. The effects of age, sex, lung size, and hyperinflation on CT lung densitometry. *AJR Am J Roentgenol* 1996;**167**:1169–73.
9. **Mishima M**, Hirai T, Itoh H, *et al*. Complexity of terminal airspace geometry assessed by lung computed tomography in normal subjects and patients with chronic obstructive pulmonary disease. *Proc Natl Acad Sci U S A* 1999;**96**:8829–34.
10. **McWilliams A**, Mayo J, MacDonald S, *et al*. Lung cancer screening: a different paradigm. *Am J Respir Crit Care Med* 2003;**168**:1167–73.
11. **American Thoracic Society**. Standardization of Spirometry, 1994 update. *Am J Respir Crit Care Med* 1995;**152**:1107–36.
12. **Crapo RO**, Morris AH, Gardner RM. Reference spirometric values using techniques and equipment that meet ATS recommendations. *Am Rev Respir Dis* 1981;**123**:659–64.
13. **Yuan R**, Mayo JR, Hogg JC, *et al*. The effects of radiation dose and CT manufacturer on measurements of lung densitometry. *Chest* 2007;**132**:617–23.
14. **Hedlund LW**, Vock P, Effmann EL. Computed tomography of the lung. Densitometric studies. *Radiol Clin North Am* 1983;**21**:775–88.
15. **Coxson HO**, Mayo JR, Behzad H, *et al*. Measurement of lung expansion with computed tomography and comparison with quantitative histology. *J Appl Physiol* 1995;**79**:1525–30.
16. **Hogg JC**, Nepszky S. Regional lung volume and pleural pressure gradient estimated from lung density in dogs. *J Appl Physiol* 1969;**27**:198–203.
17. **Crapo RO**, Morris AH, Clayton PD, *et al*. Lung volumes in healthy nonsmoking adults. *Bull Eur Physiopathol Respir* 1982;**18**:419–25.
18. **Devine BJ**. Gentamicin therapy. *Drug Intell Clin Pharm* 1974;**8**:650–5.
19. **Shohl AT**. *Mineral metabolism*. New York: Reinhold Publishing Corporation, 1939.
20. **Gevenois PA**, De Vuyst P, de Maertelaer V, *et al*. Comparison of computed density and microscopic morphometry in pulmonary emphysema. *Am J Respir Crit Care Med* 1996;**154**:187–92.
21. **Nakano Y**, Muro S, Sakai H, *et al*. Computed tomographic measurements of airway dimensions and emphysema in smokers. Correlation with lung function. *Am J Respir Crit Care Med* 2000 Sep;**162**(3 Pt 1):1102–8.
22. **Nakano Y**, Wong JC, de Jong PA, *et al*. The prediction of small airway dimensions using computed tomography. *Am J Respir Crit Care Med* 2005;**171**:142–6.
23. **Feldman HA**. Families of lines: random effects in linear regression analysis. *J Appl Physiol* 1988;**64**:1721–32.
24. **Leopold JG**, Gough J. The centrilobular form of hypertrophic emphysema and its relation to chronic bronchitis. *Thorax* 1957;**12**:219–35.
25. **McLean KH**. The histology of localized emphysema. *Australas Ann Med* 1957;**6**:282–94.
26. **Casanova C**, Cote C, de Torres JP, *et al*. Inspiratory-to-total lung capacity ratio predicts mortality in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2005;**171**:591–7.
27. **Ofir D**, Laveneziana P, Webb KA, *et al*. Mechanisms of dyspnea during cycle exercise in symptomatic patients with GOLD stage I chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2008;**177**:622–9.
28. **O'Donnell DE**. Hyperinflation, dyspnea, and exercise intolerance in chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 2006;**3**:180–4.
29. **Remy-Jardin M**, Edme JL, Boulenguez C, *et al*. Longitudinal follow-up study of smoker's lung with thin-section CT in correlation with pulmonary function tests. *Radiology* 2002;**222**:261–70.
30. **Parr DG**, White AJ, Bayley DL, *et al*. Inflammation in sputum relates to progression of disease in subjects with COPD: a prospective descriptive study. *Respir Res* 2006;**7**:136.
31. **Stolk J**, Putter H, Bakker EM, *et al*. Progression parameters for emphysema: a clinical investigation. *Respir Med* 2007;**101**:1924–30.
32. **Hogg JC**, Chu F, Utokaparch S, *et al*. The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* 2004;**350**:2645–53.
33. **Hasegawa M**, Nasuhara Y, Onodera Y, *et al*. Airflow limitation and airway dimensions in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2006;**173**:1309–15.
34. **Burrows B**, Knudson RJ, Camilli AE, *et al*. The "horse-racing effect" and predicting decline in forced expiratory volume in one second from screening spirometry. *Am Rev Respir Dis* 1987;**135**:788–93.
35. **Celli BR**, Cote CG, Marin JM, *et al*. The body-mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *N Engl J Med* 2004;**350**:1005–12.

Lung alert

The *sst1* locus controls granuloma necrosis in tuberculosis

There is broad variation in susceptibility to *Mycobacterium tuberculosis* (MTB), and mouse models play a key role in elucidating the underlying mechanisms. The authors tested the hypothesis that caseation within pulmonary lesions is a specific effect of the *sst1* (supersusceptibility to tuberculosis 1) locus on chromosome 1.

Typical mouse lesions described in the literature lack central caseation. However, the C3HeB/FeJ strain of mouse develops large necrotising granulomas after exposure to MTB. The authors compared the course of infection in an MTB-resistant mouse strain (B6) with an *sst1*-susceptible congenic strain (B6.C3H-*sst1*) which was genetically identical except for the interval containing the *sst1* locus.

They found that, although initial dissemination was similar, bacterial loads, clinical disease, lung necrosis and mortality were considerably worse in the strain containing the *sst1*-susceptible allele. In addition, relapse after 3 months of isoniazid was faster and more rampant in this group. After demonstrating significantly slower disease progression and milder histopathology in the B6.C3H-*sst1* strain compared with the C3HeB/FeJ strain, the authors concluded that the effect of the *sst1* locus is modified by the genetic background of the host. Enhanced pro-inflammatory cytokine production by macrophages was observed in the susceptible strains; however, this appears to be controlled by loci other than *sst1*.

Further characterisation of *sst1*-encoded molecular mechanisms may not only shed light on a key aspect of the pathogenesis of tuberculosis, but may also suggest therapeutic interventions to reduce lung pathology and transmission of the pathogen.

- Pichugin AV, Yan B, Sloutsky A, *et al*. Dominant role of the *sst1* locus in pathogenesis of necrotizing lung granulomas during chronic tuberculosis infection and reactivation in genetically resistant hosts. *Am J Pathol* 2009;**174**:2190–201

M Almond

Correspondence to: Dr M Almond, ST1 Oncology, Royal Marsden Hospital, London, UK; mhalmond@gmail.com

Provenance and peer review: Commissioned; not externally peer reviewed.

Thorax 2009;**64**:949. doi:10.1136/thx.2009.122879