Quorum sensing-dependent virulence during *Pseudomonas aeruginosa* colonisation and pneumonia in mechanically ventilated patients

Thilo Köhler,^{1,2} Raphael Guanella,³ Jean Carlet,⁴ Christian van Delden^{1,2}

► Supplementary patient information and methods are published online only. To view these files please visit the journal online (<http://thorax.bmj.com>).

¹Service of Infectious Diseases, University Hospital Geneva, Switzerland

²Department of Microbiology and Molecular Medicine, Faculty of Medicine, University of Geneva, Switzerland

³Service of Hemostasis, University Hospital Geneva, Switzerland

⁴Intensive Care Unit, Groupe Hospitalier Paris Saint-Joseph, France

Correspondence to

Christian van Delden, Service of Infectious Diseases, University Hospital Geneva, 4 Rue Gabrielle-Perret-Gentil, CH-1211 Geneva 14, Switzerland; christian.vandelden@unige.ch

Presented in part at the Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC), Washington, DC, 25–28 October 2008 (abstract A 2521).

Received 14 December 2009
Accepted 12 May 2010

ABSTRACT

Background *Pseudomonas aeruginosa* frequently colonises intubated patients and causes life-threatening ventilator-associated pneumonia (VAP). The role of quorum sensing (QS), regulating virulence in this pathogen, during colonisation and development of VAP is unknown.

Methods *P. aeruginosa* isolates and tracheal aspirates were prospectively collected from intubated patients. Genotypes and QS-independent virulence traits (*exoU*, *exoS*, PAPI-1 and PAPI-2) harboured by colonising isolates were identified in vitro with the CLONDIAG array. The production of elastase and rhamnolipids was measured to assess QS-dependent virulence. To monitor QS activity 'in patient', total RNA was extracted directly from tracheal aspirates and expression of QS genes was measured.

Results 320 *P. aeruginosa* isolates and tracheal aspirates were obtained from 29 patients of whom 6 developed VAP (20%). Seven patients (24%) were initially colonised by QS-proficient isolates; 57% of them developed VAP as compared with 9% of patients colonised by QS-deficient isolates ($p=0.018$). Of all tested virulence traits from the initial colonising isolates, only rhamnolipids were associated with development of VAP ($p=0.003$). VAP occurred more frequently in patients colonised during the entire observation period by isolates producing high levels of rhamnolipids ($p=0.001$). 'In patient' monitoring of QS genes showed non-induced expression profiles in patients without VAP. In contrast, exponential induction of QS circuit and target gene expression was observed for two patients with VAP, and an 'in patient' QS gene expression profile and hierarchy similar to those in vitro was measured for one patient with VAP.

Conclusions Production of the QS-dependent virulence factor rhamnolipids by colonising *P. aeruginosa* isolates is associated with development of VAP.

INTRODUCTION

Pseudomonas aeruginosa frequently colonises intubated patients. In 10–20% of these patients colonisation progresses to ventilator-associated pneumonia (VAP) associated with mortality rates of 30–40%.¹ To date, it has been impossible to accurately identify those colonised patients who will develop this complication. In particular it remains unclear whether patients prone to infection are colonised by especially virulent strains. A quorum-sensing (QS) circuit regulates most virulence determinants in *P. aeruginosa* according to cell density.^{2–6} These include many secreted factors (elastase, phospholipase C, lecithinase, rhamnolipids) as well as secondary metabolites (pyocyanin,

cyanide).^{7–8} The importance of QS-regulated virulence traits for the pathogenicity of *P. aeruginosa* has been established in both animal and plant infection models.⁹ However, evidence for QS activity in the human host remains scarce and is derived from the detection of QS signalling molecules and the detection of QS gene expression in sputum samples collected occasionally from patients with cystic fibrosis.^{10–15} Its relevance in acute human infections has even been questioned recently.^{16–17} If QS-dependent virulence traits were important for the development of infection, then preventive measures aiming at the inhibition of the QS circuit would be an attractive alternative to conventional antibiotics, which frequently lead to the selection of resistance without being able to eradicate colonisation of intubated patients.^{1–4–18}

Our primary aim was to determine the presence of particular virulence traits in *P. aeruginosa* isolates colonising intubated patients and their potential association with the development of *Paeruginosa* VAP. We also sought to determine whether induction of the QS circuit could be demonstrated 'in patient' during colonisation and/or infection of intubated patients by monitoring bacterial gene expression directly in tracheal aspirates.

METHODS

Subjects

This study is a subanalysis of a multicentre placebo-controlled trial investigating azithromycin as a QS inhibitor for the prevention of *P. aeruginosa* VAP in colonised mechanically ventilated patients (<http://ClinicalTrials.gov> ID# NCT00610623). Detailed information on the results of this clinical trial are published elsewhere (van Delden *et al*, submitted, 2010). Patients of the present study were restrictively selected from the placebo-control group on the basis of availability of both daily clinical *P. aeruginosa* isolates and tracheal aspirates. We screened intubated patients for respiratory tract colonisation by *P. aeruginosa* every 48 h. Patients with ongoing *P. aeruginosa* infection, or having received antimicrobial treatments active against the colonising isolates during the last 14 days, or during the observation period, were not included. The diagnosis of *Paeruginosa* VAP was based on the clinical picture, including a pulmonary infection score (CPIS) ≥ 6 , as well as a quantitative culture of a bronchoalveolar lavage fluid yielding $>10^4$ CFU/ml of *Paeruginosa*.¹⁹

Collection of respiratory samples

Starting the first day of proven colonisation (D0), we prospectively collected daily tracheal aspirates

and one *P. aeruginosa* isolate (collection period: 3–20 days). Samples were frozen at –80°C on site within 15 min, and sent on dry ice to the research laboratory at the University Hospital Geneva, where all RNA and DNA extractions were performed. Tracheal aspirates were used to determine *P. aeruginosa* bacterial loads and ‘in patient’ expression of QS genes, whereas *P. aeruginosa* isolates were used for genotypic and phenotypic characterisations (figure 1A). Reasons to stop sample collection were either extubation or proven *P. aeruginosa* VAP.

Genomic analysis of clinical isolates

Inpatient comparison of genotypes was performed on the entire collection of 320 isolates by random amplification of polymorphic DNA (RAPD) using primer 207.²⁰ Interpatient comparison of the genotype of the initial isolate from each of

the 29 patients was performed with the *P. aeruginosa* CLON-DIAG array.²¹ The array further detects the presence of 38 genetic markers of the accessory genome, including two genes from the pathogenicity island PAPI-1 and three from PAPI-2.²²

The *lasR* and *rhlR* genes were amplified by PCR using primers *lasR-F1* (5'-ATCTTGTGGGCTGACTGGAC-3') and *lasR8* (5'-CTGGCGCAGTCGTTTCGAGAAT-3'), and *rhlR-F1* (5'-ACGGT GCTGGCATAACAGATAGG-3'); and *rhlR-R1* (5'-CCTCTCAG TCGGAGGACATACCA-3'), respectively, from bacterial lysates. PCR conditions were as follows: 95°C for 2 min, 27 cycles at 95°C for 15 s, 57°C for 30 s and 72°C for 1 min, and a final extension at 72°C for 4 min. *lasR* and *rhlR* PCR products were purified (Qiagen, Hilden, Germany) and sequenced using primers *lasR-F1* and *rhlR-F1*, respectively.

Exoproduct analysis

Elastase activity was determined using the Elastin Congo Red assay in supernatants of cultures grown for 7 h at 37°C in PB medium.^{23 24} Rhamnolipid production was assessed on modified SW-Blue plates.²⁵ The diameter of the rhamnolipid-containing halo formed around the bacterial colony after 24 h incubation at 37°C and after 24 h at room temperature was measured and compared with that produced by the reference strain PAO1. All

Figure 1 (A) Flowchart for procedures on tracheal aspirates. From each daily aspirate one *Pseudomonas aeruginosa* isolate was selected, and DNA as well as RNA were extracted in parallel. Bacterial load was deduced from genomic DNA by quantitative reverse transcription–PCR (qRT–PCR) and gene expression was determined after reverse transcription of total RNA. (B) Bacterial load over time. Shown are the mean±SEM of bacterial loads for patients with ventilator-associated pneumonia (VAP) (filled squares) (n=5, genomic samples of patient 13128 with VAP were not available due to technical problems) and patients without VAP (open squares) (n=23). Arrows indicate days of VAP (dotted arrow indicates patient 13128, for whom no genomic DNA samples were available).

Patient	% of PAO1		sequence		Exo S/U	PAPI-1	PAPI-2	SNP type
	Ela	Rha	<i>lasR</i>	<i>rhlR</i>				
PAO1	100	100	wt	wt	S	0	0	000A
05101	151	116	wt	wt	S	0	0	6C26
19105	153	110	wt	wt	S	0	1	7C2E
16101	189	104	wt	wt	S	0	0	0C2E
13128	137	103	wt	wt	S	0	0	0C2E
13111	109	101	wt	wt	S	0	2	85AA
30101	345	91	wt	wt	U	2	3	D421
24101	292	102	wt	wt	S	0	1	2C26
21107	24	91	A231V	wt	S	0	2	6D92
13108	61	88	wt	wt	S	0	1	1BAE
13104	190	86	wt	wt	S	0	2	0C1A
26102	152	75	wt	wt	U	0	2	F469
13112	123	73	wt	wt	S	0	2	C40A
13122	122	90	wt	wt	U	0	2	F661
15102	215	79	wt	wt	U	0	1	E429
13106	127	72	wt	wt	S	0	2	4F8A
19101	34	80	Δ or IS	wt	U	0	2	F469
19102	47	80	A231V	wt	S	0	1	6D92
27104	24	78	L110Q	wt	S	0	2	0812
13118	42	41	wt	wt	S	0	2	239A
10103	35	0	199IS	wt	U	2	3	D421
15101	89	0	wt	wt	S	0	1	0C2E
13121	16	0	wt	wt	S	0	2	239A
26104	25	0	wt	T121I	U	0	2	F469
13114	28	0	L148P	Δ64 bp	U	0	2	F469
13117	19	0	L148P	Δ64 bp	U	0	2	F469
06104	0	0	47 IS	wt	S	0	2	AF9A
27101	1	0	Δ or IS	A111D	S	0	0	EC4A
22101	1	0	P74L	Δ or IS	U	0	2	F469
13116	1	0	T222I	Δ64 bp	U	0	2	F469
PA14	99	95	wt	wt	U	2	3	D421

% of PAO1		
Ela,Rha	>90%	90-10% <10%

Figure 2 Phenotypic and genotypic characterization of initial colonising *Pseudomonas aeruginosa* isolates. Production of elastase (Ela) and rhamnolipids (Rha) was scored according to the classification scheme. Alterations to the PAO1 (wild-type) sequences of the *lasR* and *rhlR* genes, as well as undefined deletions (Δ) or insertions (IS) are indicated. The presence of the cytotoxicity genes *exoS* and *exoU* from PAPI-1 (maximum of two genes) and PAPI-2 (maximum of three genes), as well as the single nucleotide polymorphism (SNP) type, were established using the CLONDIAG array. Patients with ventilator-associated pneumonia (VAP) are highlighted by black boxes.

Table 1 Virulence determinants in patients with and without ventilator-associated pneumonia (VAP)

Expression or presence of virulence determinant (total number of patients with a positive initial isolate)	VAP (n=6)	Non-VAP (n=23)	p Value
Elastase (13)*	4	9	0.364
Rhamnolipid (8)*	5	3	0.003
<i>exoS</i> (18)†	5	13	0.362
<i>exoU</i> (11)†	1	10	0.362
PAPI-1 (2)†	0	2	1.000
PAPI-2 (25)†	4	21	0.180

*≥90% activity of reference strain PAO1.

†Presence of the gene was determined using the CLONDIAG array; pathogenicity islands were considered to be present in the isolate when at least one of two (PAPI-1) and one of three (PAPI-2) genes were detected by the CLONDIAG array.

determinations were done in duplicate and expressed as a percentage of PAO1 production levels.

DNA and RNA preparation

Genomic DNA, and RNA were extracted from 0.05 ml and 0.5 ml of homogenised tracheal aspirates, respectively (figure 1A). Additional details on the extraction procedures and gene expression analyses are provided in the online data supplement.

Statistical analysis

This study is a subanalysis of a multicentre placebo-controlled trial. None of the analysis presented here was included in the primary or secondary outcomes of this trial, which was focused on the occurrence of *P. aeruginosa* VAP (primary outcome) and

A

Patient	Days of colonization																				
high (n=6)	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
5101	151	137	141	163	134																
13106	133	127	120	111	137	123	104	163	126	122	NI	NI	111	94	NI	103	102	104	115	115	128
13111	109	114	112	118	111																
13112	123	111	105	118	115	116	120	120													
13128	137	136	140	131	108																
13104	190	157	185	204	186	197															
AVG	141	130	134	141	132	145	112	141	126	122	NI	NI	111	94	NI	103	102	104	115	115	128
variable (n=8)	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
19105	153	166	169	162	135	118	5	4	22	117	78	90	4	60	59	138	166	76	11	64	83
15101	89	90	22	NI	1	NI	1	1	1	19	2	1	56	57	2	0	15	2	16		1
15102	215	164	3	188	145	182	171	4	175	3	12	5	4	193	118	2					
24101	292	301	56	23	304	NI	297	141	48	271											
26102	152	0	0																		
16101	189	165	60	21	20	25	27	1	20	0	23	21									
30101	345	138	220	173	140	23															
13122	122	NI	132	136	144	142	69	121	145	34	29	4	6	101	1	56	63	8	144	105	7
AVG	195	146	83	117	127	98	95	45	68	74	29	24	18	103	45	49	81	28	57	85	48
low (n=15)	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
6104	0	0	0	0	0	0	1	NI	1	1	NI	1	1	1	1	1					
13116	1	1	1	2	1	2	2	2	1	2											
22101	1	1	0	1	0	NI	1														
27101	1	1	NI	1	ND	0	1	NI	0	ND	ND	ND	1	ND	1	ND	1	NI	0	1	
10103	35	42	43	36	37	NI	4	5	47	6											
13108	61	57	44	70	37	22	30	81	44	45	77	68	49	39	5						
13114	28	22	25	23	22	29	24	NI	23	19	31	24	28								
13117	19	20	0	0	9	14	1	0	1	0	2	5	0	19							
13118	12	67	13	37	21	28	52	25	15												
13121	16	50	3	2	3	49	2														
19101	34	6	26	36	41	62	25	9	49	38	8	35	20	18	6	30	15	16	30	NI	3
19102	47	60	57	43	58	12	39	16	16	53	36	51	39								
21107	24	22	23	30	23	38	23	16	43	21	68										
26104	25	25	23	24	26	185	26	10	11	1	141										
27104	24	24	22	24	10	18	4	22													
AVG	22	26	20	22	21	35	16	19	21	19	52	31	20	19	3	15	8	16	15	1	3

NI, no isolate; ND, not done; AVG, average

B

Average Elastase activity (n=29)

Figure 3 Elastase production of colonising isolates from 29 patients. (A) Elastase production is expressed as a percentage of the reference strain PAO1. Patients were allocated to three different classes according to the temporal evolution of elastase production of their isolates (high, variable and low). The average (AVG) for each group is shown. Patients who developed ventilator-associated pneumonia (VAP) are shown in bold. (B) Temporal evolution of average elastase production of isolates from the three different classes shown in A. Evolution is shown only up to day 11 due to low sample sizes thereafter.

mortality (secondary outcome) in both placebo and azithromycin study arms. Two-sided Fisher exact tests were used to analyse differences between group proportions. The Mann–Whitney test was used as a non-parametric test.

RESULTS

Genotypes and virulence phenotypes of initial colonising *P aeruginosa* isolates

The collection included a total of 320 *P aeruginosa* isolates from 29 patients. Six patients developed *P aeruginosa* VAP. Detailed clinical information on the study population is presented in the online data supplement. Four patients (13122, 15101, 21107 and

26102) were colonised by two different genotypes. We first focused on the 29 initial colonising isolates. Using the *P aeruginosa* CLONDIAG genome array, we distinguished 18 clones (figure 2).²¹ The CLONDIAG array also allows determination of the presence of virulence genes including the *Pseudomonas* pathogenicity islands PAPI-1 and PAPI-2, as well as the two mutually exclusive cytotoxicity genes *exoS* and *exoU*. The genes of PAPI-1 were detected in only two isolates. In contrast, genes of PAPI-2 were found in 25 out of the 29 isolates (figure 2). The gene encoding the cytotoxic phospholipase ExoU, carried on PAPI-2, was present in isolates of 11 patients (38%), while *exoS*, not located on a pathogenicity island, was identified in isolates

NI, no isolate; AVG, average

Figure 4 Rhamnolipid production of colonising isolates from 29 patients. (A) Rhamnolipid production is expressed as a percentage of the reference strain PAO1. Patients were allocated to three different classes according to the temporal evolution of rhamnolipid production of their isolates (high, intermediate and low). The average (AVG) for each group is shown. Patients who developed ventilator-associated pneumonia (VAP) are shown in bold. (B) Temporal evolution of average rhamnolipid production of isolates from the three different classes shown in A. Evolution is shown only up to day 11 due to low sample sizes thereafter. (C) Comparison of average rhamnolipid production per patient during their entire observation period between the VAP (n=6) and non-VAP group (n=23).

from the other 18 patients (62%). Twenty-two of the initial 29 isolates (76%) produced reduced amounts of the QS-dependent virulence factors elastase and/or rhamnolipids compared with the laboratory strain PAO1 (figure 2). Sequencing revealed that lack of production of these QS-dependent virulence factors correlated for most isolates with mutations in the *lasR* and/or the *rhlR* QS-regulator genes (figure 2).²⁶

We further compared genotypes and virulence phenotypes of the initial colonising isolates from the six patients with VAP with those from the 23 patients without VAP. We found no statistically significant association between VAP and a particular genotype, or the presence of either of the *Pseudomonas* pathogenicity islands PAPI-1 and PAPI-2, or of the cytotoxicity genes *exoU* or *exoS* (table 1). In contrast four out of seven patients (57%) initially colonised by QS-proficient isolates (producing both elastase and rhamnolipids at levels comparable with PAO1) developed VAP as compared with two out of 22 patients (9%) colonised by isolates with reduced QS activities ($p=0.018$). High level production of rhamnolipids ($p=0.003$) but not of elastase was statistically associated with VAP (figure 2 and table 1).

P. aeruginosa virulence determinants during colonisation and VAP

Virulence phenotypes fluctuate during colonisation.²⁶ We therefore measured both elastase and rhamnolipid production in the 320 daily isolates from the 29 patients. According to the daily elastase production, we identified three distinct patterns. Fifteen patients were consistently colonised by low elastase-producing isolates (figure 3A, B). The average daily elastase production value in this group fluctuated between 1% and 52% of the reference strain PAO1. Eight patients were colonised by isolates producing fluctuating intermediate amounts of elastase. This category displayed a clear tendency to decreasing daily average elastase production from 195% (D0) to 24% (D11) (figure 3A, B). Six patients were colonised during their entire observation period by isolates producing high levels of elastase, with average values between 102% and 146% (figure 3A, B). The six patients with VAP (13111, 13128, 16101, 21107, 24101 and 13116) were equally distributed in all three categories, suggesting no association between elastase production of the colonising isolates and VAP. The average per patient elastase production was not different between the VAP and non-VAP groups.

The analysis of the daily rhamnolipid production levels also allowed separation of the 29 patients into three distinct groups (figure 4A, B). Over the study period, 11 patients remained colonised by non-rhamnolipid-producing isolates, with daily average values between 0 and 20% of the reference strain PAO1 (figure 4A, B). Thirteen patients were colonised by isolates producing intermediate amounts of rhamnolipids, with daily average values between 59% and 76% (figure 4A, B). Five patients (13111, 13128, 16101, 21107, 24101) were colonised during the entire study period by isolates producing high rhamnolipid levels, with average daily values between 76% and 100% of the PAO1 reference strain (figure 4A, B). All these five patients developed VAP. The sixth patient with VAP (13116), colonised by a *lasR/rhlR* double mutant, developed pneumonia in the absence of detectable rhamnolipid production. Thus VAP occurred more frequently in the group of patients colonised during the entire study period by isolates producing high rhamnolipid levels than in the other two groups ($p=0.001$). Even when patient 13116 was included in the analysis, patients with VAP were colonised over time with isolates producing higher average levels of rhamnolipids than patients without VAP (Mann–Whitney test, $p=0.014$) (figure 4C).

'In patient' QS induction

To obtain a global picture of QS activity in vivo, we analysed *P. aeruginosa* QS gene expression at the population level by extracting total genomic DNA and total RNA from daily tracheal aspirates (figure 1A). As QS is cell density dependent, we determined the global bacterial lung loads using the genomic DNA. In all patients, the daily bacterial load fluctuated within a range of 10^6 – 10^8 genomic copies/g of aspirate without a clear trend (figure 1B). Except for a transient two- to threefold increase in patients with VAP between the second and fourth day of colonisation, no difference in bacterial load was observed between those with and without VAP (figure 1B). Since all VAP cases (indicated by arrows) occurred on or before day 11, a comparison beyond this point was not possible.

We assessed the expression of two transcriptional regulator genes (*lasR* and *rhlR*), and of two autoinducer synthases genes (*lasI* and *rhlI*), which together form the QS circuit genes of *P. aeruginosa*, as well as of two major QS target genes *lasB* (elastase) and *rhlA* (rhamnosyltransferase).⁴ We detected expression of *lasI* and *rhlA* in 70% and 42%, respectively, of all aspirates (see

Figure 5 'In patient' quorum sensing (QS) induction. (A) Temporal evolution of *lasI* expression. Three patients with ventilator-associated pneumonia (VAP): 13111 (filled circles), 13128 (filled diamonds) and 21107 (filled triangles), and 19 patients without VAP (open diamonds). (B) Temporal evolution of *rhlA* expression. Three patients with VAP: 13111 (filled circles), 13128 (filled diamonds) and 21107 (filled triangles), and 16 patients without VAP (squares). Gene expression is standardised towards expression of the housekeeping gene *rpsL*.

online data supplement). The expression of *lasB* and *rhII* was measured at lower levels and in fewer samples, excluding these genes from longitudinal analysis. We therefore analysed the mean *lasI* (figure 5A) and *rhIA* (figure 5B) expression levels over time within 19 and 16 patients without VAP, respectively, from whom adequate samples for QS gene expression were available using the QS-independent housekeeping gene *rpsL* to normalise gene expression between samples. Both curves showed a similar stable non-induced profile compatible with basal gene expression. Adequate samples for QS gene expression profiles were available for four patients with VAP (13111, 13116, 13128 and 21107). Patients 13111 and 21107 with VAP showed clear QS induction profiles for both *lasI* (figure 5A) and *rhIA* (figure 5B) with an exponential increase of QS gene expression starting 48 h before VAP. Expression levels of patient 13128 with VAP remained comparable with means of patients without VAP. Consistent with the low rhamnolipid production of its colonising isolates (figure 4A), patient 13116 with VAP colonised by a double *lasR/rhlR* mutant did not show any QS induction.

To highlight the previously described cell density dependency and hierarchy of QS gene induction, we measured the expression of the QS circuit (*lasR*, *lasI*, *rhlR* and *rhlI*) and QS target (*lasB* and *rhIA*) genes during in vitro growth of the reference strain PAO1 (figure 6A).²⁷ As expected QS induction occurred only once the bacterial cell density had reached a threshold level (1×10^9 CFU/ml under our conditions), and the fold induction of the QS target genes *lasB* and *rhIA* was higher than that of the QS

regulator genes *lasR* and *rhlR*. In patient 13111 with VAP we were able to measure the expression of the same QS target and circuit genes over time (figure 6B). In agreement with in vitro data, the *rhl* QS system genes were more induced than those of the *las* system in this patient.²⁸ We observed the highest induction in gene expression (100-fold) for the QS target gene *rhIA*. In contrast to the in vitro situation, the bacterial load in the tracheal aspirates did not increase during the observation period (see inset in figure 6B).

DISCUSSION

This is the first study to report on QS-dependent virulence in *P aeruginosa* isolates in vitro at the clonal, as well as 'in patient' at the bacterial population level, during both colonisation and development of VAP. We found that (1) the majority of patients were initially colonised by QS-deficient *P aeruginosa* isolates; (2) VAP occurred more frequently in patients colonised by QS-proficient isolates; (3) the production of rhamnolipids, both by the initial colonising isolate and the subsequent colonisers, was associated with development of VAP; and (4) 'in patient' QS gene induction was observed only in patients developing VAP.

Our study included 29 colonised patients of which six developed VAP. Only 24% of our patients were initially colonised by fully QS-proficient *P aeruginosa* isolates. The incidence of VAP was much higher in this group (57%) as compared with the rest of the study population (9%). Moreover all patients initially and persistently colonised during the entire observation period by

Figure 6 In vitro and 'in patient' quorum sensing (QS) induction. (A) In vitro QS circuit induction. Growth curve of the *P aeruginosa* wild-type strain PAO1 in LB medium (figure inset). Expression (fold change of cDNA copies/ml culture) of QS circuit genes (*lasR*, *rhlR*, *lasI* and *rhlI*), QS target genes (*rhIA* and *lasB*) and the housekeeping gene (*rpsL*) determined by quantitative reverse transcription-PCR (qRT-PCR). (B) 'In patient' QS circuit induction. Daily QS gene expression of patient 13111 expressed as fold change of cDNA copies/g of aspirate with respect to day 1. Bacterial cell density (inset).

isolates producing high levels of rhamnolipids developed VAP, supporting the assertion that this particular QS-dependent virulence trait plays a role in the development of this infection. Several mechanisms by which rhamnolipids can promote infection have been proposed. Rhamnolipids are necessary and sufficient for invasion of a reconstituted human airway epithelium; they modulate biofilm architecture and lyse polymorphonuclear neutrophils.^{29–31} Taken together these data designate rhamnolipids as an essential factor promoting the early stages of mucosal invasion. However, because of the low patient number, our data do not exclude that other QS-dependent factors also play a role during colonisation and infection in intubated patients. Strikingly we did not find a direct association between elevated elastase production and the occurrence of VAP. However, we have shown previously that VAP occurs earlier in patients colonised by elastase-producing isolates, as compared with patients colonised by isolates with reduced or no elastase production.²⁶ Markedly, while rhamnolipid production remained fairly stable over time, average elastase production decreased. This is in agreement with a constant proportion of *rhlR* mutants (*rhl* mainly regulating rhamnolipid production) but an increasing proportion of *lasR* mutants (*lasR* mainly regulating elastase production) colonising the patients over time.²⁶

While the importance of QS-dependent virulence for pathogenesis has been established in different host models and is further supported by our *in vitro* data, its relevance needed to be further investigated directly in the human host.⁹ Our study provides the first 'in patient' view of QS dynamics. In patients without VAP, *lasI* and *rhlA* were only expressed at a basal, non-induced level during colonisation. In contrast, we observed global QS induction in two colonised patients progressing towards VAP. Although the number of VAP cases is not sufficient to achieve a statistically significant association with 'in patient' QS induction, this observation illustrates QS activity *in vivo* during development of VAP. As 'in patient' gene expression data reflect the mean of a global bacterial population, QS induction in spatially restricted areas such as microcolonies, where VAP might start initially, could potentially be missed. This could explain why we were unable to detect QS induction in the other patients with VAP.

In one patient, colonised by QS-proficient isolates, we observed a similar 'in patient' QS gene expression profile and hierarchy to that *in vitro*, illustrating that QS can also orchestrate *P. aeruginosa* gene expression in the lungs of the human host. This 'in patient' QS induction occurred without an increase in bacterial cell density and coincided with the development of pneumonia. In contrast, one patient with VAP was colonised by a double *lasR/rhlR* mutant. Interestingly, isolates of this patient carried the *exoU* gene, encoding a phospholipase whose expression has been associated with increased mortality from VAP.³² Since QS negatively regulates TTSS expression, these isolates might have achieved higher or earlier expression of TTSS, and development of VAP might have been mediated by cytotoxicity in this particular case.³⁵ Whether colonisation with *exoU* isolates per se is a risk factor for progression from colonisation to VAP has not been investigated, but our results are not in favour of this hypothesis.

In conclusion, this study shows that the production of rhamnolipids by colonising isolates is associated with the development of VAP in intubated patients. Further studies of the potential benefit of inhibiting this QS-dependent virulence factor in colonised intubated patients are warranted. Macrolide antibiotics, furanone derivatives and N-acylhomoserine (HSL) analogues have been shown *in vitro* to reduce QS gene expression

and the production of QS-dependent virulence factors, and might be good candidates as QS inhibitors.^{23 34 35} Such studies would provide the scientific basis for bacterial virulence inhibition to prevent *P. aeruginosa* VAP in intubated patients.

Acknowledgements We express our gratitude to the responsible physicians, especially B Francois, the nurses and medical staff involved in the collection of clinical samples and to F. Brunner-Ferber. We thank R Comte and J L Dumas for outstanding technical assistance.

Funding This work was supported by the Swiss National Science Foundation (grants 4049-063239 to TK and CVD, 320000-108106 and 320000-120011 to CVD), and partially by the Research Fund of the Department of Internal Medicine of the University Hospital and the Faculty of Medicine of Geneva; this Fund receives an unrestricted grant from AstraZeneca Switzerland, Pfizer Switzerland and the Wilsdorf Foundation. The clinical collection was financed by Anbics Corporation.

Competing interests None.

Ethics approval This study was conducted with the approval of the Ethics Committee of the Department of Medicine of the University Hospital Geneva and thereafter from all local ethical committees.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. Chastre J, Fagon JY. Ventilator-associated pneumonia. *Am J Respir Crit Care Med* 2002;**165**:867–903.
2. Pearson JP, Gray KM, Passador L, et al. Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes. *Proc Natl Acad Sci USA* 1994;**91**:197–201.
3. Pearson JP, Passador L, Iglewski BH, et al. A second N-acylhomoserine lactone signal produced by *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 1995;**92**:1490–4.
4. Van Delden C, Iglewski BH. Cell-to-cell signaling and *Pseudomonas aeruginosa* infections. *Emerg Infect Dis* 1998;**4**:551–60.
5. Wagner VE, Bushnell D, Passador L, et al. Microarray analysis of *Pseudomonas aeruginosa* quorum-sensing regulons: effects of growth phase and environment. *J Bacteriol* 2003;**185**:2080–95.
6. Schuster M, Lostroh CP, Ogi T, et al. Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: a transcriptome analysis. *J Bacteriol* 2003;**185**:2066–79.
7. Sadikot RT, Blackwell TS, Christman JW, et al. Pathogen–host interactions in *Pseudomonas aeruginosa* pneumonia. *Am J Respir Crit Care Med* 2005;**171**:1209–23.
8. Schuster M, Greenberg EP. A network of networks: quorum-sensing gene regulation in *Pseudomonas aeruginosa*. *Int J Med Microbiol* 2006;**296**:73–81.
9. Rumbaugh KP, Griswold JA, Hamood AN. The role of quorum sensing in the *in vivo* virulence of *Pseudomonas aeruginosa*. *Microbes Infect* 2000;**2**:1721–31.
10. Singh PK, Schaefer AL, Parsek MR, et al. Quorum-sensing signals indicate that cystic fibrosis lungs are infected with bacterial biofilms. *Nature* 2000;**407**:762–4.
11. Geisenberger O, Givskov M, Riedel K, et al. Production of N-acyl-L-homoserine lactones by *P. aeruginosa* isolates from chronic lung infections associated with cystic fibrosis. *FEMS Microbiol Lett* 2000;**184**:273–8.
12. Favre-Bonte S, Pache JC, Robert J, et al. Detection of *Pseudomonas aeruginosa* cell-to-cell signals in lung tissue of cystic fibrosis patients. *Microb Pathog* 2002;**32**:143–7.
13. Storey DG, Ujack EE, Rabin HR, et al. *Pseudomonas aeruginosa lasR* transcription correlates with the transcription of *lasA*, *lasB*, and *tox4* in chronic lung infections associated with cystic fibrosis. *Infect Immun* 1998;**66**:2521–8.
14. Erickson DL, Endersby R, Kirkham A, et al. *Pseudomonas aeruginosa* quorum-sensing systems may control virulence factor expression in the lungs of patients with cystic fibrosis. *Infect Immun* 2002;**70**:1783–90.
15. Son MS, Matthews WJ Jr, Kang Y, et al. *In vivo* evidence of *Pseudomonas aeruginosa* nutrient acquisition and pathogenesis in the lungs of cystic fibrosis patients. *Infect Immun* 2007;**75**:5313–24.
16. Denervaud V, TuQuoc P, Blanc D, et al. Characterization of cell-to-cell signaling-deficient *Pseudomonas aeruginosa* strains colonizing intubated patients. *J Clin Microbiol* 2004;**42**:554–62.
17. Cabrol S, Olliver A, Pier GB, et al. Transcription of quorum-sensing system genes in clinical and environmental isolates of *Pseudomonas aeruginosa*. *J Bacteriol* 2003;**185**:7222–30.
18. Smith RS, Iglewski BH. *Pseudomonas aeruginosa* quorum sensing as a potential antimicrobial target. *J Clin Invest* 2003;**112**:1460–5.
19. Pugin J, Auckenthaler R, Mili N, et al. Diagnosis of ventilator-associated pneumonia by bacteriologic analysis of bronchoscopic and nonbronchoscopic "blind" bronchoalveolar lavage fluid. *Am Rev Respir Dis* 1991;**143**:1121–9.
20. Mahenthiralingam E, Campbell ME, Foster J, et al. Random amplified polymorphic DNA typing of *Pseudomonas aeruginosa* isolates recovered from patients with cystic fibrosis. *J Clin Microbiol* 1996;**34**:1129–35.
21. Wiehlmann L, Wagner G, Cramer N, et al. Population structure of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 2007;**104**:8101–6.

22. **He J**, Baldini RL, Deziel E, *et al.* The broad host range pathogen *Pseudomonas aeruginosa* strain PA14 carries two pathogenicity islands harboring plant and animal virulence genes. *Proc Natl Acad Sci U S A* 2004;**101**:2530–5.
23. **Tateda K**, Comte R, Pechere JC, *et al.* Azithromycin inhibits quorum sensing in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2001;**45**:1930–3.
24. **Essar DW**, Eberly L, Crawfords IP. Identification and characterization of genes for a second anthranilate synthetase in *Pseudomonas aeruginosa*: interchangeability of the two anthranilate synthases and evolutionary implications. *J Bacteriol* 1990;**172**:884–900.
25. **Köhler T**, Kocjancic-Curty L, Barja F, *et al.* Swarming of *Pseudomonas aeruginosa* is dependent on cell-to-cell signaling and requires flagella and pili. *J Bacteriol* 2000;**182**:5990–6.
26. **Köhler T**, Buckling A, Van Delden C. Cooperation and virulence of clinical *Pseudomonas aeruginosa* populations. *Proc Natl Acad Sci U S A* 2009;**106**:6339–44.
27. **Pesci EC**, Pearson JP, Seed PC, *et al.* Regulation of *las* and *rhl* quorum sensing in *Pseudomonas aeruginosa*. *J Bacteriol* 1997;**179**:3127–32.
28. **Sandoz KM**, Mitzimberg SM, Schuster M. Social cheating in *Pseudomonas aeruginosa* quorum sensing. *Proc Natl Acad Sci U S A* 2007;**104**:15876–81.
29. **Zulianello L**, Canard C, Kohler T, *et al.* Rhamnolipids are virulence factors that promote early infiltration of primary human airway epithelia by *Pseudomonas aeruginosa*. *Infect Immun* 2006;**74**:3134–47.
30. **Davey ME**, Caiazza NC, O'Toole GA. Rhamnolipid surfactant production affects biofilm architecture in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* 2003;**185**:1027–36.
31. **Jensen PO**, Bjarnshold T, Phipps R, *et al.* Rapid necrotic killing of polymorphonuclear leukocytes is caused by quorum-sensing-controlled production of rhamnolipid by *Pseudomonas aeruginosa*. *Microbiology* 2007;**153**:1329–38.
32. **Shaver CM**, Hauser AR. Relative contributions of *Pseudomonas aeruginosa* ExoU, ExoS, and ExoT to virulence in the lung. *Infect Immun* 2004;**72**:6969–77.
33. **Bleves S**, Soscia C, Nogueira-Orlandi P, *et al.* Quorum sensing negatively controls type III secretion regulon expression in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* 2005;**187**:3898–902.
34. **Hentzer M**, Wu H, Andersen JB, *et al.* Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J* 2003;**22**:3803–15.
35. **Lesic B**, Lepine F, Deziel E, *et al.* Inhibitors of pathogen intercellular signals as selective anti-infective compounds. *PLoS Pathog* 2007;**3**:1229–39.

Lung alert

Clinical biomarkers in resectable NSCLC

In the last decade there have been small but real advances in the understanding and management of non-small cell lung cancer (NSCLC), with targeted treatments being a key innovation.

Hypoxia-induced upregulated tissue expression of carbonic anhydrase IX (CAIX) and elevated plasma CAIX levels are associated with more aggressive phenotypes in urological cancers. In this study, resected specimens of 555 patients with NSCLC were analysed by immunohistochemistry for CAIX and 209 preoperative plasma samples by ELISA for CAIX, with median follow-up of 35 months. 24.3% of the tissue specimens expressed high levels of CAIX and were associated with shorter overall survival in stage I and II, as was a plasma CAIX level >11 pg/ml. Tissue CAIX was underexpressed in adenocarcinoma subtypes.

This study shows that in resected early stage NSCLC, high tissue CAIX can serve as an independent predictor for shorter survival, as can plasma CAIX ELISA with 84% sensitivity and 95% specificity. Though targeted treatments directed specifically at CAIX are under development, this study also demonstrates a potential non-invasive clinical biomarker of early stage NSCLC, representing another of the technologies being developed involving proteome analysis of pretreatment peripheral blood in real time to help define the optimal therapeutic approach.

► **Ilie M**, Mazure NM, Hofman V, *et al.* High levels of carbonic anhydrase IX in tumour tissue and plasma are biomarkers of poor prognostic in patients with non-small cell lung cancer (NSCLC). *Br J Cancer* 2010;**102**:1627–35.

Burhan Khan

Darent Valley Hospital, Dartford, Kent, UK

Correspondence to Burhan Khan, Darent Valley Hospital, Dartford, Kent, UK; burhan.khan@nhs.net

Thorax 2010;**65**:710. doi:10.1136/thx.2010.145532