11. **Salome CM**, King GG, Berend N. Physiology of obesity and effects on lung function. *J Appl Physiol* 2010;**108**:206–11.
12. **Yaegashi M**, Jean R, Zuriqat M, *et al*. Outcome of morbid obesity in the intensive care unit. *J Intensive Care Med* 2005;**20**:147–54.
13. **Gajic O**, Frutos-Vivar F, Esteban A, *et al*. Ventilator settings as a risk factor for acute respiratory distress syndrome in mechanically ventilated patients. *Intensive Care Med* 2005;**31**:922–6.
14. **Gong MN**, Bajwa EK, Thompson BT, *et al*. Body mass index is associated with the development of acute respiratory distress syndrome. *Thorax* 2010;**65**:44–50.
15. **Mancuso P**. Obesity and lung inflammation. *J Appl Physiol* 2010;**108**:722–8.
16. **Neville A**, Brown C, Weng J, *et al*. Obesity is an independent risk factor of mortality in severely injured blunt trauma patients. *Arch Surg* 2004;**139**:983–7.
17. **Brown C**, Neville A, Rhee P, *et al*. The impact of obesity on the outcomes of 1,153 critically injured blunt trauma patients. *J Trauma* 2005;**59**:1048–51.
18. **Ciesla D**, Moore E, Johnson J, *et al*. Obesity increases risk for organ failure after severe trauma. *J Am Coll Surg* 2006;**203**:539–45.
19. **Slynkova K**, Mannino DM, Martin GS, *et al*. The role of body mass index and diabetes in the development of acute organ failure and subsequent mortality in an observational cohort. *Crit Care* 2006;**10**:R137.
20. **Andrews WR**. Obesity and intensive care unit survival: pass the potatoes? *Crit Care Med* 2008;**36**:369–70.

Lung alert

Vitamin B deficiency may increase risk of lung cancer

It has been hypothesised that deficiencies in B vitamin levels may influence gene expression by means of aberrant methylation patterns as vitamin B is essential for DNA synthesis and methylation. This study investigated the role of B vitamins (B2, B6, B9, B12) and methionine status on the development of lung cancer. It was conducted over 8 years and included participants from the EPIC (European Prospective Investigation into Cancer and Nutrition) cohort.

899 cases were suitable for serum analysis. Using sex, smoking and age-specific incidence rates, cumulative risks of lung cancer were greatest for current smokers followed by former smokers and never smokers. After adjusting for matching variables and cotinine (a marker of smoking), a substantially lower risk of lung cancer was seen for increasing levels of B6 and methionine. The study also revealed a high deficiency in nutrient levels of B vitamins in many western populations. Smokers were found to consume fewer fruits and vegetables (B vitamins source). Smoking intensity was inversely associated with folate, B12 and B6 levels. Serum levels of B vitamins and metabolites were ascertained to be partially determined by diet and clearly affected by vitamin supplements so low vitamin levels were found to be modifiable.

This study suggests that smoking and dietary intake can have an impact on lung cancer development. Although previous studies of vitamin B supplementation in other cancers have not been shown to be of benefit, this is clearly an interesting area for further development.

► **Johansson M**, Relton C, Magne Ueland P. Serum B vitamin levels and risk of lung cancer. *JAMA* 2010;**303**:2377–85.

Omar Hashmi

RMO, HDU, Karachi, Pakistan

Correspondence to Dr Omar Hashmi, RMO, HDU, 46-D Block 4 Federal B Area, Karachi 78400, Pakistan; drowh@live.com

Published Online First 29 September 2010

Thorax 2011;**66**:73. doi:10.1136/thx.2010.150235