Funding SBD is supported by the National Institute for Health Research Biomedical Research Centre at Guy's and St Thomas' NHS Foundation Trust/King's College London. TW was, and MA is, supported by Abbott Laboratories.

Competing interests Abbott Laboratories who supported TW and presently support MA market palivizumab, a monoclonal antibody against RSV.

Ethics approval This study was conducted with the approval of the Research Ethics Committee of King's College Hospital NHS Trust.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **Stokes GM**, Milner AD, Hodges IG, *et al*. Lung function abnormalities after acute bronchiolitis. *J Pediatr* 1981;**98**:871–4.
2. **Hall CB**, Hall WJ, Gala CL, *et al*. Long-term prospective study in children after respiratory syncytial virus infection. *J Pediatr* 1984;**105**:358–64.
3. **Noble V**, Murray M, Webb MS, *et al*. Respiratory status and allergy nine to 10 years after acute bronchiolitis. *Arch Dis Child* 1997;**76**:315–19.
4. **Korppi M**, Piippo-Savolainen E, Korhonen K, *et al*. Respiratory morbidity 20 years after RSV infection in infancy. *Pediatr Pulmonol* 2004;**38**:155–60.
5. **Broughton S**, Sylvester KP, Fox G, *et al*. Lung function in prematurely born infants after viral lower respiratory tract infections. *Pediatr Infect Dis J* 2007;**26**:1019–24.
6. **Young S**, O'Keeffe PT, Arnott J, *et al*. Lung function, airway responsiveness, and respiratory symptoms before and after bronchiolitis. *Arch Dis Child* 1995;**72**:16–24.
7. **Martinez FD**, Morgan WJ, Wright AL, *et al*. Diminished lung function as a predisposing factor for wheezing respiratory illness in infants. *N Engl J Med* 1988;**319**:1112–17.
8. **Martinez FD**, Morgan WJ, Wright AL, *et al*. Initial airway function is a risk factor for recurrent wheezing respiratory illnesses during the first three years of life. Group Health Medical Associates. *Am Rev Respir Dis* 1991;**143**:312–16.
9. **Broughton S**, Bhat R, Roberts A, *et al*. Diminished lung function, RSV infection, and respiratory morbidity in prematurely born infants. *Arch Dis Child* 2006;**91**:26–30.
10. **Clark SJ**, Beresford MW, Subhedar NV, *et al*. Respiratory syncytial virus infection in high risk infants and the potential impact of prophylaxis in a United Kingdom cohort. *Arch Dis Child* 2000;**83**:313–16.
11. **Hey EN**. The relation between environmental temperature and oxygen consumption in the new-born baby. *J Physiol* 1969;**200**:589–603.
12. **Martinez FD**, Wright AL, Taussig LM, *et al*. Asthma and wheezing in the first six years of life. *N Engl J Med* 1995;**332**:133–8.
13. **Hodgson J**, Zuckerman M, Smith M. Development of a novel internal control for a real-time PCR for HSV DNA types 1 and 2. *J Clin Virol* 2007;**38**:217–20.
14. **Schrag SJ**, Zell ER, Lynfield R, *et al*; The Active Bacterial Core Surveillance Team. A population-based comparison of strategies to prevent early-onset group B streptococcal disease in neonates. *N Engl J Med* 2002;**347**:233–9.
15. **Fowle PW**, Schmidt B. Diagnostic tests for bacterial infection from birth to 90 days—a systematic review. *Arch Dis Child Fetal Neonatal Ed* 1998;**78**:F92–8.
16. **Kuypers J**, Wright N, Ferrenberg J, *et al*. Comparison of real-time PCR assays with fluorescent-antibody assays for diagnosis of respiratory virus infections in children. *J Clin Microbiol* 2006;**44**:2382–8.
17. **van de Pol AC**, van Loon AM, Wolfs TF, *et al*. Increased detection of respiratory syncytial virus, influenza viruses, parainfluenza viruses, and adenoviruses with real-time PCR in samples from patients with respiratory symptoms. *J Clin Microbiol* 2007;**45**:2260–2.
18. **Bonroy C**, Vankeerberghen A, Boel A, *et al*. Use of a multiplex real-time PCR to study the incidence of human metapneumovirus and human respiratory syncytial virus infections during two winter seasons in a Belgian paediatric hospital. *Clin Microbiol Infect* 2007;**13**:504–9.

Journal club

A role for cigarette fumes in the development of emphysema through reduced alveolar cell proliferation and upregulation of apoptosis

This paper describes a molecular mechanism for the loss of alveolar tissue in cigarette smoke-induced lung injury. Under the influence of adverse environmental conditions including hypoxia, the intracellular stress protein Rtp801 is transcribed. This study sought to demonstrate that Rtp801 downregulates the kinase mammalian target of rapamycin (mTOR) which is responsible for alveolar cell proliferation and upregulates the apoptotic and inflammatory pathways through the activation of the transcription factor nuclear factor kappa B (NF- κ B).

Increased levels of Rtp801 were measured in alveolar cells of patients with chronic obstructive pulmonary disease compared with healthy individuals through histological staining and electrophoresis. Western blotting studies in mice showed significantly increased Rtp801 levels following exposure to cigarette smoke for 7 days. Mice treated with the antioxidant N-acetyl-L-cysteine followed by exposure to smoke showed no overall increase in Rtp801 levels. Overexpression of Rtp801 in mice without exposure to smoke led to significant alveolar apoptosis. Rtp801 knock-out mice were protected from alveolar inflammatory and apoptotic changes in the presence of smoke. Inhibition of NF- κ B activity resulted in decreased levels of inflammation and apoptosis.

These results confirm the role of this cellular stress response in alveolar damage and provide further evidence for a genetic component to emphysema.

► **Yoshida T**, Mett I, Bhunia AK, *et al*. Rtp801, a suppressor of mTOR signalling, is an essential mediator of cigarette smoke-induced pulmonary injury and emphysema. *Nat Med* 2010;**16**:767–73.

M East

Correspondence to M East, AcFY2, Department of Microbiology, Leeds Teaching Hospitals NHS Trust, Old Medical School, Leeds General Infirmary, Great George Street, West Yorkshire, UK; m.east@doctors.net.uk

Published Online First 27 September 2010

Thorax 2011;**66**:473. doi:10.1136/thx.2010.149898