Provenance and peer review Not commissioned; not externally peer reviewed.

Accepted 5 August 2010
Published Online First 14 October 2010

Thorax 2011;**66**:543–544.
doi:10.1136/thx.2010.148510

REFERENCES

1. **National Clinical Guideline Centre.** *Chronic obstructive pulmonary disease: management of adults with chronic obstructive pulmonary disease in primary and secondary care.* Updated. London: National Clinical Guideline Centre, 2010.
2. **Department of Health/Respiratory Team.** *Consultation on a strategy for services for chronic obstructive pulmonary disease (COPD) in England.* London, Department of Health, 2010.
3. **National Collaborating Centre for Chronic Conditions.** *Chronic obstructive pulmonary disease. National clinical guideline on management of chronic obstructive pulmonary disease in adults in primary and secondary care.* *Thorax* 2004;**59** (Suppl 1):1–232.
4. **Puhan MA, Garcia-Aymerich J, Frey M, et al.** Expansion of the prognostic assessment of patients with chronic obstructive pulmonary disease: the updated BODE index and the ADO index. *Lancet* 2009;**374**:704–11.
5. **Jones RC, Donaldson GC, Chavannes NH, et al.** Derivation and validation of a composite index of severity in chronic obstructive pulmonary disease: the DOSE index. *Am J Respir Crit Care Med* 2009;**180**:1189–95.

Reversed halo sign in pulmonary zygomycosis

We read with interest the article ‘The atoll sign’ by Walsh and Robertson¹ in the November issue of *Thorax*. The authors report a case of cryptogenic organising pneumonia with the atoll sign, also called the reversed halo sign. As mentioned by the authors, this CT sign was first described in

cryptogenic organising pneumonia and was initially considered to be specific for this disease.² It was subsequently reported in a variety of pulmonary diseases, including paracoccidioidomycosis (South American blastomycosis), tuberculosis, pulmonary sarcoidosis, lymphomatoid granulomatosis, Wegener’s granulomatosis, lipoid pneumonia and pneumococcal pneumonia.^{3–5}

We would like to highlight another important cause of the reversed halo sign: invasive pulmonary fungal infections, particularly pulmonary zygomycosis (PZ) (figure 1). In immunosuppressed patients, the presence of the reversed halo sign on CT should be considered as invasive fungal disease until proven otherwise. It is an early sign that is more frequently seen in patients with PZ than invasive pulmonary aspergillosis (IPA).⁶

Early institution of high-dose antifungal therapy is associated with improved outcomes; therefore, early recognition of invasive fungal disease is important. Moreover, because the therapy for presumed fungal pneumonia in this population is often aimed at IPA due to its higher incidence and the preferred antifungal agent for IPA is voriconazole, which is not effective against PZ, it is important to differentiate between the two entities. The presence of the reversed halo sign can be used to optimise antifungal therapy to cover PZ.

Myrna C B Godoy, Edith M Marom

University of Texas M.D. Anderson Cancer Center, Department of Diagnostic Radiology, Houston, Texas, USA

Correspondence to Myrna C B Godoy, University of Texas M.D. Anderson Cancer Center, Department of Diagnostic Radiology, 1515 Holcombe Blvd., Unit 371, Houston, TX 77030, USA; mgodoy@mdanderson.org

Competing interests None.

Provenance and peer review Not commissioned; not externally peer reviewed.

Accepted 26 January 2011
Published Online First 8 March 2011

Thorax 2011;**66**:544. doi:10.1136/thx.2010.157776

REFERENCES

1. **Walsh SL, Robertson BJ.** Images in thorax. The atoll sign. *Thorax* 2010;**65**:1029–30.
2. **Voloudaki AE, Froudarakis ME, Datsis GE, et al.** Crescentic and ring-shaped opacities. CT features in two cases of bronchiolitis obliterans organizing pneumonia (BOOP). *Acta Radiol* 1996;**37**:889–92.
3. **Gasparetto EL, Escuissato DL, Davaus T, et al.** Reversed halo sign in pulmonary paracoccidioidomycosis. *AJR Am J Roentgenol* 2005;**184**:1932–4.
4. **Marchiori E, Grando RD, Simoes Dos Santos CE, et al.** Pulmonary tuberculosis associated with the reversed halo sign on high-resolution CT. *Br J Radiol* 2010;**83**:e58–60.
5. **Tzilas V, Provata A, Koti A, et al.** The “reversed halo” sign in pneumococcal pneumonia: a review with a case report. *Eur Rev Med Pharmacol Sci* 2010;**14**:481–6.
6. **Wahba H, Truong MT, Lei X, et al.** Reversed halo sign in invasive pulmonary fungal infections. *Clin Infect Dis* 2008;**46**:1733–7.

Serum 25-hydroxy vitamin D and exercise capacity in COPD

Janssens and colleagues¹ have recently reported that vitamin D deficiency is very common in patients affected by chronic obstructive pulmonary disease (COPD) and that vitamin D status correlates with lung function. In the same issue of *Thorax*, Quint and Wedzicha² discuss potential effects of vitamin D deficiency and supplementation

Figure 1 Pulmonary zygomycosis in a 24-year-old woman undergoing chemotherapy for recurrence of acute myelogenous leukaemia, 1 year following allogeneic stem cell transplantation, with a 2-week history of fever and dry cough. (A) Chest CT shows a focal round area of ground-glass attenuation surrounded by a ring of consolidation in the right lower lobe, consistent with the reversed halo sign. (B) Photomicrograph of the specimen from a transthoracic biopsy of the lesion shows pauciseptated hyphae with non-parallel walls and 90° branching, characteristic of *Zygomycetes* species, which is confirmed by culture. (C) One-month follow-up CT shows cavitation of the lesion, which eventually resolved under antifungal therapy.