3. **Brochard L**, Mancebo J, Wysocki M, *et al*. Noninvasive ventilation for acute exacerbations of chronic obstructive pulmonary disease. *N Engl J Med* 1995;**28**:817–22.
4. **Man WD-C**, Polkey MI, Donaldson N, *et al*. Community pulmonary rehabilitation after hospitalisation for acute exacerbations of chronic obstructive pulmonary disease: randomised controlled study. *BMJ* 2004;**329**:1209.
5. **Schönhofer B**. Non-invasive positive pressure ventilation in patients with stable hypercapnic COPD: light at the end of the tunnel? *Thorax* 2010;**65**:765–7.

CORRESPONDENCE

Comment on: Morbidity and mortality associated with the restrictive spirometric pattern: a longitudinal study

We read with interest the paper by Guerra *et al*¹ profiling the demographic/clinical characteristics and prospectively assessing the prognosis of subjects with a restrictive spirometric pattern enrolled in the TESAOD population-based study. The manuscript has the merit of following up a large number of patients for 14 years and investigating how selected co-morbidities are relevant to survival.²

The results of this study deserve comparison with those published in 2008 by our research group on 1265 subjects aged 65–97 years.³ First, in an older population (mean age 73.4 years) we found a comparable prevalence of restriction at spirometry (12% vs 10.9%), although restriction is an age-related phenomenon and, thus, is expected to be much lower in the younger population (ie, 6.6% in a population aged 42.2 years⁴). Secondly, the study by Guerra and colleagues strongly reproduces the increased mortality risk rates (MRRs) observed in our work: for all (HR 1.89; 95% CI 1.15 to 3.11 (Scarlata *et al*) vs HR 1.7; 95% CI 1.3 to 2.3 (Guerra *et al*)), cardiac (MRR 1.51; 95% CI 0.60 to 3.78 (Scarlata *et al*) vs HR 2.0; 95% CI 1.3 to 3.1 (Guerra *et al*)) and cerebrovascular (MRR 4.79; 95% CI 1.54 to 14.84 (Scarlata *et al*) vs HR 2.4; 95% CI 0.9 to 6.3 (Guerra *et al*)). This finding is consistent with restrictive lung dysfunction affecting survival in a predictable manner.

At variance with our study, that of Guerra and colleagues lacks information about clinical correlates of restrictive pulmonary disease. The Cox proportional hazard models are adjusted only for sex, age and body mass index, but not for concomitant conditions known to be associated with restriction. Indeed, we found that co-morbidities such as kyphosis of the spine (OR 2.40; 95% CI 1.58 to 3.64) and diabetes mellitus (OR 1.66; 95% CI 1.00 to 2.74) as well as the physical (Activities of Daily Living scale, OR 2.17; 95% CI 1.32 to 3.58; 6 minute walking test, OR 1.75; 95% CI 1.15 to 2.67) and cognitive (Mini Mental State Examination, OR 2.05; 95% CI 1.27 to 3.32) status are strong independent correlates of restriction.

In conclusion, the paper by Guerra and colleagues has the merit of confirming that the restrictive spirometric pattern is highly prevalent and is associated with a clinical profile and risk factors differing from those of obstructive lung disease. However, research is needed to expand our knowledge of the mechanisms underlying restriction as well as to explain the link between restriction and frailty. Clarifying these issues will allow the implementation of both guided screening and preventive interventions.

Simone Scarlata,¹ Claudio Pedone,¹ Filippo L Fimognari,^{1,2} Vincenzo Bellia,³ Francesco Forastiere,⁴ Raffaele Antonelli Incalzi^{1,5}

¹Unit of Respiratory Pathophysiology, Chair of Geriatrics, University Campus Bio-Medico, Rome, Italy; ²Unit of Respiratory Physiopathology, Internal Medicine Division, Ospedale 'L. Parodi Delfino', Colleferro, Rome, Italy;

³Institute of General Medicine and Pneumology, Università di Palermo, Palermo, Italy; ⁴Department of Epidemiology, Roma E Health Authority, Rome, Italy;

⁵San Raffaele Foundation, Cittadella della Carità, Taranto, Italy

Correspondence to Simone Scarlata, Servizio di Fisiopatologia Respiratoria, Università Campus Bio-Medico, Via A. del Portillo, 200, Rome, Italy; s.scarlata@unicampus.it

Competing interests None.

Provenance and peer review Not commissioned; not externally peer reviewed.

Accepted 30 August 2010

Published Online First 22 October 2010

Thorax 2011;**66**:826. doi:10.1136/thx.2010.149419

REFERENCES

1. **Guerra S**, Sherrill DL, Venker C, *et al*. Morbidity and mortality associated with the restrictive spirometric pattern: a longitudinal study. *Thorax* 2010;**65**:499–504.
2. **Mannino DM**, Buist AS, Petty TL, *et al*. Lung function and mortality in the United States: data from the First National Health and Nutrition Examination Survey follow up study. *Thorax* 2003;**58**:388–93.
3. **Scarlata S**, Pedone C, Fimognari FL, *et al*. Restrictive pulmonary dysfunction at spirometry and mortality in the elderly. *Respir Med* 2008;**102**:1349–54.
4. **Mannino DM**, Ford ES, Redd SC. Obstructive and restrictive lung disease and functional limitation: data from the Third National Health and Nutrition Examination. *J Intern Med* 2003;**254**:540–7.

Authors' response

We appreciate the interest shown in our study by Scarlata and colleagues.¹ We agree with them that, in line with several previous reports,^{2–3} the prevalence of the restrictive spirometric pattern was strikingly similar in the TESAOD⁴ and SARA⁵ studies, despite the substantial difference in age distribution between the two populations (the TESAOD report included subjects ≥21 years and the SARA study subjects ≥65 years of age).

However, a direct comparison of cause-specific mortality between the two studies should be interpreted with caution because of the different assessment of the restrictive

spirometric pattern, which was evaluated at a single point in time in the SARA study and prospectively in the TESAOD study. Not all subjects with the restrictive spirometric pattern at baseline will have a consistent restrictive spirometric pattern (the one that was used for comparison in the letter by Scarlata *et al*) over time. Actually, in TESAOD only one out of three such subjects did. The remaining two-thirds either had an inconsistent restrictive longitudinal pattern or developed airflow limitation at some point during the follow-up. Profiles of cause-specific mortality risk differed notably across these three longitudinal groups. For example, hazard ratios for mortality by cardiac disease were 2.0, 2.7 and 1.6, respectively.

We believe that the most novel contribution of our study does not lie in confirming the mortality risk associated with the cross-sectional restrictive spirometric pattern, but rather in assessing spirometric patterns prospectively, for two main reasons. First, our data indicate that up to 38% of subjects with a restrictive spirometric pattern at enrolment developed airflow limitation during the study follow-up. These subjects were more likely to be smokers, to have a physician-confirmed diagnosis of asthma at enrolment, and—unlike those with recurrent or inconsistent restrictive patterns—to die of COPD during follow-up. These results suggest that an underlying airway obstruction may be present in a significant proportion of cases with spirometric restriction assessed at a single time point, and this may explain the finding (apparently conflicting with ours) of an increased pulmonary mortality risk associated with spirometric restriction in the SARA study. Second, the prospective analyses of our study demonstrate that, among subjects who do not develop an obstructive pattern over time, both the recurrent and the inconsistent spirometric restriction increase all-cause mortality risk by a substantial magnitude.

Although what causes increased mortality in these groups remains to be determined, our findings do suggest that this pulmonary condition predisposes to (or at least is linked to) other extrapulmonary conditions such as cardiovascular disease and diabetes. Thus, these comorbidities may be in the causal pathway from spirometric restriction to mortality and we therefore elected not to include them among covariates in our Cox proportional hazards models. We definitely agree with Scarlata and colleagues that further research is required to understand the factors that are related aetiologically to spirometric restriction, the molecular mechanisms that drive its effects on all-cause and cause-specific mortality, and the possible implications of these findings for early identification of subjects at risk.

Stefano Guerra,^{1,2} Fernando D Martinez¹

¹Arizona Respiratory Center, University of Arizona, Tucson, Arizona, USA; ²CREAL, Barcelona, Spain

Correspondence to Fernando D Martinez, Arizona Respiratory Center, University of Arizona, 1501 N Campbell, Tucson, AR 85724, USA; fernando@arc.arizona.edu

Competing interests None.

Provenance and peer review Not commissioned; not externally peer reviewed.

Accepted 30 August 2010

Published Online First 22 October 2010

Thorax 2011;**66**:826–827.

doi:10.1136/thx.2010.150227

REFERENCES

1. **Scarlata S**, Pedone C, Fimognari FL, *et al.* Comment on: Morbidity and mortality associated with the restrictive spirometric pattern: a longitudinal study. *Thorax* 2011;**66**:826.
2. **Mannino DM**, Buist AS, Petty TL, *et al.* Lung function and mortality in the United States: data from the First National Health and Nutrition Examination Survey follow-up study. *Thorax* 2003;**58**:388–93.
3. **Mannino DM**, Davis KJ. Lung function decline and outcomes in an elderly population. *Thorax* 2006;**61**:472–7.
4. **Guerra S**, Sherrill DL, Venker C, *et al.* Morbidity and mortality associated with the restrictive spirometric pattern: a longitudinal study. *Thorax* 2010;**65**:499–504.
5. **Scarlata S**, Pedone C, Fimognari FL, *et al.* Restrictive pulmonary dysfunction at spirometry and mortality in the elderly. *Respir Med* 2008;**102**:1349–54.

Validity of using Hospital Episode Statistics data on monitoring disease trends

We read with interest the article by Koshy *et al.*¹ The findings are important in documenting changes in admission rates of childhood pneumonia and empyema since the introduction of heptavalent pneumococcal conjugate vaccine (PCV7). We are concerned that undue emphasis has been placed on Hospital Episode Statistics (HES) data to define the aetiology of childhood pneumonia, particularly ‘bacterial pneumonia’.²

Given the magnitude of the case numbers reported, it would appear that the analyses are based on all pneumonia codes collectively. This would also (although it is not clear from the article) include ‘unspecified pneumonia’, which describes pneumonia of any aetiology. Our analysis of national HES data on childhood pneumonia (1997–2006) showed that 91% of cases were coded as unspecified pneumonia. This may be of significance given that much unspecified pneumonia in children is likely to be viral; in routine clinical practice it can be difficult to differentiate between viral and bacterial pneumonia.³

The authors¹ also assert that ‘PCV7 offers protection against the most common serotypes accounting for most of the bacterial pneumonias in children’. The references provided do not support this statement. There are international variations in serotype distributions of laboratory-confirmed pneumococcal disease.⁴ There are no published

data on the serotype distribution of pneumococcal pneumonia for UK children.

We have evaluated the accuracy of HES data for paediatric pneumonia in the North East of England. The incidence was previously established in a prospective study,⁵ and we repeated it prospectively between 2008 and 2009. Of 50 subjects identified during prospective recruitment, 14 (28%) had misattributed codes and were not identified in the coding list. These patients were coded, for example, as unspecified acute upper respiratory tract infection (J06.9), dyspnoea (R06.0) and cough (R05), despite a clinical diagnosis of pneumonia. Among those identified by HES codes, pneumonia (N=5) and lower respiratory tract infection (N=2) were coded as secondary diagnoses. These figures suggest that reliance on primary diagnostic codes on the basis of HES data could underestimate the levels of pneumonia. There are no reasons to think that levels of miscoding have changed over time.

This article does not describe trends in bacterial pneumonia as stated throughout the paper but all causes of pneumonia. We suggest that use of HES data should be limited to analysis of changes in the overall incidence of pneumonia.

Mohamed A Elemraid,^{1,2} Kerry Pollard,¹ Matthew F Thomas,³ Andrew R Gennery,^{1,2} Katherine M Eastham,⁴ Stephen P Rushton,³ Fiona Hampton,⁵ Pauline Singleton,⁵ Russell Gorton,⁶ David A Spencer,⁷ Julia E Clark,^{1,2} on behalf of North East of England Paediatric Respiratory Infection Study Group

¹Department of Paediatric Infectious Disease and Immunology, Newcastle upon Tyne Hospitals NHS Trust, Newcastle upon Tyne, UK; ²Institute of Cellular Medicine, Newcastle University, Newcastle upon Tyne, UK; ³School of Biology, Newcastle University, Newcastle upon Tyne, UK; ⁴Department of Paediatrics, Sunderland Royal Hospital, Sunderland, UK; ⁵Department of Paediatrics, James Cook University Hospital, Middlesbrough, UK; ⁶Regional Epidemiology Unit, Health Protection Agency North East, Newcastle upon Tyne, UK; ⁷Department of Respiratory Paediatrics, Newcastle upon Tyne Hospitals NHS Trust, Newcastle upon Tyne, UK

Correspondence to Dr Julia Clark, Great North Children's Hospital, Queen Victoria Road, Newcastle upon Tyne, NE1 4LP, UK; julia.clark@nuth.nhs.uk

Competing interests None.

Provenance and peer review Not commissioned; internally peer reviewed.

Accepted 20 October 2010

Published Online First 2 December 2010

Thorax 2011;**66**:827. doi:10.1136/thx.2010.153551

REFERENCES

1. **Koshy E**, Murray J, Bottle A, *et al.* Impact of the seven-valent pneumococcal conjugate vaccination (PCV7) programme on childhood hospital admissions for bacterial pneumonia and empyema in England: national time-trends study, 1997–2008. *Thorax* 2010;**65**:770–4.
2. **Lorence DP**, Ibrahim IA. Disparity in coding concordance: do physicians and coders agree? *J Health Care Finance* 2003;**29**:43–53.

3. **British Thoracic Society**. Guidelines for the management of community acquired pneumonia in childhood. *Thorax* 2002;**57**(Suppl 1):i1–24.
4. **Hausdorff WP**, Bryant J, Paradiso PR, *et al.* Which pneumococcal serogroups cause the most invasive disease: implications for conjugate vaccine formulation and use, Part I. *Clin Infect Dis* 2000;**30**:100–21.
5. **Clark JE**, Hammal D, Hampton F, *et al.* Epidemiology of community-acquired pneumonia in children seen in hospital. *Epidemiol Infect* 2007;**135**:262–9.

Authors' response

Elemraid and colleagues raise important points about potential diagnostic misclassification and under-ascertainment using the Hospital Episodes Statistics (HES) database, as well as the absence of national pneumococcal serotype data for children.

The HES database covers all NHS hospital activity in England and has been widely used to report disease trends. It also provides the opportunity to estimate the clinical impact of major clinical policies on disease burden but, as with any large epidemiological dataset, has inherent weaknesses at the individual level.

Our study aimed to focus on common community-acquired bacterial pneumonia trends to evaluate the impact of the heptavalent pneumococcal conjugate vaccine (PCV7). We agree that pneumonia is a clinical diagnosis and that it is difficult to differentiate between bacterial and viral causes. Furthermore, a significant proportion of cases (8–40%) of community-acquired pneumonia can be of mixed aetiology.¹ Hence, we applied broad pneumonia definitions. We aimed to focus on bacterial pneumonias and so excluded specific viral pneumonia ICD-10 codes (eg, ‘viral pneumonia, not elsewhere classified’—all J12 codes). The codes we searched are listed in the Appendix.

The authors highlight a useful point that some children diagnosed with pneumonia may have symptoms and/or signs recorded in the primary diagnosis field. Hence, we acknowledge under-ascertainment is possible for some pneumonia admissions. HES coding is dependent on the recording of the ‘reason for admission’ by clinicians and the subsequent coding by the trained staff, and we included this as a potential limitation in our discussion. We agree that such levels of miscoding are unlikely to have significantly changed over time. Therefore, this would suggest that the pneumonia admission trends that we observed are likely to represent real changes.

We used the Health Protection Agency cumulative weekly incidence reports of PCV7 and non-PCV7 isolates for children under 5 years,² together with the national serotype surveillance for all ages,³ as the best available source of information on pneumococcal serotypes causing invasive pneumococcal disease. Admittedly, this covers a broader spectrum of invasive diseases. In