REFERENCES

- Havelock T, Teoh R, Laws D, *et al*. Pleural procedures and thoracic ultrasound: British Thoracic Society pleural disease guideline 2010. *Thorax* 2010;**65** (Suppl 2):ii61–76.
- Kirkpatrick AW, Sirois M, Laupland KB, *et al*. Hand-held thoracic sonography for detecting post-traumatic pneumothoraces: the Extended Focused Assessment with Sonography for Trauma (EFAST). *J Trauma* 2004;**57**:288–95.
- Soldati G, Testa A, Sher S, *et al*. Occult traumatic pneumothorax: diagnostic accuracy of lung ultrasonography in the emergency department. *Chest* 2008;**133**:204–11.
- Soldati G, Testa A, Pignataro G, *et al*. The ultrasonographic deep sulcus sign in traumatic pneumothorax. *Ultrasound Med Biol* 2006;**32**:1157–63.
- Elbarbary M, Melniker L, Volpicelli G. Development of evidence based clinical recommendations and consensus statements in the critical ultrasound field: why and how? *Crit Ultrasound J* 2010;**2**(3):93–5.
- Ball CG, Hameed SM, Evans D, *et al*. Occult pneumothorax in the mechanically ventilated trauma patient. *Can J Surg* 2003;**46**:373–9.
- Ball CG, Kirkpatrick AW, Feliciano DV. The occult pneumothorax: what have we learned? *Can J Surg* 2009;**52**:E173–9.
- Wilkerson RG, Stone MB. Sensitivity of bedside ultrasound and supine anteroposterior chest radiographs for the identification of pneumothorax after blunt trauma. *Acad Emerg Med* 2010;**17**:11–17.
- Alsalam W, Lewis D. Towards evidence based emergency medicine: best BETs from the Manchester Royal Infirmary. BET 1: Is ultrasound or chest x ray best for the diagnosis of pneumothorax in the emergency department? *Emerg Med J* 2009;**26**:434–5.
- Elbarbary M, Melniker L, Volpicelli G. Development of evidence based clinical recommendations and consensus statements in the critical ultrasound field: why and how? *Crit Ultrasound J* 2010;**2**(3):93–5.
- Zhang M, Liu ZH, Yang JX, *et al*. Rapid detection of pneumothorax by ultrasonography in patients with multiple trauma. *Crit Care* 2006;**10**:R112.
- Blaivas M, Lyon M, Duggal S. A prospective comparison of supine chest radiography and bedside ultrasound for the diagnosis of traumatic pneumothorax. *Acad Emerg Med* 2005;**12**:844–9.
- Kirkpatrick AW, Sirois M, Laupland KB, *et al*. Hand-held thoracic sonography for detecting post-traumatic pneumothoraces: the Extended Focused Assessment with Sonography for Trauma (EFAST). *J Trauma* 2004;**57**:288–95.
- Knudtson JL, Dort JM, Helmer SD, *et al*. Surgeon-performed ultrasound for pneumothorax in the trauma suite. *J Trauma* 2004;**56**:527–30.
- Dulchavsky SA, Schwarz KL, Kirkpatrick AW, *et al*. Prospective evaluation of thoracic ultrasound in the detection of pneumothorax. *J Trauma* 2001;**50**:201–5.
- Ball CG, Kirkpatrick AW, Laupland KB, *et al*. Incidence, risk factors, and outcomes for occult pneumothoraces in victims of major trauma. *J Trauma* 2005;**59**:917–24; discussion 24–5.
- Rowan KR, Kirkpatrick AW, Liu D, *et al*. Traumatic pneumothorax detection with thoracic US: correlation with chest radiography and CT—initial experience. *Radiology* 2002;**225**:210–14.
- Soldati G, Testa A, Pignataro G, *et al*. The ultrasonographic deep sulcus sign in traumatic pneumothorax. *Ultrasound Med Biol* 2006;**32**:1157–63.
- Soldati G, Testa A, Sher S, *et al*. Occult traumatic pneumothorax: diagnostic accuracy of lung ultrasonography in the emergency department. *Chest* 2008;**133**:204–11.
- Reissig A, Kroegel C. Accuracy of transthoracic sonography in excluding post-interventional pneumothorax and hydropneumothorax. Comparison to chest radiography. *Eur J Radiol* 2005;**53**:463–70.
- Goodman TR, Traill ZC, Phillips AJ, *et al*. Ultrasound detection of pneumothorax. *Clin Radiol* 1999;**54**:736–9.
- Chung MJ, Goo JM, Im JG, *et al*. Value of high-resolution ultrasound in detecting a pneumothorax. *Eur Radiol* 2005;**15**:930–5.
- Sartori S, Tombesi P, Trevisani L, *et al*. Accuracy of transthoracic sonography in detection of pneumothorax after sonographically guided lung biopsy: prospective comparison with chest radiography. *AJR Am J Roentgenol* 2007;**188**:37–41.
- Slater A, Goodwin M, Anderson KE, *et al*. COPD can mimic the appearance of pneumothorax on thoracic ultrasound. *Chest* 2006;**129**:545–50.
- Tocino IM, Miller MH, Fairfax WR. Distribution of pneumothorax in the supine and semirecumbent critically ill adult. *AJR Am J Roentgenol* 1985;**144**:901–5.

as shown in the prospective study by Goodman *et al*¹⁸ using CT as the gold standard, small pneumothoraces may fail to be detected. The CXR is undoubtedly unreliable in the detection of small pneumothoraces in the supine patient²² and in specific clinical circumstances, as suggested by Agricola *et al*, ultrasound may be of value. This being the case, we agree that if a suitably skilled operator and ultrasound equipment are available at the patient's bedside then ultrasound may provide useful diagnostic information, but we maintain that it is unlikely to obviate the need for a formal CXR.

We are surprised that the authors experienced significant delays in obtaining 'stat' portable CXRs in their critical care and emergency departments and that patients may have died as a consequence. Clinicians managing critically ill trauma patients require rapid access to portable CXRs and all institutions managing such patients should be able to deliver this.

In conclusion, we agree that in supine and trauma patients ultrasound may be a valuable tool in the detection of pneumothorax. In these patients, ultrasound may have increased sensitivity compared with a CXR, although difficulty with pneumothorax quantification suggests that ultrasound is unlikely to completely replace the need for a radiograph. In the majority of cases of spontaneous or postprocedure pneumothorax, ultrasound is unlikely to provide additional benefit over the combination of CXR and clinical judgement when deciding management.

Tom Havelock,¹ Richard Teoh,² Diane Laws,³ Nick Maskell,⁴ Fergus Gleeson⁵

¹Wellcome Trust Clinical Research Facility, Southampton General Hospital, Southampton, UK; ²Department of Respiratory Medicine, Castle Hill Hospital, Cottingham, East Yorkshire, UK; ³Department of Thoracic Medicine, Royal Bournemouth Hospital, Bournemouth, UK; ⁴Department of Clinical Sciences, Southmead Hospital, University of Bristol, Bristol, UK; ⁵Department of Radiology, Churchill Hospital, Oxford, UK

Correspondence to Dr Tom Havelock, Wellcome Trust Clinical Research Facility, Southampton General Hospital, Tremona Road, Southampton. SO16 6YD, UK; t.havelock@soton.ac.uk

Competing interests None.

Provenance and peer review Not commissioned; not externally peer reviewed.

Accepted 30 November 2010

Published Online First 30 December 2010

Thorax 2011;**66**:829. doi:10.1136/thx.2010.156398

REFERENCES

- Agricola E, Arbelot C, Blaivas M, *et al*. Ultrasound performs better than radiographs. *Thorax* 2011;**66**:828–9.
- Havelock T, Teoh R, Laws D, *et al*. Pleural procedures and thoracic ultrasound: British Thoracic Society Pleural Disease Guideline 2010. *Thorax* 2010;**65**(Suppl 2):ii61–76.

Authors' response

We thank Agricola and colleagues¹ for their compliments on our guideline² and their contribution to the discussion on the role of ultrasound in the detection of pneumothorax, but we maintain that the medical community should proceed with caution when using ultrasound in the detection and management of pneumothoraces. If the reviews^{3–7} referenced are not considered (5 papers), 13 of the remaining 24 papers referenced are in two well-defined patient groups—trauma^{8–16} and post-intervention.^{17–20} None of the papers published prospectively demonstrated improved outcomes and management change using ultrasound in comparison with chest x-ray (CXR), and perhaps more significantly only one prospective blinded study in medical patients with varying degrees of respiratory compromise has been reported and this demonstrated an unacceptably high false positive rate.²¹

We maintain that ultrasound is limited in its usefulness in the assessment of cases of spontaneous pneumothorax and following pleural procedures particularly in settings outside critical care. Many of these patients have underlying lung disease, particularly chronic obstructive pulmonary disease, which reduces the accuracy of pneumothorax detection by ultrasound.²¹ If a pneumothorax is detected by ultrasound, a CXR is usually required to assess its size (unless a CT scan is then performed). If the pneumothorax is so small as to be undetectable on CXR, then it is unlikely to require intervention and the use of ultrasound will not have changed the management.

We acknowledge that in the assessment of a supine patient thoracic ultrasound performed by a skilled operator may detect even small pneumothoraces (and effusions) and that if these patients require positive pressure ventilation detecting even a small amount of pleural air may be relevant. Even so, in this group, caution is needed because,