BLOOD AND ELECTROLYTE BALANCE IN THE POST-OPERATIVE CARDIAC PATIENT*

BY

M. V. BRAIMBRIDGE

(RECEIVED FOR PUBLICATION MAY 30, 1960)

A great deal has been written about post-operative fluid and electrolyte balance after abdominal operations. Relatively little attention has been paid to similar problems after cardiac surgery, particularly when extracorporeal circulation has been used. It may be valuable therefore to outline a post-operative régime used by most cardiac centres with minor modifications.

Consideration of blood and electrolyte balance after a cardiac operation may conveniently be divided into three sections: the maintenance of a normal blood volume, the problem of the unexpected blood pressure fall, and the necessity for supplying water and electrolyte requirements.

MAINTENANCE OF NORMAL BLOOD VOLUME

After cardiac surgery, the most important single factor in the immediate post-operative period is the maintenance of the normal blood volume. Too large a blood volume causes overloading of the ventricles, venous stasis, and pulmonary oedema. Too small a volume results in vasoconstriction, peripheral anoxia, and the picture of surgical shock.

In order to keep the volume constant, blood loss must be accurately replaced as it occurs. Blood is lost mainly into the thoracic cavity—pleura, pericardium, and mediastinum—though loss from thoracic and femoral wounds and blood withdrawn for haematological studies may assume significant proportions in children.

The immediate appearance of blood in calibrated chest drainage bottles is essential for its prompt replacement. These bottles should be tall and thin, so that small amounts of blood may be easily appreciated. A litre measuring cylinder is suitable if the top is cut off to take a bung, and the bottom anchored to a wider base, with 100 ml. of water for an underwater seal (Fig. 1).

Efficient drainage allows of immediate recognition of blood loss. Pleural drainage is commonplace in thoracic surgery and needs little mention, except to emphasize the importance of having the basal drain in the paravertebral gutter where blood accumulates. Tubes tend to swing forwards with the weight of their external length and it is often desirable to anchor the drain posteriorly inside the chest.

Pericardial drainage is more important than pleural. Tamponade is a common cause of death following cardiac surgery and accumulations too

FIG. 1.—Thoracic cavity drainage bottle used at the Brompton Hospital, London.

*A lecture given at the Institute of Diseases of the Chest, Brompton Hospital, London.

measurements, the technique is only possible in larger centres and is too complicated for routine use. It may be useful as an occasional check on the accuracy of the other methods.

UNEXPECTED BLOOD PRESSURE FALL

Attention to the régime described makes the post-operative course of the average cardiac patient straightforward and manageable by the nursing staff. Nevertheless the problem of an unexpected fall in blood pressure may occur. Haemorrhage is the immediate thought that arises, but there are at least six other conditions that may cause difficulty in diagnosis. These are cardiac failure, heart block, tamponade, vasodilatation, water and electrolyte imbalance, and adrenal exhaustion.

Unrecognized haemorrhage is the commonest cause of an unexpected blood pressure fall, and its frequency is proportional to the care that has been taken with haemostasis and drainage of the thoracic cavity. Haemostasis is particularly important if heparin has been used. Haemorrhage may occur with normal or with abnormal clotting. With normal clotting, haemorrhage may occur if a bleeding point has been left uncontrolled within the chest or if bleeding starts afresh. Haematological investigations will be normal, there will be clots in drainage tubes and bottles, and the chest wound will be dry. If bleeding is severe the patient must be returned to the operating theatre to secure proper haemostasis.

Failure to clot may be due to low levels of platelets or fibrinogen, the presence of fibrinolysins, or excessive use of anticoagulants such as heparin and "dindevan." Such bleeding may occur after any cardiac operation, normothermic, hypothermic, or extracorporeal. It is most frequent with deeply cyanosed patients after hypothermia where the circulation has been occluded longer than 10 minutes and after extracorporeal operations with inadequately cleaned or traumatic apparatus. It is recognized by an oozing thoracic wound, the complete absence of clots in chest drainage bottles and drains, and abnormal haematological studies. Most measurable clotting factors may be normal in the presence of an obvious bleeding state.

The differentiation of the type of haemorrhage is important. If haemostasis is at fault ligation of the bleeding vessel is essential. If there is abnormal clotting re-operation is useless and often dangerous. Re-opening the chest confronts the surgeon with widespread oozing and the patient is returned to bed with bleeding uncontrolled and sometimes with superadded cerebral symptoms.

Since detection of the clotting factor that is at fault is time consuming and often impossible, empirical treatment is necessary. If anticoagulants have been used before or during the operation, an extra dose of the appropriate antagonist is indicated. The triad of calcium, fibrinogen, and fresh blood will then be found to be effective, if used sufficiently early before the patient's blood volume has been replaced with stale, citrated, stored blood.

Enough calcium (10 ml. of 10% calcium gluconate per pint of transfused citrated blood) is injected. A bottle of fibrinogen is reconstituted and given intravenously. A donor is obtained and a sample of blood taken so that cross-matching may be begun at once. The fresh blood is then taken into a plastic bag and the patient immediately transfused. It is useful to cross-match relatives in advance before a severe operation where bleeding is anticipated since they will be available quickly if required. The use of fibrinogen is unpopular with blood banks since it is costly and inconvenient to make. This disadvantage must be weighed against its sometimes dramatic action in stopping haemorrhage even when the fibrinogen level of the patient is apparently normal.

Cardiac failure is the main differential diagnosis of haemorrhage. It is vital to differentiate these two, as rapid transfusion is the treatment of the latter but may be fatal in the former. Tachycardia, a low blood pressure, and vasoconstriction are common to both, but pulmonary oedema and a raised jugular venous pressure will favour cardiac failure. In cases of difficulty, a therapeutic test with the rapid injection of a small amount of blood, some 50 to 100 ml., may be helpful. If the condition is failure there will be little response; if due to haemorrhage the blood pressure usually rises sharply.

Heart block usually follows immediately after trauma to the bundle of His during closure of a septal defect. Complete heart block is recognized by a slow ventricular rate and cannon waves in the venous pressure. Continuous recording of the electrocardiogram on an oscilloscope has been found useful by both nurses and doctors post-operatively, since sudden alterations in rhythm or cardiac arrest are apparent and delay in treatment is minimal. Isoprenaline, 5-10 mg., may be given per rectum as often as is necessary to maintain an adequate heart rate. If this fails external electrodes may be placed over the heart and connected to a pacemaker. A higher voltage is necessary than when leads have been left in the heart, and twitching of the pectoral muscles, pain, and even burns distress

the patient. The insertion of an insulated stainless steel wire into the ventricle through a wide-bore needle avoids these disadvantages.

Tamponade is a real hazard in cardiac surgery and minor degrees of it often go unrecognized, causing unexplained tachycardia and poor immediate post-operative progress. It is due to inadequate pericardial drainage. Measures to counteract this have already been outlined. Tamponade mimics cardiac failure, with tachycardia, low blood pressure, and a raised jugular venous pressure. A marked pulsus paradoxus has been found to be the best differentiating feature and has allowed immediate thoracotomy to avert disaster on more than one occasion.

Vasodilatation, particularly following extracorporeal circulation, sometimes causes a fall in blood pressure. A patient may return to the ward vasoconstricted, pale, with poor peripheral pulses. Some hours later he may suddenly vasodilate, become pink and warm, look much better but be markedly hypotensive. The reason for this vasodilatation is obscure, since it occurs even when the blood volume is normal, whether calculated by mathematical balance of blood loss, the patient's weight, or radioactive isotope studies. The treatment is blood transfusion in excess of theoretical requirements, with an amount that may vary from 300 ml. in children to 1,000 ml. in adults. With experience, this vasodilatory episode may be anticipated by maintaining a positive blood balance during the vasoconstrictive phase, so that, when the anticipated vasodilatation occurs, serious hypotension will be avoided.

Water and electrolyte imbalance does not cause a sudden drop in blood pressure, but may result in troublesome hypotension if oral intake is restricted for any reason, and care has not been taken to replace them. If the calculated inevitable loss of water is replaced as a routine post-operative measure, imbalance may be ignored in the diagnosis of a fall in blood pressure.

Adrenal exhaustion is a rare cause of post-operative hypotension. Sometimes, even when intravenous noradrenaline fails to raise the blood pressure, a good response may be obtained by an injection of cortisone. Consequently cortisone is often given during the treatment of obscure post-operative hypotension, but it is rare for it to be of proven value.

WATER AND ELECTROLYTE REQUIREMENTS

The necessity for post-operative blood balance is apparent. Less obvious is the importance of fluid balance. There are two policies that may be adopted in this respect. The estimated

inevitable loss of water and electrolytes may be replaced as a routine, or they may be ignored and oral feeding relied upon to correct any discrepancy. The latter is the common practice in most thoracic surgical centres and works satisfactorily in the main. For cardiac cases, particularly after extracorporeal circulation operations, the maintenance of fluid balance is always important and sometimes vital. Oral feeding is not always resumed at once; nausea, vomiting, or cerebral symptoms may cause further delay, and, if provision has not been made to cover each day's losses as they occur, a dangerous state of dehydration may result. Replacement of expected fluid loss is simple, and will avert the hypotension and anuria of water and electrolyte depletion.

WATER.—There are few figures available for the amount of water inevitably lost by a patient following cardiac surgery. Careful studies have been made at the Mayo Clinic on the average losses of water following total heart/lung bypass operations (Sturtz, Kirklin, Burke, and Power, 1957). Water was lost as obligatory urine, insensible water loss from lungs and skin, and abnormal losses due to vomiting and thoracic drainage not replaced with blood. Against this was set the gain of water of oxidation and of catabolism of fat, carbohydrate, and protein, the so-called metabolic water (Table I).

TABLE I
POST-PERFUSION WATER LOSS (MAYO CLINIC)

Water Loss (ml./m. ² /day)	First Day	Subsequent Days
Obligatory urine	294	465
Insensible loss	469	420
Abnormal loss	0	150
minus		
Metabolic water	270	270
	493 ml./m. ²	765 ml./m. ²

From these figures it can be seen that during the first 24 hours after operation some 500 ml. of water per square metre of surface area of the patient was lost. This is two-thirds of the loss on the second day that was some 750 ml. per sq. metre. The loss on the third day was almost identical.

Nurses change and fluid balances are drawn up at 8 a.m., so it is convenient to reckon a day as starting at this time. After a major cardiac operation, there are usually about 18 of the 24 hours remaining, that is, three-quarters of the day. Only two-thirds of the subsequent day's water requirements are needed on the first day, $\frac{2}{3} \times \frac{3}{4} = \frac{1}{2}$; so it may be said that 750 ml. of water per sq.

metre will replace inevitable loss on the second and third days with half of this on the day of operation. At the Mayo Clinic, cold, humid oxygen tents were used post-operatively, and it is wise to increase this amount as a routine to 800 ml., with 400 ml. on the day of operation.

SODIUM.—The question of whether cardiac patients should post-operatively be given sodium is difficult to answer dogmatically. On the one hand, it may be said that a patient in heart failure will retain sodium, which, combined with the normal post-operative sodium retention, will make further administration of it dangerous. On the other hand, most patients with any degree of heart failure will have been energetically treated with a low salt intake and diuretics. If too energetic a régime of dehydration has been adopted before the operation, sodium will definitely be required post-operatively.

Sodium is less obviously required when salt has not been restricted. At Stanford University Hospital, San Francisco, dextrose 5% was at one time given as intravenous fluid replacement following all cardiac surgery. Electrolyte imbalance, sometimes serious, tended to occur if there were any delay in the resumption of oral intake. Nausea, vomiting, and hypotension were the sequels and a vicious circle was begun.

Normal sodium requirements are 50 mEq. per sq. metre of surface area per day (Snively and Sweeney, 1956). Taking into consideration post-operative sodium retention, 35 mEq. per sq. metre was arbitrarily chosen as the amount to be given on the average post-operative day with half of this on the day of operation. On this régime, electrolyte imbalance ceased to be a problem.

POTASSIUM.—Potassium is only rarely required following cardiac surgery. It is necessary when cerebral symptoms have prevented oral intake for more than 48 hours, or when too enthusiastic a course of dehydration has been carried out with chlorothiazide or hydroflumethiazide without potassium supplements.

ENERGY.—Energy requirements need to be covered, particularly in children, in order to prevent ketosis. Normal energy requirements are met by 50 g. of carbohydrate per sq. metre of surface area per day, so 40 g. is enough, with half of this amount on the day of operation.

ADMINISTRATION OF WATER AND ELECTROLYTE REQUIREMENTS.—On an average post-operative day 800 ml. of water, 35 mEq. of sodium, and 40 g. of carbohydrate per sq. metre of surface area will cover the minimal needs of the cardiac patient

without danger of overloading the circulation. Eight hundred millilitres of one-third normal saline in 5% dextrose contains these quantities of water, sodium, and energy. Half this amount, 400 ml. per sq. metre, may be given on the day of operation ending at 8 a.m. the following morning. One-third normal saline is not a standard solution in Great Britain. It may be made up by dispensaries, or one-fifth normal saline in dextrose used. This will give 22 mEq. of sodium in the same 800 ml. of solution, which is probably adequate. The surface area is read from the DuBois nomogram from the height and weight of the patient. The exact quantity of fluid necessary is calculated therefrom.

The route of administration is initially intravenous. A blood drip is always running in the post-operative cardiac patient, and a separate transfusion set for the dextrose/saline may be inserted into the side of the blood transfusion tubing. Whenever the blood balance is level, blood may be turned off and dextrose/saline continued instead. This is a positive advantage in post-operative nursing, since the blood does not have to be transfused so slowly that the cannula in the vein clots. The calculated quantity of fluid is given by 8 a.m. the following morning, both blood and fluid balance being maintained by relatively simple adjustments with the transfusions.

Oral feeding may be begun when convenient. Thirst is a common symptom after severe cardiac operations, and is particularly prominent after extracorporeal circulation when it is seldom relieved even by large quantities of water by mouth. These quantities given orally may cause vomiting with further disturbance of electrolyte balance, and it is wise to restrict oral intake on the day of operation to small amounts. If this adds up to a significant volume, the intravenous administration of fluid is curtailed by the same quantity.

The régime set out here, while based on that originally recommended by the Mayo Clinic for the post-operative care of their open heart cases, differs from it in two important respects. Oral fluid is not totally restricted during the immediate post-operative period, and sodium is given intravenously. Too literal following of the Mayo Clinic dicta resulted in severe electrolyte difficulties at Stanford University Hospital before it was realized that the restriction of oral fluid at the Mayo Clinic was due to the nausea of ether anaesthesia, and of sodium due to the relatively large amount transfused by the Gibbon pump oxygenator.

SUMMARY

The normal blood volume is maintained in the post-operative cardiac patient by the accurate replacement of blood loss as it occurs. Efficient drainage of pleura, pericardium, and mediastinum ensures the prompt appearance of blood loss in calibrated chest drainage bottles. Accurate replacement is then simple.

Monitors of blood balance are mathematical, using a chart of progressive totals of blood loss and replacement; clinical, taking into account arterial and venous pressures and weight; and special, with radioactive isotope tagged red cells.

Unexpected hypotension occurs. While this is often due to unrecognized haemorrhage, associated with either normal or abnormal clotting factors, other conditions may cause difficulty in diagnosis. These are cardiac failure, heart block, tamponade, vasodilatation, water and electrolyte imbalance, and adrenal exhaustion.

Water and electrolyte balance may be little less important than blood balance. A simple régime, covering inevitable loss without circulatory overloading, is the administration of 800 ml. of one-third normal saline in 5% dextrose per sq. metre of surface area on an average post-operative day with half of this on the day of operation. Any delay in the start of oral feeding will not then result in dehydration.

I am particularly grateful to Dr. Frank Gerbode for stimulating my interest in this subject while working in his department as Heller Fellow in Surgery at Stanford University, San Francisco. I am also indebted to Sir Russell Brock for further opportunities to study this subject and for his help and encouragement in the preparation of this paper.

REFERENCES

- Emanuel, R. (1960). *Thorax*, 15, 00.
 Snively, W. D., Jr., and Sweeney, M. J. (1956). *Fluid Balance Handbook for Practitioners*. Charles C. Thomas, Springfield, Illinois.
 Sturtz, G. S., Kirklin, J. W., Burke, E. C., and Power, M. H. (1957). *Circulation*, 16, 988.