Gallium-67 as a diagnostic tool in the evaluation of peripheral pulmonary lesions

B. T. le ROUX and A. E. HOULDER

The Thoracic Surgical Unit, Wentworth Hospital, and the Department of Radiophysics, Addington Hospital, Durban

le Roux, B. T., and Houlder, A. E. (1974). *Thorax*, 29, 355–358. **Gallium-67 as a diagnostic tool in the evaluation of peripheral pulmonary lesions.** While Gallium-67 shows a high affinity for primary pulmonary tumours, benign pulmonary lesions also concentrate the isotope. In 50 consecutive patients in whom the diagnosis of pulmonary carcinoma was not achieved by other means, the exhibition of ^{67}Ga showed a positive scan in 36, of whom three, on histological evidence, certainly did not have a pulmonary carcinoma, and in three others radiographic diminution in the extent of the peripheral pulmonary lesion makes most unlikely the diagnosis of carcinoma. In seven of 14 patients in whom the scan was negative it was judged necessary to undertake surgical management. Scanning with ^{67}Ga may modify the extent of surgical resection in some with a negative scan but does not establish the diagnosis of carcinoma in those with a positive scan, is unsafe as a screening test on which to base the decision on the need for referral to a thoracic surgeon, and in particular does not distinguish between the three common causes of 'chronic lung abscess'—chronic destructive pneumonia, cavitated carcinoma, and tuberculosis with cavitation and without tubercle bacilli in the sputum.

Gallium-67 is a cyclotron-produced isotope which in a carrier-free form has been shown to have a high affinity for malignant tissues which appears to be independent of histological type. The initial studies of ^{67}Ga were performed by Bruner, Hayes, and Perkinson (1953) as part of an attempt to use gallium isotope for specific therapy of bone tumours. It was while scanning the bones of a patient with Hodgkin's disease that the tumour-specific properties of ^{67}Ga citrate were discovered. Edward and Hayes (1969; 1970) and, since that time, numerous others have confirmed the tumour-specific properties of this isotope. Reports concerning the uptake of ^{67}Ga citrate in primary lung cancer show that the detection efficiency is greater than 90%.

However, benign lesions in the lungs also concentrate ^{67}Ga citrate. Positive scans have been obtained in active pulmonary tuberculosis, silicosis, acute infections, and especially with recently developed sarcoid. Healed tuberculosis, chronic infections, bacterial and viral pneumonias, pulmonary infarction, and pleural lesions generally give negative results. van der Schoot, van Marle-van der Goot, Groen, and de Jong (1972) found

a positive scan in almost 40% of 58 cases with established benign pulmonary lesions.

The purpose of this paper is to report a study of the use of ^{67}Ga in 50 consecutive patients. The objects of the trial were:

1. to confirm the high detection efficiency of ^{67}Ga in primary pulmonary tumours;
2. to evaluate its use in a series of patients in whom the chest radiograph showed a lesion of 'coin' type and in whom sputum had been shown not to contain tubercle bacilli;
3. to determine whether the routine use of ^{67}Ga as a diagnostic tool in the evaluation of 'peripheral' pulmonary lesions increased the accuracy of preoperative diagnosis, so as not only to modify the extent of pulmonary resection but also possibly to obviate the need for an aggressive surgical approach.

The trial was limited to 50 patients in whom the diagnosis of pulmonary carcinoma was not achieved by other means. It follows that most had peripheral, spherical, pulmonary opacities, some cavitated. In only one patient had tumour cells been demonstrated in a single specimen of sputum. In all but two patients the appearances at bron-

TABLE

Group	Age (yr)	Caucasian		Mulatto		Asiatic		African		Total
		M	F	M	F	M	F	M	F	
A Positive ⁶⁷ Ga scan with histological confirmation of the diagnosis of carcinoma	< 30							2	1	0
	31-40							7		3
	41-50	1	1			1		4		2
	51-60	1						1		2
	61-70	3						1		4
										16
B Positive ⁶⁷ Ga scan. Diagnosis of carcinoma likely on clinical grounds. No histological confirmation	< 30									0
	31-40									0
	41-50	1	1					2		4
	51-60	1						4		5
	61-70	2	1(72)					2		5
										14
C Positive ⁶⁷ Ga scan. Lesion shown histologically (3 patients) or presumed on radiological evidence (3 patients) not to be a carcinoma	< 30									0
	31-40							2	1	3
	41-50							2		2
	51-60							1		1
	61-70									0
										6
D Negative ⁶⁷ Ga scan. Diagnosis of carcinoma excluded on histological evidence or reasonable clinical grounds	< 30			1(21)						1
	31-40					1		1		2
	41-50					1		1		2
	51-60	1		1			1	4		7
	61-70	2								2
										14
Total		12	3	2	—	3	1	27	2	50

Males 44; Females 6

choscopy were normal. In two, each with the radiographic appearance of shrinkage of the left upper lobe, the left upper bronchus was seen at bronchoscopy to be compressed, but the mucosa looked normal and a biopsy from within the compressed bronchus showed normal bronchial epithelium.

The age range and sex of the patients are shown in the Table. In this the patients are grouped according to the final assessment of the results of the scan and not in the order in which they were investigated. Most (44) were men. The youngest was a mulatto male of 21 years and the oldest a Caucasian female of 72. All but nine patients were older than 40 years; 11 were older than 60 and 32 were in the fifth and sixth decades. The race groups are represented in a more or less proper local proportion—there is a slight emphasis on Caucasians.

METHOD

⁶⁷Ga citrate was injected intravenously with an approximate dose of 35 μ Ci per kilogram (2 to 3 mCi per adult). ⁶⁷Ga decays by electron capture with a half-life of 78 hours yielding no beta rays. The gamma ray energies are 93 KeV (40%), 184 KeV (24%), 296 KeV (22%), and 388 KeV (7%). The 184 and 296 KeV gamma rays are used for imaging purposes. Scanning was carried out at 48 hours after the injection of the dose, using a rectilinear colour scanner with 5-inch crystal. Only the anterior or the posterior surface of the chest was normally scanned.

RESULTS

Of the 50 patients the ⁶⁷Ga scan was positive in 36 and negative in 14. In 16 patients (group A) with a positive ⁶⁷Ga scan the diagnosis of a malignant tumour was confirmed by pulmonary resection or some other form of surgical manoeuvre which allowed of biopsy. In one patient the tumour was mediastinal and a reticulum-cell sarcoma; in 15 the tumour was a pulmonary carcinoma—squamous in eight, an adenocarcinoma in three, and undifferentiated in four (called oat-cell in one of these). Pneumonectomy was undertaken in six patients and a lobectomy in four. In one patient with a squamous carcinoma tumour cells had been found on examination of one specimen of sputum, a finding which could not be repeated. In the patient with the mediastinal lymphoma the diagnosis was achieved at mediastinoscopy. Formal thoracotomy was undertaken in 12 patients in this group of 16, and in the remaining four a diagnosis was achieved histologically as a preliminary to a form of treatment alternative to pulmonary resection because it was recognized that the diagnosis of tumour was virtually certain and that this had extended beyond the limits of surgical management.

In 14 patients (group B) in whom the ⁶⁷Ga scan was positive, the diagnosis of carcinoma seems inescapable despite absence of histological confirmation. In three of these 14 patients the tumour was of the Pancoast variety and associated with

radiographic evidence of rib destruction. One patient with a left peripheral tumour was hoarse of voice; in two others there was gross oesophageal displacement, also accepted as evidence of mediastinal invasion. Three, who on all counts were suitable for surgical management, declined thoracotomy; a fourth had such severe angina that management by irradiation was preferred and the lesion has been seen to diminish; four were so frail that they were judged unsuitable for any form of management. The three with radiographic evidence of parietal invasion, the three with mediastinal invasion, and the one with angina whose lesion has regressed with irradiation certainly have pulmonary tumours, and, on all clinical criteria, these are primarily pulmonary. The three who continue to decline thoracotomy and the four too frail for surgical, or other, management would be regarded, on all available evidence, as having primary pulmonary malignant tumours, and growth of the primary lesion observed in four of these seven over the past six months tends further to substantiate the diagnosis.

In one patient in group B a large left peripheral pulmonary shadow, abutting on the chest wall and associated with medial rib erosion, has been shown by percutaneous aspiration biopsy to be a squamous carcinoma. In this patient there is an additional contralateral 2 cm peripheral pulmonary opacity presumed on radiological grounds to represent a pulmonopulmonary metastasis. The large left-sided lesion showed a strongly positive ^{67}Ga scan; the small right lesion did not. Medial chest wall invasion was the contra-indication to surgical management. On clinical grounds the right-sided lesion would, in the absence of contralateral medial chest wall invasion, have been accepted as evidence of haematogenous metastasis and therefore a contra-indication to management by left thoracotomy. The negative right-sided scan is judged on this evidence to represent failure of the metastasis either to take up ^{67}Ga or, because of size, to be demonstrable by the scanning technique used.

In six patients (group C), each with a positive ^{67}Ga scan, there is histological (in three) or radiographic (in three) evidence that the lesion is not a carcinoma. In two, the resected lesion was shown to be a chronic destructive pneumonia (chronic pyogenic lung abscess) and in the third the resected lesion was caseating tuberculosis, which was radiographically ingravescent despite a well-supervised four-week course of antituberculosis chemotherapy. Rapid diminution of a

right upper pulmonary cavity in response to chemotherapy and physiotherapy makes the diagnosis of carcinoma most unlikely in a fourth, in whom bronchoscopic appearances were normal; in the remaining two, who have bronchoscopically normal bronchi, non-cavitated pulmonary opacities are diminishing slowly and are judged, on radiographic grounds, unlikely to represent carcinomata. In these last-mentioned two the evidence is presumptive and inconclusive but it has been judged on clinical and radiographic grounds unnecessary to proceed to diagnostic thoracotomy, despite the positive ^{67}Ga scan.

In 14 patients (group D) the results of scanning after the exhibition of ^{67}Ga were negative. Seven of these 14 patients were submitted to thoracotomy because of a well-defined peripheral pulmonary opacity, cavitated in one. In three patients the lesion was an hydatid cyst; in two the lesion was tuberculoma; in one a chondroadenoma (hamartoma); and in the last a 'plasma-cell granuloma'. In four of these patients the lesions were enucleated. In three—one an effete infected hydatid, another the 'plasma-cell granuloma', and the third a cavitated tuberculous lesion which was radiographically ingravescent despite good drug therapy—management was by lobectomy in the belief that the lesion was probably a carcinoma despite the negative scan. If cognisance had been taken of the negative scan two of these lesions could probably have been managed successfully by enucleation or subsegmental resection. The cavitated tuberculous lesion occupied most of the right upper lobe. In the remaining seven patients response in two to antituberculosis chemotherapy and in three to other forms of chemotherapy make certain the diagnosis of a tuberculous or other inflammatory lesion. The remaining two patients are elderly and frail and on general grounds unsuitable for surgical management. One has multiple lesions and has been known in the past to have tuberculosis and these lesions are probably tuberculous. The final diagnosis in these two patients is uncertain.

DISCUSSION

There are a number of reports concerning the uptake of ^{67}Ga in lung cancer. Langhammer *et al.* (1972) reviewed the results of four European university centres, Amsterdam, Berlin, Giessen, and Munich, where striking tumour concentration of ^{67}Ga was found in 62 of 70 patients with bronchial carcinoma. (In a previous communication van der Schoot *et al.* (1971) from Amsterdam

University reported on 92% positive ^{67}Ga scans in bronchial carcinoma in 48 out of 54 cases.) Merrick *et al.* (1972) found gallium concentrated in all primary lung cancers in 21 patients. Higasi *et al.* (1972) found that all but one of 19 primary lung cancers showed positive scans, the size of the tumour missed being a 2×1.5 cm adenocarcinoma. At Tohoku University, Ito *et al.* (1971) found that 21 out of 21 cases of pulmonary cancer were positive with ^{67}Ga scanning. These results indicate that ^{67}Ga is taken up in almost all primary pulmonary cancers; lesions which have been missed may be too small. These facts are undisputed. But does the exhibition of ^{67}Ga in any way modify the surgical attitude towards an isolated peripheral pulmonary lesion in a patient in whom the use of alternative diagnostic aids has failed to provide an answer and a sufficient period of time has been allowed to elapse to give inflammatory lesions the opportunity to regress? Is a positive scan unequivocal evidence for a diagnosis of carcinoma? Does a negative scan exonerate the surgeon from an aggressive approach? Is the easily available technique of scanning with ^{67}Ga one which may be used by physicians as a screening test on which to base the decision on whether to submit a patient for thoracic surgical opinion?

The analysis of the results of scanning after the exhibition of ^{67}Ga in 50 consecutive patients makes clear that the answer to all these questions is an unequivocal no. Surgical attitude is unmodified by the use of ^{67}Ga additional to other routine diagnostic aids; the extent of surgical resection may occasionally be modified if the ^{67}Ga scan is negative and frozen section examination is available at the time of thoracotomy; a negative scan does not obviate the need for thoracotomy; it may be that a lesion which does not show a positive scan with gallium is in fact a carcinoma; and a positive scan is not in itself evidence of carcinoma. In particular, ^{67}Ga failed to distinguish between the three common local

causes of 'chronic lung abscess'—destructive pneumonia, cavitated carcinoma, and tuberculosis without tubercle bacilli in the sputum.

REFERENCES

- Bruner, H. D., Hayes, R. L., and Perkinson, J. D. (1953). Preliminary data on gallium-67. *Radiology*, **61**, 602.
- Edwards, C. L. and Hayes, R. L. (1969). Tumour scanning with gallium-67 citrate. *Journal of Nuclear Medicine*, **10**, 103.
- and — (1970). Scanning malignant neoplasms with gallium-67. *Journal of the American Medical Association*, **212**, 1182.
- Higasi, T., Nakayama, Y., Murata, A., Nakamura, K., Sugiyama, M., Kawaguchi, T., and Suzuki, S. (1972). Clinical evaluation of gallium-67 citrate scanning. *Journal of Nuclear Medicine*, **13**, 196.
- Ito, Y., Okuyama, S., Awano, T., Takahashi, K., Sato, T., and Kanno, I. (1971). Diagnostic evaluation of gallium-67 scanning of lung cancer and other disease. *Radiology*, **101**, 355.
- Langhammer, H., Galubitt, G., Grebe, S. F., Hampe, J. F., Haubold, U., Hör, G., Kaul, A., Koeppe, P., Koppenhagen, J., Roedler, H. D., and van der Schoot, J. B. (1972). Gallium-67 for tumor scanning. *Journal of Nuclear Medicine*, **13**, 25.
- Merrick, M. V., Gunasekera, S. W., Lavender, J. P., Nunn, A. D., Thakur, M. L., and Williams, E. D. (1972). The use of indium-111 for tumour localization. International Atomic Energy Agency.
- van der Schoot, J. B., Groen, A. S., and De Jong, J. (1971). Gallium-67 scintigraphy in lung disease. 2nd Congress of the European Association of Radiology.
- , van Marle-van der Goot, M., Groen, A. S., and De Jong, J. (1972). Symposium on medical radioisotope scintigraphy. Gallium-67 scintigraphy in benign lung disease. International Atomic Energy Agency.

Requests for reprints to: Professor B. T. le Roux, The Thoracic Surgical Unit, Wentworth Hospital, Durban.