THORAX

Editorials

Leukotrienes and aspirin induced asthma

Leukotrienes are lipid mediators formed from the oxidative metabolism of arachidonic acid via the 5-lipoxygenase enzyme cascade.¹ The sulphidopeptide leukotrienes comprise leukotriene (LT) C₄ and LTD₄, and LTE₄. These molecules are responsible for the activity previously recognised as slow reacting substance of anaphylaxis.

The sulphidopeptide leukotrienes are potent constrictors of the smooth muscle of the airways,² and may also contribute to the bronchial hyperresponsiveness characteristic of asthma.³ Lam *et al* used fast atom bombardment mass spectroscopy to analyse bronchoalveolar lavage fluid for the presence of leukotrienes.⁴ LTE₄ was the predominant leukotriene recovered, although LTD₄ was also identified. Significant quantities of LTC₄ have been detected by radioimmunoassay in the bronchoalveolar lavage fluid of symptomatic asthmatic patients compared with normal control subjects,⁵ and increased quantities of leukotrienes have been detected in bronchoalveolar lavage fluid after local endobronchial challenge with allergen.⁶

In man LTC₄ is converted enzymatically in the blood to LTD₄ and LTE₄ which is excreted in the urine. After infusion of radiolabelled LTC₄ 12-20% appears in the urine, of which 4-6% of the total infused dose is seen as LTE₄.⁷ Measurement of the urinary LTE₄ concentration may therefore act as a marker for the systemic release of the sulphidopeptide leukotrienes. Urinary LTE₄ levels are increased during acute severe asthma,⁸ following allergen challenge,^{8,9} and in aspirin induced asthmatic responses.¹⁰

There is compelling evidence that aspirin induced asthma is mediated by 5-lipoxygenase pathway products and, in particular, by the sulphidopeptide leukotrienes. A proportion of patients with asthma are intolerant of aspirin.¹¹ In these individuals aspirin ingestion is followed by the onset of wheezing, rhinitis, urticaria, or anaphylaxis. These symptoms may occur singly or in combination. It has been suggested that aspirin induced asthma may relate to the inhibition of cyclooxygenase, resulting in the "shunting" of arachidonate metabolism to the 5-lipoxygenase pathway. This, in turn, will lead to the increased generation of leukotrienes in susceptible individuals.¹² While there is increasing evidence for the enhanced generation of leukotrienes in aspirin induced asthmatic responses, the evidence that this is a result of shunting of arachidonic acid metabolism from the cyclooxygenase to the 5-lipoxygenase pathway is lacking.

Ferreri *et al* measured mediator release in the nasal lavage fluids of aspirin sensitive asthmatic patients after aspirin challenge.¹³ The release of LTC₄ into nasal secretions was noted but there was no decrease in the levels of prostaglandin E₂ (PGE₂). Urinary levels of LTE₄ have also been measured before and after aspirin provocation in patients with aspirin sensitivity. The baseline LTE₄ levels are significantly raised in asthmatic patients with aspirin sensitivity compared with those who are tolerant of aspirin.¹⁴ Following provocation with oral aspirin there

is a further fourfold increase in urinary LTE₄ excretion in aspirin sensitive individuals.¹⁴⁻¹⁷ Urinary concentrations of the thromboxane A₂ metabolite, 11-dehydrothromboxane B₂, were unchanged.¹⁷ Increased release of LTE₄ into the urine of patients with aspirin sensitivity is also seen after asthma induced by lysine-aspirin in susceptible subjects,^{17,18} suggesting a pulmonary source of the leukotrienes. These cumulative observations strongly support the view that there is an upregulation of arachidonate metabolism in aspirin sensitive patients resulting in the augmented production of the sulphidopeptide leukotrienes.

If sulphidopeptide leukotrienes play a significant part in the pathogenesis of asthma, then attempts to inhibit their generation or antagonise their action should be of some benefit. A number of selective leukotriene receptor antagonists and 5-lipoxygenase inhibitors have now been developed and many are undergoing clinical trials in patients with disease. These assessments are still in their early stages, but several studies have shown bronchodilatation after administration of leukotriene antagonists and 5-lipoxygenase inhibitors in asthmatic patients but not in normal subjects, suggesting that leukotrienes may contribute to the resting airways tone in asthma.¹⁹⁻²² Furthermore, leukotriene receptor antagonists inhibit asthmatic responses induced by aspirin,^{23,24} exercise,^{25,26} and allergens.^{27,28} In one study leukotriene receptor antagonism reduced allergen induced enhancement in bronchial hyperresponsiveness.²⁷ Zileuton, a novel 5-lipoxygenase inhibitor, protected against asthma induced by hyperventilation²⁹ and attenuated aspirin induced asthmatic responses.³⁰

Only limited data are available on the effects of leukotriene antagonism and 5-lipoxygenase inhibitors in chronic asthma. Preliminary evidence suggests that these new classes of drugs produce a significant mild improvement in airways function and a reduction in symptoms.³¹⁻³⁵

The interesting study reported by Dahlén and colleagues in this issue of *Thorax* (pp 1205-10) in aspirin intolerant patients³⁶ extends the previous observations on the bronchodilating properties of sulphidopeptide leukotriene antagonists. Eight subjects were studied on two separate days, when they received 825 mg MK-0679 or placebo, orally, in a double blind randomised crossover study. MK-0679 elicited bronchodilatation which lasted for at least nine hours. The maximal improvement in FEV₁ ranged from 5% to 34% above predrug baseline values, and the degree of bronchodilatation correlated with the severity of asthma and aspirin sensitivity. These findings provide further evidence that leukotriene dependent tone may be an important component of the persistent airway obstruction in asthmatic patients. In view of the clinical heterogeneity of bronchial asthma, it would be interesting to define a subgroup of individuals who respond particularly well to leukotriene antagonism. In this regard it would be informative to

evaluate whether there is a correlation between the bronchodilating effect of MK-0679 and basal urinary LTE₄ concentrations. Although there is strong evidence for the involvement of leukotrienes in aspirin induced asthma, it is not possible to establish from the study by Dahlén *et al* whether it was the severity of asthma or the degree of aspirin sensitivity which was the critical determinant of the bronchodilator response to MK-0679. Nevertheless, the present findings support the view that leukotriene receptor antagonists offer promise in the treatment of patients with asthma; patients with aspirin induced asthma may be particularly helped by these drugs. The recent observation that inhalation of LTE₄ results in eosinophilic accumulation in asthmatic airways³⁷ has suggested that this new class of therapeutic agents may have anti-inflammatory and steroid-sparing potential.

TAK H LEE
PANDORA E CHRISTIE
Department of Allergy and
Allied Respiratory Disorders,
Guy's Hospital,
London SE1 9RT

Reprint requests to: Professor T H Lee

- Samuelsson B. Leukotrienes: mediators of hypersensitivity reactions and inflammation. *Science* 1983;220:568-75.
- Drazen JM, Austen KF, Lewis RA, Corey EJ. Comparative airway and vascular activities of leukotrienes C-1 and D *in vivo* and *in vitro*. *Proc Natl Acad Sci USA* 1980;77:4354-8.
- Arm JP, Spur BW, Lee TH. The effects of inhaled leukotriene E₄ on the airway responsiveness to histamine in subjects with asthma and normal subjects. *J Allergy Clin Immunol* 1988;82:654-60.
- Lam S, Chan H, LeRiche JC, Chan-Yeung M, Salari H. Release of leukotrienes in patients with bronchial asthma. *J Allergy Clin Immunol* 1988;81:711-7.
- Wardlaw AJ, Hay H, Cromwell O, Collins JV, Kay AB. Leukotrienes, LTC₄ and LTB₄, bronchoalveolar lavage in bronchial asthma and other respiratory diseases. *J Allergy Clin Immunol* 1989;84:19-26.
- Wenzel SE, Larsen GL, Johnston K, Voelkel NF, Westcoff JY. Elevated levels of leukotriene C₄ in bronchoalveolar lavage from atopic asthmatics after endobronchial allergen challenge. *Am Rev Respir Dis* 1990;142:112-9.
- Maltby NH, Taylor GW, Ritter JM, Moore K, Fuller RW, Dollery CT. Leukotriene C₄ elimination and metabolism in man. *J Allergy Clin Immunol* 1990;85:3-9.
- Taylor GW, Taylor I, Black P, Turner N, Taylor I, Maltby NH, *et al*. Urinary leukotriene E₄ after antigen challenge and in acute asthma and allergic rhinitis. *Lancet* 1989;i:584-8.
- Manning PJ, Rokach J, Malo J-L, Ethier D, Cartier A, Girard Y, *et al*. Urinary leukotriene E₄ levels during early and late asthmatic responses. *J Allergy Clin Immunol* 1990;86:211-20.
- Smith CM, Christie PE, Hawksworth RJ, Thien F, Lee TH. Urinary leukotriene E₄ levels following allergen and exercise challenge in bronchial asthma. *Am Rev Respir Dis* 1991;143:1411-3.
- Samter M, Beers RF. Intolerance to aspirin: clinical studies and consideration of its pathogenesis. *Ann Intern Med* 1968;68:975-83.
- Szczeklik A, Gryglewski RJ, Czerniawska-Mysik G. Relationship of inhibition of prostaglandin biosynthesis by analgesics to asthma attacks in aspirin-sensitive patients. *BMJ* 1975;1:67-9.
- Ferreri NR, Howland WC, Stevenson DD, Spiegelberg HL. Release of leukotrienes, prostaglandins and histamine into nasal secretions of aspirin-sensitive asthmatics during reaction to aspirin. *Am Rev Respir Dis* 1988;137:847-54.
- Christie PE, Tagari P, Ford-Hutchinson AW, Charlessen S, Chee P, Arm JP, *et al*. Urinary leukotriene E₄ concentrations increase after aspirin challenge in aspirin-sensitive asthmatic subjects. *Am Rev Respir Dis* 1991;143:1025-9.
- Smith CM, Hawksworth RJ, Thien FCK, Christie PE, Lee TH. Urinary leukotriene E₄ in bronchial asthma. *Eur Respir J* 1992;5:693-9.
- Knapp HR, Sladek K, Fitzgerald GA. Increased excretion of leukotriene E₄ during aspirin-induced asthma. *J Lab Clin Med* 1992;119:48-51.
- Kumlin M, Dahlén B, Björck T, Zetterstrom O, Granstrom E, Dahlén S-E. Urinary excretion of leukotriene E₄ and 11-dehydrothromboxane B₂ in response to bronchial provocations with allergen, aspirin, leukotriene D₄ and histamine in asthmatics. *Am Rev Respir Dis* 1992;146:96-103.
- Christie PE, Tagari P, Ford-Hutchinson AW, Black C, Markendorf A, Schmitz-Schumann M, *et al*. Urinary leukotriene E₄ after lysine-aspirin inhalation in asthmatic subjects. *Am Rev Respir Dis* 1992;146:1531-4.
- Hui KP, Barnes NC. Lung function improvement in asthma with a cysteinyl-leukotriene receptor antagonist. *Lancet* 1991;337:1062-3.
- Gaddy JN, Margolskee DJ, Bush RK, Williams VC, Busse WW. Bronchodilation with a potent and selective leukotriene D₄ (LTD₄) receptor antagonist (MK-571) in asthma patients. *Am Rev Respir Dis* 1992;146:358-63.
- Israel E, Rubin P, Pearlman H, Cohn J, Drazen J. 5-lipoxygenase inhibition by Zileuton causes acute bronchodilation in asthma. *Am Rev Respir Dis* 1992;145:A16.
- Kips JC, Joos GF, De Lepeleire I, Joos G, Williams V, Buntix A, *et al*. MK-571, a potent antagonist of leukotriene D₄-induced bronchoconstriction in the human. *Am Rev Respir Dis* 1991;144:617-21.
- Christie PE, Spur BW, Lee TH. The effect of inhalation of the leukotriene receptor antagonist SK&F 104353, on leukotriene C₄ and leukotriene E₄-induced bronchoconstriction in subjects with asthma. *J Allergy Clin Immunol* 1991;88:193-8.
- Dahlén B, Kumlin M, Margolskee DJ, Larsson C, Blomqvist H, Williams VC, *et al*. The leukotriene-receptor antagonist MK-0679 blocks airway obstruction induced by bronchial provocation with lysine-aspirin in aspirin-sensitive asthmatics. *Eur Respir J* 1993;6:1018-26.
- Manning PJ, Watson RM, Margolskee DJ, Williams VC, Schwartz JL, O'Byrne PM. Inhibition of exercise-induced bronchoconstriction by MK-571, a potent leukotriene D₄ antagonist. *N Engl J Med* 1990;323:1736-9.
- Finnerty JP, Wood-Baker R, Thomson H, Holgate ST. Role of leukotrienes in exercise-induced asthma; inhibitory effect of ICI 204,219, a potent leukotriene D₄ receptor antagonist. *Am Rev Respir Dis* 1992;145:746-9.
- Taylor IK, O'Shaughnessy KM, Fuller RW, Dollery CT. Effects of cysteinyl-leukotriene receptor antagonist ICI 204,219 on allergen-induced bronchoconstriction and airway hyperreactivity in atopic subjects. *Lancet* 1991;337:690-4.
- Findlay SR, Barden JM, Easley CB, Glass M. Effect of the oral leukotriene antagonist ICI 204,219 on the antigen-induced bronchoconstriction in subjects with asthma. *J Allergy Clin Immunol* 1992;89:1040-5.
- Israel E, Dermarkarian R, Rosenberg M, Sperling R, Taylor G, Rubin P, *et al*. The effects of a 5-lipoxygenase inhibitor on asthma induced by cold, dry air. *N Engl J Med* 1990;323:1740-4.
- Israel E, Fischer AR, Rosenberg MA, Lilly CM, Callery JC, Shapiro J, *et al*. The pivotal role of 5-lipoxygenase products in the reaction of aspirin-sensitive asthmatics to aspirin. *Am Rev Respir Dis* 1993 (in press).
- Gaddy J, Bush RK, Margolskee D, Williams VC, Busse W. The effects of a leukotriene D₄ (LTD₄) antagonist (MK-571) in mild to moderate asthma (abstract). *J Allergy Clin Immunol* 1990;85:197.
- Cloud ML, Enas GG, Kemp J, Platts-Mill T, Altman LC, Townley R, *et al*. A specific LTD₄/LTE₄ receptor antagonist improves pulmonary function in patients with mild, chronic asthma. *Am Rev Respir Dis* 1989;140:1335-9.
- Wahedna I, Wisniewski AFZ, Wong CS, Tattersfield AE. Effect of multiple doses of RG 12525, an oral leukotriene D₄ antagonist, in chronic asthma (abstract). *Am Rev Respir Dis* 1992;145:A16.
- Spector SL, Glass M, Minkwitz MC, ICI Asthma Trial Group. The effect of six weeks of therapy with oral doses of ICI 204,219 in asthmatics (abstract). *Am Rev Respir Dis* 1992;145:A16.
- Israel E, Drazen J, Pearlman H, Cohn J, Rubin P. A double blind multicenter study of Zileuton, a potent 5-lipoxygenase (5-LO) inhibitor, versus placebo in the treatment of spontaneous asthma in adults. *J Allergy Clin Immunol* 1992;89:236.
- Dahlén B, Margolskee DJ, Zetterström O, Dahlén S-E. Effect of the leukotriene receptor antagonist MK-0679 on baseline pulmonary function in aspirin sensitive asthmatic subjects. *Thorax* 1993;48:1205-10.
- Laitinen LA, Laitinen A, Haahtela T, Vilks V, Spur BW, Lee TH. Leukotriene E₄ and granulocytic infiltration into asthmatic airways. *Lancet* 1993;341:989-90.